

# Higher-spin holography from the bulk

S. Didenko (with M. A. Vasiliev)

August 3, 2017 Dubna

# Plan

- 1 Introduction
  - Locality issue
- 2 Summary of results
  - Goals
  - Results
- 3 Vasiliev system in  $d = 4$ 
  - Free level
  - Boundary conditions
  - Propagators
  - Second order
- 4 Correlators
  - Green's function
- 5 Conclusion

# *AdS/CFT* in HS

- Weak-weak duality which does not require supersymmetry (Sundborg, Klebanov, Polyakov, Leigh, Petkou, Sezgin, Sundell)

# *AdS/CFT* in HS

- Weak-weak duality which does not require supersymmetry (Sundborg, Klebanov, Polyakov, Leigh, Petkou, Sezgin, Sundell)
- Vasiliev theory is dual to vectorial CFT models

# AdS/CFT in HS

- Weak-weak duality which does not require supersymmetry (Sundborg, Klebanov, Polyakov, Leigh, Petkou, Sezgin, Sundell)
- Vasiliev theory is dual to vectorial CFT models

$$S_{HS}[\phi] \sim \langle J \dots J \rangle_{CFT}$$

# AdS/CFT in HS

- Weak-weak duality which does not require supersymmetry (Sundborg, Klebanov, Polyakov, Leigh, Petkou, Sezgin, Sundell)
- Vasiliev theory is dual to vectorial CFT models

$$S_{HS}[\phi] \sim \langle J \dots J \rangle_{CFT}$$

- Giombi and Yin tests from equations of motion: substantial piece of evidence that many of  $3pt$  functions match.

# AdS/CFT in HS

- Weak-weak duality which does not require supersymmetry (Sundborg, Klebanov, Polyakov, Leigh, Petkou, Sezgin, Sundell)
- Vasiliev theory is dual to vectorial CFT models

$$S_{HS}[\phi] \sim \langle J \dots J \rangle_{CFT}$$

- **Giombi and Yin** tests from equations of motion: substantial piece of evidence that many of  $3pt$  functions match.
- Generic structure of  $3pt$ -correlators (**Maldacena, Zhiboedov**)

$$\langle JJJ \rangle = \cos^2 \phi \langle JJJ \rangle_b + \sin^2 \phi \langle JJJ \rangle_f + \frac{1}{2} \sin(2\phi) \langle JJJ \rangle_o$$

# Locality

- **GY:** Interaction vertices are sometimes so nonlocal that produce infinities in correlation functions.



# Locality

- **GY:** Interaction vertices are sometimes so nonlocal that produce infinities in correlation functions. Some correlators can be reached via analytic continuation, but some happened to be inconsistent with crossing.

# Locality

- **GY:** Interaction vertices are sometimes so nonlocal that produce infinities in correlation functions. Some correlators can be reached via analytic continuation, but some happened to be inconsistent with crossing. Free boson and free fermion were captured.

# Locality

- **GY:** Interaction vertices are sometimes so nonlocal that produce infinities in correlation functions. Some correlators can be reached via analytic continuation, but some happened to be inconsistent with crossing. Free boson and free fermion were captured. Parity odd correlators  $\langle J_0 J_{S_1} J_{S_2} \rangle$  were accessible only in one channel.

# Locality

- **GY:** Interaction vertices are sometimes so nonlocal that produce infinities in correlation functions. Some correlators can be reached via analytic continuation, but some happened to be inconsistent with crossing. Free boson and free fermion were captured. Parity odd correlators  $\langle J_0 J_{S_1} J_{S_2} \rangle$  were accessible only in one channel.
- **Locality**

$$S = \int \phi \square \phi + D \dots D \phi D \dots D \phi D \dots D \phi + D \dots D \phi D \dots D \phi D \dots D \phi$$

# Locality

- **GY:** Interaction vertices are sometimes so nonlocal that produce infinities in correlation functions. Some correlators can be reached via analytic continuation, but some happened to be inconsistent with crossing. Free boson and free fermion were captured. Parity odd correlators  $\langle J_0 J_{S_1} J_{S_2} \rangle$  were accessible only in one channel.
- **Locality**

$$S = \int \phi \square \phi + D \dots D \phi D \dots D \phi D \dots D \phi + D \dots D \phi D \dots D \phi D \dots D \phi$$

Vasiliev equation encode HS vertices in a field-redefinition independent way through certain differential equations in a twistor space. The price to pay – one has to specify solution in a proper class of functions.

# (Non)locality

- Potential nonlocalities in Vasiliev system

# (Non)locality

- Potential nonlocalities in Vasiliev system

$$d\omega(Y|x) = -\omega * \omega + \Upsilon(\omega, \omega, C \dots C),$$

$$dC(Y|x) = -[\omega, C]_* + \Upsilon(\omega, C \dots C).$$

# (Non)locality

- Potential nonlocalities in Vasiliev system

$$d\omega(Y|x) = -\omega * \omega + \Upsilon(\omega, \omega, C \dots C),$$

$$dC(Y|x) = -[\omega, C]_* + \Upsilon(\omega, C \dots C).$$

- Origin of nonlocalities is the star-product



# (Non)locality

- Potential nonlocalities in Vasiliev system

$$d\omega(Y|x) = -\omega * \omega + \Upsilon(\omega, \omega, C \dots C),$$

$$dC(Y|x) = -[\omega, C]_* + \Upsilon(\omega, C \dots C).$$

- Origin of nonlocalities is the star-product

$$e^{y\bar{y}} * e^{-y\bar{y}} = \infty$$

# (Non)locality

- Potential nonlocalities in Vasiliev system

$$d\omega(Y|x) = -\omega * \omega + \Upsilon(\omega, \omega, C \dots C),$$

$$dC(Y|x) = -[\omega, C]_* + \Upsilon(\omega, C \dots C).$$

- Origin of nonlocalities is the star-product

$$e^{y\bar{y}} * e^{-y\bar{y}} = \infty$$

- Structure of perturbation theory

$$d_Z \text{Field}(Z, Y) = I(Z, Y)$$

# (Non)locality

- Potential nonlocalities in Vasiliev system

$$d\omega(Y|x) = -\omega * \omega + \Upsilon(\omega, \omega, C \dots C),$$

$$dC(Y|x) = -[\omega, C]_* + \Upsilon(\omega, C \dots C).$$

- Origin of nonlocalities is the star-product

$$e^{y\bar{y}} * e^{-y\bar{y}} = \infty$$

- Structure of perturbation theory

$$d_Z \text{Field}(Z, Y) = I(Z, Y), \quad \text{Field}(Z, Y) = d_Z^{-1} I(Z, Y) + H(Y)$$

# (Non)locality

- Potential nonlocalities in Vasiliev system

$$d\omega(Y|x) = -\omega * \omega + \Upsilon(\omega, \omega, C \dots C),$$

$$dC(Y|x) = -[\omega, C]_* + \Upsilon(\omega, C \dots C).$$

- Origin of nonlocalities is the star-product

$$e^{y\bar{y}} * e^{-y\bar{y}} = \infty$$

- Structure of perturbation theory

$$d_Z \text{Field}(Z, Y) = I(Z, Y), \quad \text{Field}(Z, Y) = d_Z^{-1} I(Z, Y) + H(Y)$$

If  $H(Y) = 0$  and  $d_Z^{-1}$  is homotopy operator, then  $\Upsilon(C, C)$  is nonlocal.

# (Non)locality

- Nonlocalities in  $\omega$ -sector (N. Boulanger, P. Kessel, E.D. Skvortsov, M. Taronna)

# (Non)locality

- Nonlocalities in  $\omega$ -sector (N. Boulanger, P. Kessel, E.D. Skvortsov, M. Taronna)– Infinite coefficients in front of cubic vertices within central gauge.

# (Non)locality

- Nonlocalities in  $\omega$ -sector (N. Boulanger, P. Kessel, E.D. Skvortsov, M. Taronna)– Infinite coefficients in front of cubic vertices within central gauge. Breakdown of predictivity of Vasiliev theory – *any* coefficients result from nonlocal field redefinitions.

# (Non)locality

- Nonlocalities in  $\omega$ -sector (N. Boulanger, P. Kessel, E.D. Skvortsov, M. Taronna)– Infinite coefficients in front of cubic vertices within central gauge. Breakdown of predictivity of Vasiliev theory – *any* coefficients result from nonlocal field redefinitions.
- **Vasiliev+Gelfond**: Unique field redefinition that brings equations into a local form with fixed coefficients

$$C(Y|x) + \eta \int_{[0,1]^3} e^{iu_A v^A} \delta'(1-t_1-t_2-t_3) J(t_3 u + t_1 y, v - t_2 y, \bar{y} + \bar{u}, \bar{y} + \bar{v})$$

$$J(y_1, y_2, \bar{y}_1, \bar{y}_2) = C(y_1, \bar{y}_1) C(y_2, \bar{y}_2)$$



# (Non)locality

- Nonlocalities in  $\omega$ -sector (N. Boulanger, P. Kessel, E.D. Skvortsov, M. Taronna)– Infinite coefficients in front of cubic vertices within central gauge. Breakdown of predictivity of Vasiliev theory – *any* coefficients result from nonlocal field redefinitions.
- **Vasiliev+Gelfond**: Unique field redefinition that brings equations into a local form with fixed coefficients

$$C(Y|x) + \eta \int_{[0,1]^3} e^{iu_A v^A} \delta'(1-t_1-t_2-t_3) J(t_3 u + t_1 y, v - t_2 y, \bar{y} + \bar{u}, \bar{y} + \bar{v})$$

$$J(y_1, y_2, \bar{y}_1, \bar{y}_2) = C(y_1, \bar{y}_1) C(y_2, \bar{y}_2)$$

- Fierz identities at higher orders bring in the new homotopy operator that renders equations local automatically.

# Goals and summary

## GOALS:

- Study the local form of second order higher-spin equations and their boundary limit

# Goals and summary

## GOALS:

- Study the local form of second order higher-spin equations and their boundary limit
- *AdS/CFT* driven HS reductions

# Goals and summary

## GOALS:

- Study the local form of second order higher-spin equations and their boundary limit
- $AdS/CFT$  driven HS reductions
- Calculate three-point correlation functions

# Goals and summary

## RESULTS:

- The local form of HS equations was shown to be perfectly consistent with *AdS/CFT* expectations

# Goals and summary

## RESULTS:

- The local form of HS equations was shown to be perfectly consistent with *AdS/CFT* expectations
- Known bosonic truncations of Vasiliev equations in four dimensions admit no *CFT* duals other than free theories

# Goals and summary

## RESULTS:

- The local form of HS equations was shown to be perfectly consistent with  $AdS/CFT$  expectations
- Known bosonic truncations of Vasiliev equations in four dimensions admit no  $CFT$  duals other than free theories
- Parity broken  $CFT$ 's require different truncation of the full Vasiliev system which is available at least in perturbation theory

# Goals and summary

## RESULTS:

- The local form of HS equations was shown to be perfectly consistent with *AdS/CFT* expectations
- Known bosonic truncations of Vasiliev equations in four dimensions admit no *CFT* duals other than free theories
- Parity broken *CFT*'s require different truncation of the full Vasiliev system which is available at least in perturbation theory
- Correlation functions  $\langle J_{s_1} J_{s_2} J_{s_3} \rangle$  with  $s_3 \geq s_1 + s_2$  were found including in the parity broken case



# Vasiliev equations

## Vasiliev equations in $d = 4$

$$dW + W * W = 0,$$

$$dS + [W, S]_* = 0,$$

$$dB + [W, B]_* = 0,$$

$$S * S = -i\theta_\alpha \wedge \theta^\alpha (1 + \eta B * k\mathcal{K}) - i\bar{\theta}_{\dot{\alpha}} \wedge \bar{\theta}^{\dot{\alpha}} (1 + \bar{\eta} B * \bar{k}\bar{\mathcal{K}}),$$

$$[S, B]_* = 0.$$

# Vasiliev equations

## Vasiliev equations in $d = 4$

$$dW + W * W = 0,$$

$$dS + [W, S]_* = 0,$$

$$dB + [W, B]_* = 0,$$

$$S * S = -i\theta_\alpha \wedge \theta^\alpha (1 + \eta B * k\mathcal{K}) - i\bar{\theta}_{\dot{\alpha}} \wedge \bar{\theta}^{\dot{\alpha}} (1 + \bar{\eta} B * \bar{k}\bar{\mathcal{K}}),$$

$$[S, B]_* = 0.$$

Note extra Klein operators  $k$  and  $\bar{k}$

# Field-current correspondence

Free HS equations for 0-form

$$DC(y, \bar{y}; k, \bar{k}) = 0.$$

## Field-current correspondence

Free HS equations for 0-form

$$DC(y, \bar{y}; k, \bar{k}) = 0.$$

Using Poincare connection (Vasiliev)

$$C(y, \bar{y}; k, \bar{k}) = ze^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}),$$

$$w = \sqrt{z}y, \quad \bar{w} = \sqrt{z}\bar{y}.$$

## Field-current correspondence

Free HS equations for 0-form

$$DC(y, \bar{y}; k, \bar{k}) = 0.$$

Using Poincare connection (Vasiliev)

$$C(y, \bar{y}; k, \bar{k}) = ze^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}),$$
$$w = \sqrt{z}y, \quad \bar{w} = \sqrt{\bar{z}}\bar{y}.$$

Gives current conservation

$$d_x T - \frac{i}{2} dx^{\alpha\alpha} \partial_\alpha \bar{\partial}_\alpha T = 0, \quad \partial \cdot J = 0$$

## Field-current correspondence

Free HS equations for 0-form

$$DC(y, \bar{y}; k, \bar{k}) = 0.$$

Using Poincare connection (Vasiliev)

$$C(y, \bar{y}; k, \bar{k}) = ze^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}),$$

$$w = \sqrt{z}y, \quad \bar{w} = \sqrt{\bar{z}}\bar{y}.$$

Gives current conservation

$$d_x T - \frac{i}{2} dx^{\alpha\alpha} \partial_\alpha \bar{\partial}_\alpha T = 0, \quad \partial \cdot J = 0$$

Boundary HS connections

$$d_x \omega_x = \frac{1}{4} H_{xx}^{\alpha\beta} \frac{\partial^2}{\partial w^\alpha \partial \bar{w}^\beta} (\bar{\eta} T(w, 0) \bar{k} - \eta T(0, iw) k).$$

## Boundary conditions

- Parity preserving cases  $\eta = 1$  or  $\eta = i$

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k}.$$

## Boundary conditions

- Parity preserving cases  $\eta = 1$  or  $\eta = i$

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k}.$$

- Parity broken case,  $\eta \neq 1, i$



## Boundary conditions

- Parity preserving cases  $\eta = 1$  or  $\eta = i$

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k}.$$

- Parity broken case,  $\eta \neq 1, i$

$$C(Y; k, \bar{k}) = C(Y)k + \bar{C}(Y)\bar{k}$$

## Boundary conditions

- Parity preserving cases  $\eta = 1$  or  $\eta = i$

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k}.$$

- Parity broken case,  $\eta \neq 1, i$

$$C(Y; k, \bar{k}) = C(Y)k + \bar{C}(Y)\bar{k}$$

$$C(Y) := C^+(Y), \quad \bar{C}(Y) := C^-(Y),$$

## Boundary conditions

- Parity preserving cases  $\eta = 1$  or  $\eta = i$

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k}.$$

- Parity broken case,  $\eta \neq 1, i$

$$C(Y; k, \bar{k}) = C(Y)k + \bar{C}(Y)\bar{k}$$

$$C(Y) := C^+(Y), \quad \bar{C}(Y) := C^-(Y),$$

$$\bar{\eta}T^+(w, \bar{w}) = \eta T^-(-i\bar{w}, iw).$$

# Propagators

0-form  $\Delta = 1$  propagators

$$C^+ = \eta K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \xi^\alpha y_\alpha}, \quad C^- = \bar{\eta} K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \bar{\xi}^{\dot{\alpha}} \bar{y}_{\dot{\alpha}}}$$

# Propagators

## 0-form $\Delta = 1$ propagators

$$C^+ = \eta K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \xi^\alpha y_\alpha}, \quad C^- = \bar{\eta} K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \bar{\xi}^{\dot{\alpha}} \bar{y}_{\dot{\alpha}}}$$

$$K = \frac{z}{(x - x_0)^2 + z^2},$$

$$f_{\alpha\dot{\alpha}} = -\frac{2z}{(x - x_0)^2 + z^2} (x - x_0)_{\alpha\dot{\alpha}} - i \frac{(x - x_0)^2 - z^2}{(x - x_0)^2 + z^2} \epsilon_{\alpha\dot{\alpha}},$$

$$\xi_\alpha = \Pi_{\alpha\beta} \mu_\beta, \quad \Pi_{\alpha\beta} = K \left( \frac{1}{\sqrt{z}} (x - x_0)_{\alpha\beta} - i \sqrt{z} \epsilon_{\alpha\beta} \right)$$

# Propagators

## 0-form $\Delta = 1$ propagators

$$C^+ = \eta K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \xi^\alpha y_\alpha}, \quad C^- = \bar{\eta} K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \bar{\xi}^{\dot{\alpha}} \bar{y}_{\dot{\alpha}}}$$

$$K = \frac{z}{(x - x_0)^2 + z^2},$$

$$f_{\alpha\dot{\alpha}} = -\frac{2z}{(x - x_0)^2 + z^2} (x - x_0)_{\alpha\dot{\alpha}} - i \frac{(x - x_0)^2 - z^2}{(x - x_0)^2 + z^2} \epsilon_{\alpha\dot{\alpha}},$$

$$\xi_\alpha = \Pi_{\alpha\beta} \mu_\beta, \quad \Pi_{\alpha\beta} = K \left( \frac{1}{\sqrt{z}} (x - x_0)_{\alpha\beta} - i \sqrt{z} \epsilon_{\alpha\beta} \right)$$

## $\Delta = 2$ scalar propagator

$$C_{\Delta=2} = K^2 (1 + i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}}) \times e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}}}.$$

## Boundary limit

### Current sources

$$T^+ = \frac{\eta}{|x - x_0|^2} e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha + i(x-x_0)_{\alpha\beta} \mu^\beta w^\alpha},$$

$$T^- = \frac{\bar{\eta}}{|x - x_0|^2} e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha + i(x-x_0)_{\alpha\beta} \bar{\mu}^\beta \bar{w}^\alpha}$$

## Boundary limit

### Current sources

$$T^+ = \frac{\eta}{|x - x_0|^2} e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha + i(x-x_0)_{\alpha\beta} \mu^\beta w^\alpha},$$

$$T^- = \frac{\bar{\eta}}{|x - x_0|^2} e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha + i(x-x_0)_{\alpha\beta} \bar{\mu}^\beta \bar{w}^\alpha}$$

### $\Delta = 2$ source

$$T_{\Delta=2} = \frac{w_\alpha \bar{w}^\alpha}{|x - x_0|^4} \times e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha}.$$



## Boundary limit

### Current sources

$$T^+ = \frac{\eta}{|x - x_0|^2} e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha + i(x-x_0)_{\alpha\beta} \mu^\beta w^\alpha},$$

$$T^- = \frac{\bar{\eta}}{|x - x_0|^2} e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha + i(x-x_0)_{\alpha\beta} \bar{\mu}^\beta \bar{w}^\alpha}$$

### $\Delta = 2$ source

$$T_{\Delta=2} = \frac{w_\alpha \bar{w}^\alpha}{|x - x_0|^4} \times e^{-2i(x-x_0)_{\alpha\alpha}^{-1} w^\alpha \bar{w}^\alpha}.$$

### Boundary condition

$$\bar{\eta} T^+(w, \bar{w}) = \eta T^-(-i\bar{w}, iw).$$

## Second order

Local form in the 0-form sector

$$DC = \frac{i}{2} \eta e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha} (t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k$$

$$J = CC$$

## Second order

### Local form in the 0-form sector

$$DC = \frac{i}{2} \eta e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha}(t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k$$

$$J = CC$$

### Vertices in 1-form sector are phase-independent

$$D\omega = \frac{i}{8} \eta \bar{\eta} \int \frac{1}{t_4^2} \delta(1-t_3-t_4) \delta'(1-t_1-t_2) \times$$

$$\times [\bar{H}^{\dot{\alpha}\dot{\alpha}} \left(\frac{\partial}{\partial \bar{y}^{\dot{\alpha}}}\right)^2 e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} J(t_1y, -t_2y, t_2t_4\bar{y} + t_3\bar{u}, -t_1t_4\bar{y} + \bar{v}) +$$

$$+ H^{\alpha\alpha} \left(\frac{\partial}{\partial y^{\alpha}}\right)^2 e^{iu_{\alpha}v^{\alpha}} J(t_1t_4y + t_3u, -t_2t_4y + v, t_2\bar{y}, -t_1\bar{y})],$$

## Boundary limit: currents

Carrying out boundary limit in the 0-form sector  $z \rightarrow 0$

$$C(y, \bar{y}; k, \bar{k}) = z e^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}), \quad l = TT$$

## Boundary limit: currents

Carrying out boundary limit in the 0-form sector  $z \rightarrow 0$

$$C(y, \bar{y}; k, \bar{k}) = ze^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}), \quad I = TT$$

$$\begin{aligned} d_x T - \frac{i}{2} dx^{\alpha\alpha} \partial_\alpha \bar{\partial}_\alpha T = \\ - \frac{\eta}{4} dx^{\alpha\alpha} w_\alpha \int_0^1 (t \bar{\partial}_{2\alpha} - (1-t) \bar{\partial}_{1\alpha}) I(tw, -(1-t)w, \bar{w} + i(1-t)w, \bar{w} - itw) k \end{aligned}$$

## Boundary limit: currents

Carrying out boundary limit in the 0-form sector  $z \rightarrow 0$

$$C(y, \bar{y}; k, \bar{k}) = z e^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}), \quad I = TT$$

$$\begin{aligned} d_x T - \frac{i}{2} dx^{\alpha\alpha} \partial_\alpha \bar{\partial}_\alpha T = \\ - \frac{\eta}{4} dx^{\alpha\alpha} w_\alpha \int_0^1 (t \bar{\partial}_{2\alpha} - (1-t) \bar{\partial}_{1\alpha}) I(tw, -(1-t)w, \bar{w} + i(1-t)w, \bar{w} - itw) k \end{aligned}$$

HS currents do not conserve  $\partial \cdot J \neq 0$

## Boundary limit: currents

Carrying out boundary limit in the 0-form sector  $z \rightarrow 0$

$$C(y, \bar{y}; k, \bar{k}) = ze^{y_\alpha \bar{y}^\alpha} T(w, \bar{w}; k, \bar{k}), \quad I = TT$$

$$\begin{aligned} d_x T - \frac{i}{2} dx^{\alpha\alpha} \partial_\alpha \bar{\partial}_\alpha T = \\ - \frac{\eta}{4} dx^{\alpha\alpha} w_\alpha \int_0^1 (t \bar{\partial}_{2\alpha} - (1-t) \bar{\partial}_{1\alpha}) I(tw, -(1-t)w, \bar{w} + i(1-t)w, \bar{w} - itw) k \end{aligned}$$

HS currents do not conserve  $\partial \cdot J \neq 0$

They do for the free theories boundary conditions

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k} \quad \Rightarrow \quad \partial \cdot J = 0$$

## Boundary limit: HS connections

Boundary limit in the 1-form sector  $z \rightarrow 0$

$$\begin{aligned}
 D_x \omega_x &= \frac{i}{8} \eta \bar{\eta} \int d^2 t \delta'(1 - t_1 - t_2) H_{xx}^{\alpha\alpha} \left( \frac{\partial}{\partial u^\alpha} \right)^2 \times \\
 &\times \left\{ I(t_1(w + u), -t_2(w + u), it_2 w, -it_1 w) - \right. \\
 &\left. - I(t_1 w, -t_2 w, it_2(w + u), -it_1(w + u)) \right\} \Big|_{u=0}
 \end{aligned}$$



## Boundary limit: HS connections

Boundary limit in the 1-form sector  $z \rightarrow 0$

$$D_x \omega_x = \frac{i}{8} \eta \bar{\eta} \int d^2 t \delta'(1 - t_1 - t_2) H_{xx}^{\alpha\alpha} \left( \frac{\partial}{\partial u^\alpha} \right)^2 \times \\
 \times \left\{ I(t_1(w+u), -t_2(w+u), it_2 w, -it_1 w) - \right. \\
 \left. - I(t_1 w, -t_2 w, it_2(w+u), -it_1(w+u)) \right\} \Big|_{u=0}$$

Again, for free theory boundary conditions

$$T(w, \bar{w})k = \pm T(-i\bar{w}, iw)\bar{k} \quad \Rightarrow D_x \omega_x = 0$$

## Boundary limit: HS connections

Boundary limit in the 1-form sector  $z \rightarrow 0$

$$D_x \omega_x = \frac{i}{8} \eta \bar{\eta} \int d^2 t \delta'(1 - t_1 - t_2) H_{xx}^{\alpha\alpha} \left( \frac{\partial}{\partial u^\alpha} \right)^2 \times \\
 \times \left\{ I(t_1(w+u), -t_2(w+u), it_2 w, -it_1 w) - \right. \\
 \left. - I(t_1 w, -t_2 w, it_2(w+u), -it_1(w+u)) \right\} \Big|_{u=0}$$

Again, for free theory boundary conditions

$$T(w, \bar{w}) k = \pm T(-i\bar{w}, iw) \bar{k} \quad \Rightarrow \quad D_x \omega_x = 0$$

For parity broken theory  $\eta \neq 1, i$

$$D_x \omega_x \neq 0$$

# Green's function

- **Giombi-Yin:** Correlators from equations of motion

$$DC = J[C, C], \quad \langle JJJ \rangle \sim \lim_{z \rightarrow 0} z^{-1} G(wz^{-\frac{1}{2}}, \bar{w}z^{-\frac{1}{2}}) \Big|_{\bar{w}=0}$$

# Green's function

- **Giombi-Yin:** Correlators from equations of motion

$$DC = J[C, C], \quad \langle JJJ \rangle \sim \lim_{z \rightarrow 0} z^{-1} G(wz^{-\frac{1}{2}}, \bar{w}z^{-\frac{1}{2}}) \Big|_{\bar{w}=0}$$

- $G(w, \bar{w}|x, z)$  – is the Green's function,  $w_\alpha$  – to be associated with outgoing spinor polarization

## Green's function

- **Giombi-Yin:** Correlators from equations of motion

$$DC = J[C, C], \quad \langle JJJ \rangle \sim \lim_{z \rightarrow 0} z^{-1} G(wz^{-\frac{1}{2}}, \bar{w}z^{-\frac{1}{2}}) \Big|_{\bar{w}=0}$$

- $G(w, \bar{w}|x, z)$  – is the Green's function,  $w_\alpha$  – to be associated with outgoing spinor polarization

$$C^{2s}(y|x, z) \rightarrow z^{s+1} \int \frac{dz_0 d^3x_0}{z_0^4} G(y, \partial_{y_0}|x - x_0, z_0) J(y_0|x_0, z_0)$$

## Green's function

- **Giombi-Yin:** Correlators from equations of motion

$$DC = J[C, C], \quad \langle JJJ \rangle \sim \lim_{z \rightarrow 0} z^{-1} G(wz^{-\frac{1}{2}}, \bar{w}z^{-\frac{1}{2}}) \Big|_{\bar{w}=0}$$

- $G(w, \bar{w}|x, z)$  – is the Green's function,  $w_\alpha$  – to be associated with outgoing spinor polarization

$$C^{2s}(y|x, z) \rightarrow z^{s+1} \int \frac{dz_0 d^3x_0}{z_0^4} G(y, \partial_{y_0}|x - x_0, z_0) J(y_0|x_0, z_0)$$

$$G_s(y, \lambda|x, z) \sim \int_0^\infty \frac{dt}{(1+t)^{-2s}} \left\{ \frac{(y\lambda\theta)^{2s}}{(2s)!} \left[ \Gamma[2-2s] x^{2s-3} \sin(2(s-1) \arctan \frac{x}{z}) \right] \right\} \Big|_{z \rightarrow (2t+1)z}$$

## Green's function

- **Giombi-Yin:** Correlators from equations of motion

$$DC = J[C, C], \quad \langle JJJ \rangle \sim \lim_{z \rightarrow 0} z^{-1} G(wz^{-\frac{1}{2}}, \bar{w}z^{-\frac{1}{2}}) \Big|_{\bar{w}=0}$$

- $G(w, \bar{w}|x, z)$  – is the Green's function,  $w_\alpha$  – to be associated with outgoing spinor polarization

$$C^{2s}(y|x, z) \rightarrow z^{s+1} \int \frac{dz_0 d^3x_0}{z_0^4} G(y, \partial_{y_0}|x - x_0, z_0) J(y_0|x_0, z_0)$$

$$G_s(y, \lambda|x, z) \sim \int_0^\infty \frac{dt}{(1+t)^{-2s}} \left\{ \frac{(y\lambda\theta)^{2s}}{(2s)!} \left[ \Gamma[2-2s] x^{2s-3} \sin(2(s-1) \arctan \frac{x}{z}) \right] \right\} \Big|_{z \rightarrow (2t+1)z}$$

**GY Green's function does not take into account consistency constraint on  $J[C, C]$**

# Homotopy Green's function

Equation to solve

$$DG = \frac{i}{2} e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha}(t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k.$$



# Homotopy Green's function

Equation to solve

$$DG = \frac{i}{2} e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha} (t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k.$$

$$J = CC, \quad DC = 0$$

# Homotopy Green's function

Equation to solve

$$DG = \frac{i}{2} e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha} (t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k.$$

$$J = CC, \quad DC = 0$$

We're looking for solution outside a spin triangle

$$s \geq s_1 + s_2$$

# Homotopy Green's function

Equation to solve

$$DG = \frac{i}{2} e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha} (t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k.$$

$$J = CC, \quad DC = 0$$

We're looking for solution outside a spin triangle

$$s \geq s_1 + s_2$$

The Green's function:

$$G = \frac{1}{2} \int_{[0,1]^3 \times \mathbf{R}} \delta'(1-t_1-t_2-t_3) e^{iu_{AV}^A} J(u+t_1y, t_3v-t_2y, \bar{y}+\bar{u}, \bar{y}+\bar{v}) k$$

## Homotopy Green's function

Equation to solve

$$DG = \frac{i}{2} e^{\alpha\dot{\alpha}} \int e^{i\bar{u}_{\dot{\alpha}}\bar{v}^{\dot{\alpha}}} y_{\alpha} (t\bar{u}_{\dot{\alpha}} + (1-t)\bar{v}_{\dot{\alpha}}) J(ty, -(1-t)y, \bar{y} + \bar{u}, \bar{y} + \bar{v}) k.$$

$$J = CC, \quad DC = 0$$

We're looking for solution outside a spin triangle

$$s \geq s_1 + s_2$$

The Green's function:

$$G = \frac{1}{2} \int_{[0,1]^3 \times \mathbf{R}} \delta'(1-t_1-t_2-t_3) e^{iu_A v^A} J(u+t_1y, t_3v-t_2y, \bar{y}+\bar{u}, \bar{y}+\bar{v}) k$$

Valid for opposite helicity signs only!

## 3pt functions

- Recall that propagators get naturally split into positive and negative helicity parts

$$C^+ = \eta K e^{if_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i\xi^\alpha y_\alpha}, \quad C^- = \bar{\eta} K e^{if_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i\bar{\xi}^{\dot{\alpha}} \bar{y}_{\dot{\alpha}}}$$

## 3pt functions

- Recall that propagators get naturally split into positive and negative helicity parts

$$C^+ = \eta K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \xi^\alpha y_\alpha}, \quad C^- = \bar{\eta} K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \bar{\xi}^{\dot{\alpha}} \bar{y}_{\dot{\alpha}}}$$

- all different three-point correlators arrange in

$$\begin{aligned} \langle JJJ \rangle_{boson} &\sim G^{++} + G^{--} + G^{+-} + G^{-+}, \\ \langle JJJ \rangle_{fermion} &\sim G^{++} + G^{--} - G^{+-} - G^{-+}, \\ \langle JJJ \rangle_{odd} &\sim G^{++} - G^{--}, \end{aligned}$$

## 3pt functions

- Recall that propagators get naturally split into positive and negative helicity parts

$$C^+ = \eta K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \xi^\alpha y_\alpha}, \quad C^- = \bar{\eta} K e^{i f_{\alpha\dot{\alpha}} y^\alpha \bar{y}^{\dot{\alpha}} + i \bar{\xi}^{\dot{\alpha}} \bar{y}_{\dot{\alpha}}}$$

- all different three-point correlators arrange in

$$\begin{aligned} \langle JJJ \rangle_{boson} &\sim G^{++} + G^{--} + G^{+-} + G^{-+}, \\ \langle JJJ \rangle_{fermion} &\sim G^{++} + G^{--} - G^{+-} - G^{-+}, \\ \langle JJJ \rangle_{odd} &\sim G^{++} - G^{--}, \end{aligned}$$

- Simple Gaussian integration gives

$$G^{+-} = \int d^3 t \frac{K_1 K_2}{\Delta} \delta'(1-t_1-t_2-t_3) e^{2 \frac{t_1 t_2}{\Delta} Q + \frac{t_1}{\Delta} ((1-t_3) P_1 + z t_3 \tilde{S}_1) - \frac{t_1}{\Delta} ((1-t_3) P_2 + z t_3 \tilde{S}_2)}$$

## 3pt functions

$$\begin{aligned}
 G^{++} &= \frac{z}{2} K'_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (-\tau P_2 + S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{--} &= \frac{z}{2} K'_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 - S_1)^{2s_1} (-\tau P_2 - S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{+-} &= \frac{z}{2} K_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (-\tau P_2 - S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{-+} &= \frac{z}{2} K_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 - S_1)^{2s_1} (-\tau P_2 + S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}.
 \end{aligned}$$



## 3pt functions

$$\begin{aligned}
 G^{++} &= \frac{z}{2} K'_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (-\tau P_2 + S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{--} &= \frac{z}{2} K'_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 - S_1)^{2s_1} (-\tau P_2 - S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{+-} &= \frac{z}{2} K_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (-\tau P_2 - S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{-+} &= \frac{z}{2} K_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 - S_1)^{2s_1} (-\tau P_2 + S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}.
 \end{aligned}$$

$$K_{s_1 s_2 s} = \frac{2^{s-s_1-s_2} (s+s_1+s_2)!}{(2s)!(2s_1)!(2s_2)!}, \quad K'_{s_1 s_2 s} = \frac{2^{s-s_1-s_2} (s+s_1-s_2)!(s+s_2-s_1)!}{(2s)!(2s_1)!(2s_2)!(s-s_1-s_2)!}$$

## 3pt functions

$$\begin{aligned}
 G^{++} &= \frac{z}{2} K'_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (-\tau P_2 + S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{--} &= \frac{z}{2} K'_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 - S_1)^{2s_1} (-\tau P_2 - S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{+-} &= \frac{z}{2} K_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (-\tau P_2 - S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}, \\
 G^{-+} &= \frac{z}{2} K_{s_1 s_2 s} \frac{Q^{s-s_1-s_2}}{|x_{01}||x_{02}||x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s} (\tau P_1 - S_1)^{2s_1} (-\tau P_2 + S_2)^{2s_2}}{(1+\tau^2)^{s+s_1+s_2+1}}.
 \end{aligned}$$

$$K_{s_1 s_2 s} = \frac{2^{s-s_1-s_2} (s+s_1+s_2)!}{(2s)!(2s_1)!(2s_2)!}, \quad K'_{s_1 s_2 s} = \frac{2^{s-s_1-s_2} (s+s_1-s_2)!(s+s_2-s_1)!}{(2s)!(2s_1)!(2s_2)!(s-s_1-s_2)!}$$

$$\begin{aligned}
 P_1 &= i \frac{(x_{01})_{\alpha\alpha} \mu_0^\alpha \mu_1^\alpha}{|x_{01}|^2}, & P_2 &= i \frac{(x_{02})_{\alpha\alpha} \mu_0^\alpha \mu_2^\alpha}{|x_{02}|^2}; & Q &= \left( \frac{x_{01}}{|x_{01}|^2} - \frac{x_{02}}{|x_{02}|^2} \right)_{\alpha\alpha} \mu_0^\alpha \mu_0^\alpha, \\
 S_1 &= \frac{(x_{02})^{\beta\alpha} (x_{12})_{\alpha\gamma} \mu_{1\gamma} \mu_{0\beta}}{|x_{01}||x_{02}||x_{12}|}, & S_2 &= \frac{(x_{01})^{\beta\alpha} (x_{12})_{\alpha\gamma} \mu_{2\gamma} \mu_{0\beta}}{|x_{01}||x_{02}||x_{12}|}.
 \end{aligned}$$

## 3pt functions

- Free theory correlators

$$G^{++} + G^{--} = \frac{K'_{s_1 s_2 s} Q^{s-s_1-s_2}}{|x_{01}| |x_{02}| |x_{12}|} \int_{-\infty}^{\infty} d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (\tau P_2 + S_2)^{2s_2}}{(1 + \tau^2)^{s+s_1+s_2+1}}$$

$$G^{+-} + G^{-+} = \frac{K_{s_1 s_2 s} Q^{s-s_1-s_2}}{|x_{01}| |x_{02}| |x_{12}|} \int_{-\infty}^{\infty} d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (\tau P_2 - S_2)^{2s_2}}{(1 + \tau^2)^{s+s_1+s_2+1}}$$

## 3pt functions

- Free theory correlators

$$G^{++} + G^{--} = \frac{K'_{s_1 s_2 s} Q^{s-s_1-s_2}}{|x_{01}| |x_{02}| |x_{12}|} \int_{-\infty}^{\infty} d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (\tau P_2 + S_2)^{2s_2}}{(1 + \tau^2)^{s+s_1+s_2+1}}$$

$$G^{+-} + G^{-+} = \frac{K_{s_1 s_2 s} Q^{s-s_1-s_2}}{|x_{01}| |x_{02}| |x_{12}|} \int_{-\infty}^{\infty} d\tau \frac{\tau^{2s} (\tau P_1 + S_1)^{2s_1} (\tau P_2 - S_2)^{2s_2}}{(1 + \tau^2)^{s+s_1+s_2+1}}$$

Match with free boson and free fermion

## 3pt functions

- Parity broken correlators  $\eta \neq 1, i$

$$\langle J_{s_1} J_{s_2} J_s \rangle_{odd} \sim \frac{1}{2} \frac{K'_{s_1 s_2 s}}{|x_{01}| |x_{02}| |x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s}}{(1 + \tau^2)^{s+s_1+s_2+1}} Q^{s-s_1-s_2} \times \\ ((\tau P_1 + S_1)^{2s_1} (\tau P_2 - S_2)^{2s_2} - (\tau P_1 - S_1)^{2s_1} (\tau P_2 + S_2)^{2s_2})$$

## 3pt functions

- Parity broken correlators  $\eta \neq 1, i$

$$\langle J_{s_1} J_{s_2} J_s \rangle_{\text{odd}} \sim \frac{1}{2} \frac{K'_{s_1 s_2 s}}{|x_{01}| |x_{02}| |x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s}}{(1 + \tau^2)^{s+s_1+s_2+1}} Q^{s-s_1-s_2} \times \\ ((\tau P_1 + S_1)^{2s_1} (\tau P_2 - S_2)^{2s_2} - (\tau P_1 - S_1)^{2s_1} (\tau P_2 + S_2)^{2s_2})$$

$$\langle J_{s_1} J_{s_2} J_s \rangle \sim \int_0^\infty d\tau \frac{\tau^{2s}}{(1 + \tau^2)^{s+s_1+s_2+1}} (\tau a + b)^{2s_1} (\tau c + d)^{2s_2}$$

## 3pt functions

- Parity broken correlators  $\eta \neq 1, i$

$$\langle J_{s_1} J_{s_2} J_s \rangle_{\text{odd}} \sim \frac{1}{2} \frac{K'_{s_1 s_2 s}}{|x_{01}| |x_{02}| |x_{12}|} \int_0^\infty d\tau \frac{\tau^{2s}}{(1 + \tau^2)^{s+s_1+s_2+1}} Q^{s-s_1-s_2} \times \\ ((\tau P_1 + S_1)^{2s_1} (\tau P_2 - S_2)^{2s_2} - (\tau P_1 - S_1)^{2s_1} (\tau P_2 + S_2)^{2s_2})$$

$$\langle J_{s_1} J_{s_2} J_s \rangle \sim \int_0^\infty d\tau \frac{\tau^{2s}}{(1 + \tau^2)^{s+s_1+s_2+1}} (\tau a + b)^{2s_1} (\tau c + d)^{2s_2}$$

$$\int_0^{\pi/2} d\phi \sin^{2s} \phi \sin^{2s_1}(\phi + \phi_1) \sin^{2s_2}(\phi + \phi_2), \quad \tau = \tan \phi$$

## Examples

- Quasi-bosonic scalar (S. Giombi 1, V. Gurucharan, V. Kirilin, S. Prakash, E.D. Skvortsov)

$$\langle J_1 J_0 J_3 \rangle \sim Q_3^2 S_2,$$

$$\langle J_2 J_0 J_4 \rangle \sim Q_3^2 S_2 (4P_2^2 + Q_1 Q_3),$$

$$\langle J_1 J_0 J_5 \rangle \sim Q_3^4 S_2,$$

$$\langle J_3 J_0 J_5 \rangle \sim Q_3^2 S_2 (2P_2^2 + Q_1 Q_3) (6P_2^2 + Q_1 Q_3),$$

$$\langle J_2 J_0 J_6 \rangle \sim Q_3^4 S_2 (6P_2^2 + Q_1 Q_3),$$

$$\langle J_4 J_0 J_6 \rangle \sim Q_3^2 S_2 (107P_2^4 Q_1 Q_3 + 40P_2^2 Q_1^2 Q_3^2 + 102P_2^6 + 3Q_1^3 Q_3^3)$$



## Examples

- Quasi-bosonic scalar (S. Giombi 1, V. Gurucharan, V. Kirilin, S. Prakash, E.D. Skvortsov)

$$\langle J_1 J_0 J_3 \rangle \sim Q_3^2 S_2,$$

$$\langle J_2 J_0 J_4 \rangle \sim Q_3^2 S_2 (4P_2^2 + Q_1 Q_3),$$

$$\langle J_1 J_0 J_5 \rangle \sim Q_3^4 S_2,$$

$$\langle J_3 J_0 J_5 \rangle \sim Q_3^2 S_2 (2P_2^2 + Q_1 Q_3) (6P_2^2 + Q_1 Q_3),$$

$$\langle J_2 J_0 J_6 \rangle \sim Q_3^4 S_2 (6P_2^2 + Q_1 Q_3),$$

$$\langle J_4 J_0 J_6 \rangle \sim Q_3^2 S_2 (107P_2^4 Q_1 Q_3 + 40P_2^2 Q_1^2 Q_3^2 + 102P_2^6 + 3Q_1^3 Q_3^3)$$

All match.

## Conclusion

- Vasiliev equations are analyzed in the current interaction sector  $s \geq s_1 + s_2$  to the second order. Perfect agreement with *CFT* expectation is found.
- Boundary limit of equations was investigated. It is shown that known bosonic HS system has no CFT dual other than free theories. Parity broken CFTs result from different truncation of Vasiliev equations. Boundary conditions require nonlinear modifications at higher orders.
- Homotopy Green's function in the current interaction sector is found.
- Parity broken three-point functions  $\langle J_{s_1} J_{s_2} J_s \rangle$  were extracted from equations of motion.