

# Standard Model at High Energies

**N. A. Gromov**

Institute of Physics and Mathematics, Komi Science Center RAS,  
E-mail: [gromov@dm.komisc.ru](mailto:gromov@dm.komisc.ru)

**Supersymmetries and Quantum Symmetries**  
**Dubna, Russia, July 31 – August 5, 2017**

# The Standard Model

- ▶ The modern knowledge of the particle world is concentrated in Standard Model (SM). This theory consist of two parts: Electroweak Model (EWM), which unified electromagnetic and weak interactions, and Quantum Chromodynamics (QCD), describing their strong interactions.
- ▶ The Electroweak Model is a gauge theory with the gauge group  $SU(2) \times U(1)$ , which act in the boson, lepton and quark sectors.
- ▶ QCD is  $SU(3)$  gauge theory based on the local color degrees of freedom of quarks.

# Elementary particles content of SM

Gauge bosons:

$\gamma$  (photon),

$W^\pm$  (charged weak bosons),  $Z^0$  (neutral weak boson),

$A^k$ ,  $k = 1, \dots, 8$  (gluons).

Special particle:

$\chi$  (Higgs boson).

Leptons:

$$\begin{pmatrix} \nu_e \\ e \end{pmatrix}, \quad \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix}, \quad \begin{pmatrix} \nu_\tau \\ \tau \end{pmatrix} \in \mathbb{C}_2.$$

Quarks:

$$\begin{pmatrix} u \\ d \end{pmatrix}, \quad \begin{pmatrix} c \\ s \end{pmatrix}, \quad \begin{pmatrix} t \\ b \end{pmatrix} \in \mathbb{C}_2.$$

## Electroweak Model

The Lagrangian of the model is given by the sum of the boson  $L_B$ , of the lepton  $L_L$  and of the quark  $L_Q$  Lagrangians:

$$L = L_B + L_L + L_Q.$$

It is invariant under the action of the gauge group  $SU(2) \times U(1)$  in the 2-dim. complex space  $\mathbb{C}_2$ :

$$SU(2) : \vec{z}' = G\vec{z},$$

$$\begin{pmatrix} z'_1 \\ z'_2 \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}, \quad |\alpha|^2 + |\beta|^2 = 1,$$

$$U(1) : \vec{z}' = e^{i\omega} \vec{z}, \quad \omega \in \mathbf{R}.$$

**Boson sector**  $L_B = L_A + L_\phi$  involve two parts:  
**the gauge field Lagrangian**

$$L_A = -\frac{1}{4}[(F_{\mu\nu}^1)^2 + (F_{\mu\nu}^2)^2 + (F_{\mu\nu}^3)^2] - \frac{1}{4}(B_{\mu\nu})^2,$$

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu], \quad B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$$

and **the matter field Lagrangian**

$$L_\phi = \frac{1}{2}(D_\mu\phi)^\dagger D_\mu\phi - \frac{\lambda}{4}(\phi^\dagger\phi - v^2)^2,$$

where  $\phi = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} \in \mathbb{C}_2$  are the matter fields.

**Fermion** sector is represented by **the lepton**  $L_L$  and **quark**  $L_Q$  Lagrangians.

For the first generation **the lepton** Lagrangian is taken in the form:

$$L_{L,e} = L_l^\dagger i \tilde{\tau}_\mu D_\mu L_l + e_r^\dagger i \tau_\mu D_\mu e_r - h_e [e_r^\dagger (\phi^\dagger L_l) + (L_l^\dagger \phi) e_r],$$

where  $L_l = \begin{pmatrix} \nu_l \\ e_l \end{pmatrix} \in \mathbb{C}_2$  is the  $SU(2)$ -doublet,  $e_r$  is the  $SU(2)$ -singlet,  $h_e$  is constant,  $\tau_0 = \tilde{\tau}_0 = \mathbf{1}$ ,  $\tilde{\tau}_k = -\tau_k$ ,  $\tau_\mu$  are Pauli matrices and  $e_r, e_l, \nu_l$  are the two component Lorentz spinors. First and second terms in  $L_{L,e}$  describe free movement of left and right fermions and their interactions with gauge fields. Last term corresponds to the electron mass.

The quark Lagrangian is given by

$$L_Q = Q_l^\dagger i \tilde{\tau}_\mu D_\mu Q_l + u_r^\dagger i \tau_\mu D_\mu u_r + d_r^\dagger i \tau_\mu D_\mu d_r - \\ - h_d [d_r^\dagger (\phi^\dagger Q_l) + (Q_l^\dagger \phi) d_r] - h_u [u_r^\dagger (\tilde{\phi}^\dagger Q_l) + (Q_l^\dagger \tilde{\phi}) u_r],$$

where left quark fields form the  $SU(2)$ -doublet

$Q_l = \begin{pmatrix} u_l \\ d_l \end{pmatrix} \in \mathbb{C}_2$ , right quark fields  $u_r, d_r$  are the

$SU(2)$ -singlets,  $\tilde{\phi}_i = \epsilon_{ik} \bar{\phi}_k$ ,  $\epsilon_{00} = 1$ ,  $\epsilon_{ii} = -1$  is the conjugate representation of  $SU(2)$  group and  $h_u, h_d$  are constants. All fields  $u_l, d_l, u_r, d_r$  are two component Lorentz spinors.

## The new gauge fields

$$W_{\mu}^{\pm} = \frac{1}{\sqrt{2}} (A_{\mu}^1 \mp iA_{\mu}^2), \quad Z_{\mu} = \frac{1}{\sqrt{g^2 + g'^2}} (gA_{\mu}^3 - g'B_{\mu}),$$

$$A_{\mu} = \frac{1}{\sqrt{g^2 + g'^2}} (g'A_{\mu}^3 + gB_{\mu})$$

are expressed through the old fields

$$A_{\mu}(x) = -ig \sum_{k=1}^3 T_k A_{\mu}^k(x), \quad B_{\mu}(x) = -ig' B_{\mu}(x),$$

and take their values in the Lie algebras  $su(2)$  and  $u(1)$ , respectively.



# Chromodynamics

- ▶ QCD is a gauge theory based on the local color degrees of freedom. The QCD gauge group is  $SU(3)$ , acting in 3-dime. complex space  $\mathbb{C}_3$  of color quark states

$$q = \begin{pmatrix} q_1 \\ q_2 \\ q_3 \end{pmatrix} \equiv \begin{pmatrix} q_R \\ q_G \\ q_B \end{pmatrix} \in \mathbb{C}_3,$$

where  $q(x)$  are quark fields  $q = u, d, s, c, b, t$  and  $R$  (red),  $G$  (green),  $B$  (blue) are color degrees of freedom.

- ▶ The  $SU(3)$  gauge bosons are called **gluons**. There are eight gluons in total, which are the force carrier of the theory between quarks.

- ▶ QCD Lagrangian is taken in the form

$$\mathcal{L} = \sum_q \bar{q}^i (i\gamma^\mu) (D_\mu)_{ij} q^j - \frac{1}{4} \sum_{\alpha=1}^8 F_{\mu\nu}^\alpha F^{\mu\nu\alpha},$$

- ▶ where  $D_\mu q$  are covariant derivatives of quark fields

$$D_\mu q = \left( \partial_\mu - ig_s \left( \frac{\lambda^a}{2} \right) A_\mu^a \right) q,$$

$g_s$  is the strong coupling constant,  $t^a = \lambda^a/2$  are generators of  $SU(3)$ ,  $\lambda^a$  — Gell-Mann matrices

- ▶ and gluon stress tensor has the form

$$F_{\mu\nu}^\alpha = \partial_\mu A_\nu^\alpha - \partial_\nu A_\mu^\alpha + g_s f^{\alpha\beta\gamma} A_\mu^\beta A_\nu^\gamma.$$

## Contraction of the Electroweak Model

- ▶ The **infinite energy limit** of EWM corresponds to the **consistent rescaling** of  $SU(2)$  and  $\mathbb{C}_2$

$$\begin{pmatrix} z'_1 \\ \epsilon z'_2 \end{pmatrix} = \begin{pmatrix} \alpha & \epsilon\beta \\ -\epsilon\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} z_1 \\ \epsilon z_2 \end{pmatrix}, \quad |\alpha|^2 + \epsilon^2|\beta|^2 = 1,$$

when contraction parameter tends to zero  $\epsilon \rightarrow 0$ .

- ▶ Substitution  $\beta \rightarrow \epsilon\beta$  induces another ones for the gauge fields:

$$W_\mu^\pm \rightarrow \epsilon W_\mu^\pm, \quad Z_\mu \rightarrow Z_\mu, \quad A_\mu \rightarrow A_\mu.$$

and substitution  $z_2 \rightarrow \epsilon z_2$  induces the following transformation of lepton and quark fields:

$$e_l \rightarrow \epsilon e_l, \quad d_l \rightarrow \epsilon d_l, \quad \nu_l \rightarrow \nu_l, \quad u_l \rightarrow u_l.$$

- ▶ The next reason is the special mechanism of spontaneous symmetry breaking, which is used to generate mass for the vector bosons. One of  $L_B$  ground states

$$\phi^{vac} = \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad A_\mu^k = B_\mu = 0$$

is taken as vacuum and then small field excitations  $v + \chi(x)$  with respect to this vacuum are regarded.

- ▶ So, Higgs boson field  $\chi$ , constant  $v$  and particle masses  $m_p$ , which depend on  $v$ , are multiplied by contraction parameter:

$$\chi \rightarrow \epsilon\chi, \quad v \rightarrow \epsilon v, \quad m_p \rightarrow \epsilon m_p, \quad p = \chi, W, Z, e, u, d.$$

- ▶ After these transformations the Lagrangian can be represented in the form

$$L(\epsilon) = L_\infty - \epsilon m_u (u_r^\dagger u_l + u_l^\dagger u_r) + \\ + \epsilon^2 L_2 + \epsilon^3 L_3 + \epsilon^4 L_4.$$

For  $\epsilon \rightarrow 0$  the terms with the higher powers of  $\epsilon$  contribute less than the terms with lower powers.

- ▶ So the Electroweak Model demonstrates **five stages of behaviour** in the infinite energy limit, which are distinguished by the powers of the contraction parameter.

## QCD with contracted gauge group

The contracted group  $SU(3; \epsilon)$  is defined by the action

$$q'(\epsilon) = \begin{pmatrix} q'_1 \\ \epsilon q'_2 \\ \epsilon^2 q'_3 \end{pmatrix} = \begin{pmatrix} u_{11} & \epsilon u_{12} & \epsilon^2 u_{13} \\ \epsilon u_{21} & u_{22} & \epsilon u_{23} \\ \epsilon^2 u_{31} & \epsilon u_{32} & u_{33} \end{pmatrix} \begin{pmatrix} q_1 \\ \epsilon q_2 \\ \epsilon^2 q_3 \end{pmatrix} = U(\epsilon)q(\epsilon)$$

on the color space  $\mathbb{C}_3(\epsilon)$ .

The quark and gluon fields are transformed as follows:

$$q_1 \rightarrow q_1, \quad q_2 \rightarrow \epsilon q_2, \quad q_3 \rightarrow \epsilon^2 q_3,$$

$$A_\mu^{GR} \rightarrow \epsilon A_\mu^{GR}, \quad A_\mu^{BG} \rightarrow \epsilon A_\mu^{BG}, \quad A_\mu^{BR} \rightarrow \epsilon^2 A_\mu^{BR},$$

and diagonal gauge fields are not changed

$$A_\mu^{RR} \rightarrow A_\mu^{RR}, \quad A_\mu^{GG} \rightarrow A_\mu^{GG}, \quad A_\mu^{BB} \rightarrow A_\mu^{BB}.$$

- ▶ With this substitution, we obtain the quark part of QCD Lagrangian in the form

$$\begin{aligned}
 \mathcal{L}_q(\kappa) = & \sum_q i\bar{q}_1\gamma^\mu\partial_\mu q_1 + \frac{g_s}{2}|q_1|^2\gamma^\mu A_\mu^{RR} + \\
 & + \epsilon^2 \left\{ i\bar{q}_2\gamma^\mu\partial_\mu q_2 + \frac{g_s}{2} \left( |q_2|^2\gamma^\mu A_\mu^{GG} + q_1\bar{q}_2\gamma^\mu A_\mu^{GR} + \bar{q}_1q_2\gamma^\mu \bar{A}_\mu^{GR} \right) \right\} + \\
 & + \epsilon^4 \left[ i\bar{q}_3\gamma^\mu\partial_\mu q_3 + \frac{g_s}{2} \left( |q_3|^2\gamma^\mu A_\mu^{BB} + q_1\bar{q}_3\gamma^\mu A_\mu^{BR} + \bar{q}_1q_3\gamma^\mu \bar{A}_\mu^{BR} + \right. \right. \\
 & \left. \left. + q_2\bar{q}_3\gamma^\mu A_\mu^{BG} + \bar{q}_2q_3\gamma^\mu \bar{A}_\mu^{BG} \right) \right] = L_q^\infty + \epsilon^2 L_q^{(2)} + \epsilon^4 L_q^{(4)}.
 \end{aligned}$$

- ▶ Gluon part  $L_{gl} = -\frac{1}{4}F_{\mu\nu}^\alpha F^{\mu\nu\alpha}$  of Lagrangian is very cumbersome, therefore we omit its general form.

- ▶ The contracted QCD Lagrangian has the form

$$\mathcal{L}(\epsilon) = L^\infty + \epsilon^2 L^{(2)} + \epsilon^4 L^{(4)} + \epsilon^6 L^{(6)} + \epsilon^8 L^{(8)},$$

with the explicit expressions for each  $L^{(k)}$ .

- ▶ The contraction parameter is monotonous function of the average energy  $E$  (or temperature  $T$ ) with the property  $\epsilon(E) \rightarrow 0$  for  $E \rightarrow \infty$ .



Very higher energies (temperatures) can exist in the early Universe after inflation and reheating on the first stages of the Hot Big Bang ( $1eV = 10^4 K$ ).




Рис.: History of the Universe (V. Rubakov, D. Gorbunov, INR RAS)

- ▶ When the contraction of some mathematical or physical structure is performed **one can reconstruct the initial structure** moving back along the contraction way.
- ▶ Similar approach as applied to the Standard Model gives a chance to describe some part of the very early history of the Universe.

## Estimation of boundary values

- ▶ The contraction of gauge groups of SM gives an opportunity to order in time different stages of its development, but does not make it possible to bear their absolute date.
- ▶ To estimate the absolute date we use additional assumptions. The equality of the contraction parameters for QCD and the EWM is one of these assumptions.
- ▶ Then we use the fact that the electroweak epoch starts at the temperature  $T_4 = 100 \text{ GeV}$  ( $1 \text{ GeV} = 10^{13} \text{ K}$ ) and the QCD epoch begins at  $T_8 = 0,2 \text{ GeV}$ , i.e. we assume that complete reconstruction of the EWM, whose Lagrangian has minimal terms  $\approx \epsilon^4$ , and QCD with minimal terms  $\approx \epsilon^8$ , take place at these temperatures.

- ▶ Let us denote by  $\Delta$  cutoff level for  $\epsilon^k$ ,  $k = 1, 2, 4, 6, 8$ , i.e. for  $\epsilon^k < \Delta$  all the terms proportionate to  $\epsilon^k$  are negligible quantities in Lagrangian.
- ▶ At last we suppose that the contraction parameter inversely depends on temperature

$$\epsilon(T) = \frac{A}{T}, \quad A = \text{const.} \quad (1)$$

- ▶ From the equation for QCD  $\epsilon^8(T_8) = A^8 T_8^{-8} = \Delta$  we obtain  $A = T_8 \Delta^{1/8} = 0,2 \Delta^{1/8} \text{ GeV}$ .

From the similar equation for EWM we obtain the cutoff level  $\Delta = (T_8 E_4^{-1})^8 = (0,2 \cdot 10^{-2})^8 \approx 10^{-22}$ .

- ▶ From the equation for  $k$ -th power  $\epsilon^k(T_k) = A^k T_k^{-k} = \Delta$  we obtain

$$T_k = T_8 \Delta^{\frac{k-8}{8k}} \approx 10^{\frac{88-15k}{4k}} \text{ GeV}$$

and easily find (GeV):  $T_1 = 10^{18}$ ,  $T_2 = 10^7$ ,  $T_3 = 10^3$ ,  $T_4 = 10^2$ ,  $T_6 = 1$ ,  $T_8 = 2 \cdot 10^{-1}$ .

- ▶ **The obtained estimation for "infinity" temperature  $T_1 \approx 10^{18} \text{ GeV}$  is comparable with Planck energy  $\approx 10^{19} \text{ GeV}$ , where the gravitation effects are important.**
- ▶ So the developed evolution of the elementary particles does not exceed the range of the problems described by electroweak and strong interactions.

## Evolution of elementary particles

$$(T > 10^{18} \text{ GeV}).$$

- ▶ We can give some conclusions already at the level of classical fields. At the infinite temperature ( $\epsilon = 0$ ) the EWM Lagrangian is as follows

$$L_\infty = -\frac{1}{4}Z_{\mu\nu}^2 - \frac{1}{4}F_{\mu\nu}^2 + \nu_l^\dagger i\tilde{\tau}_\mu \partial_\mu \nu_l + u_l^\dagger i\tilde{\tau}_\mu \partial_\mu u_l + \\ + e_r^\dagger i\tau_\mu \partial_\mu e_r + d_r^\dagger i\tau_\mu \partial_\mu d_r + u_r^\dagger i\tau_\mu \partial_\mu u_r + L_\infty^{int}(A_\mu, Z_\mu).$$

- ▶ So the Electroweak Model includes only **massless particles**: photons  $A_\mu$  and neutral bosons  $Z_\mu$ , left quarks  $u_l$  and neutrinos  $\nu_l$ , right electrons  $e_r$  and quarks  $u_r, d_r$ .
- ▶ **The electroweak interactions become long-range** because they are mediated by the massless  $Z$ -bosons and photons.

- ▶ From the explicit form of the interaction part

$$\begin{aligned}
 L_{\infty}^{int}(A_{\mu}, Z_{\mu}) = & \frac{g}{2 \cos \theta_w} \nu_l^{\dagger} \tilde{\tau}_{\mu} Z_{\mu} \nu_l + \frac{2e}{3} u_l^{\dagger} \tilde{\tau}_{\mu} A_{\mu} u_l + \\
 & + \frac{g}{\cos \theta_w} \left( \frac{1}{2} - \frac{2}{3} \sin^2 \theta_w \right) u_l^{\dagger} \tilde{\tau}_{\mu} Z_{\mu} u_l + g' \sin \theta_w e_r^{\dagger} \tau_{\mu} Z_{\mu} e_r - \\
 & - g' \cos \theta_w e_r^{\dagger} \tau_{\mu} A_{\mu} e_r - \frac{1}{3} g' \cos \theta_w d_r^{\dagger} \tau_{\mu} A_{\mu} d_r + \frac{1}{3} g' \sin \theta_w d_r^{\dagger} \tau_{\mu} Z_{\mu} d_r + \\
 & + \frac{2}{3} g' \cos \theta_w u_r^{\dagger} \tau_{\mu} A_{\mu} u_r - \frac{2}{3} g' \sin \theta_w u_r^{\dagger} \tau_{\mu} Z_{\mu} u_r
 \end{aligned}$$

it follows that **there are no interactions between particles of different kind**, for example neutrinos interact only with each other by neutral currents.

- ▶ It looks like some **stratification of the Electroweak Model** with only one sort of particles in each stratum.

- ▶ It follows from the limit QCD Lagrangian

$$\mathcal{L}_\infty = L_q^\infty + L_{gl}^\infty = \sum_q i\bar{q}_1 \gamma^\mu \partial_\mu q_1 + \frac{g_s}{2} |q_1|^2 \gamma^\mu A_\mu^{RR} - \frac{1}{4} (F_{\mu\nu}^{RR})^2 - \frac{1}{4} (F_{\mu\nu}^{GG})^2 - \frac{1}{4} F_{\mu\nu}^{RR} F_{\mu\nu}^{GG}.$$

that only dynamic terms for the first color components of massless quarks survive under infinite temperature, which means that **quarks are monochromatic**.

- ▶ The terms also survive, which describe the interactions of these components with  $R$ -gluons. So **the stratification is conserved in the QCD sector**.



## Evolution of elementary particles

$$(\approx \epsilon, \quad 10^{18} \text{ GeV} \geq T > 10^7 \text{ GeV}).$$

- ▶ The mass term of  $u$ -quark in the complete Lagrangian  $L(\epsilon)$  is proportional to  $\epsilon$

$$\epsilon m_u (u_r^\dagger u_l + u_l^\dagger u_r).$$

The same is held for  $c$ - and  $t$ -quark. So  $u$ -,  $c$ - and  $t$ -quark first restores its mass in the evolution of the Universe.

## Evolution of elementary particles

$$(\approx \epsilon^2, \quad 10^7 \text{ GeV} \geq T > 10^3 \text{ GeV}).$$

- ▶ The mass terms of electron and  $d$ -quark are multiplied by  $\epsilon^2$

$$\epsilon^2 \left[ m_e (e_r^\dagger e_l + e_l^\dagger e_r) + m_d (d_r^\dagger d_l + d_l^\dagger d_r) \right],$$

The same is true for  $\mu^-$  and  $\tau$ -lepton, for  $s$ - and  $b$ -quark.  
**These particles become massive in the second stage.**

- ▶ The quarks obtain the second color degree of freedom.
- ▶ The main part of electroweak and color interactions are restored in this epoch.

## Evolution of elementary particles

$$(\approx \epsilon^3, \quad 10^3 \text{ GeV} \geq T > 10^2 \text{ GeV}).$$

There is one term in Lagrangian  $L_3 = gW_\mu^+ W_\mu^- \chi$  proportionate to  $\epsilon^3$ , which describe Higgs boson interaction with charged  $W$ -bosons.

$$(\approx \epsilon^4, \quad 10^2 \text{ GeV} \geq T > 1 \text{ GeV}).$$

- ▶ Higgs boson  $\chi$  and charged  $W$ -boson last restore their masses after all other particles of SM.
- ▶ The final reconstruction of the EWM takes place in this epoch.
- ▶ The quarks obtain the third color degree of freedom.

## Evolution of elementary particles

$$(\approx \epsilon^6, \quad 1 \text{ GeV} \geq T > 2 \cdot 10^{-1} \text{ GeV}).$$

Next part of color interactions is restored.

$$(\approx \epsilon^8, \quad T \leq 2 \cdot 10^{-1} \text{ GeV}).$$

- ▶ The color interactions are completely valid.
- ▶ **The QCD is fully reconstructed** .
- ▶ **Start the time of the Standard Model.**

## Conclusion

The hypothesis of the particles evolution in the early Universe starting from Planck temperature  $10^{19} GeV$  was offered, using the modern knowledge of the particle world, which is concentrated in SM.

The particles evolution was developed from the first principles of the gauge theory with the help of the contractions of the SM gauge groups.

- ▶ more details:

**Journal of Cosmology and Astroparticle  
Physics 03(2016) 053.**

- ▶ **Thank you for attention.**