

# *Hyperbolic spin Ruijsenaars-Schneider model from Poisson reduction*

*Gleb Arutyunov*

*II. Institute for Theoretical Physics  
Hamburg University*

*“Supersymmetries and Quantum Symmetries SQS’19” 26-31.08.2019*

*Yerevan*

# Calogero-Moser-Sutherland models

## Rational model

Invert oscillator potential

$$H = \frac{1}{2}p^2 + \frac{1}{2}\omega^2 q^2 \rightarrow H = \frac{1}{2}p^2 + \frac{\varkappa^2}{q^2}$$

Integrable generalisation to many particles  $\{q\}_{i=1,\dots,N}$

$$H = \frac{1}{2} \sum_{i=1}^N p_i^2 + \varkappa^2 \sum_{i<j}^N \frac{1}{q_{ij}^2}, \quad q_{ij} = q_i - q_j$$

## Hyperbolic model

$$H = \frac{1}{2} \sum_{i=1}^N p_i^2 + \varkappa^2 \sum_{i<j}^N \frac{1}{\sinh q_{ij}^2}$$

The models were discovered in 1970's.  
Wide applications

- soliton theory
- quantum field theory
- solvable models of stat. mechanics
- black hole physics
- condensed matter
- quantum chaos
- representation theory
- harmonic analysis
- random matrix theory
- complex geometry

## Elliptic model

$$H = \frac{1}{2} \sum_{i=1}^N p_i^2 + \varkappa^2 \sum_{i<j}^N \wp(q_{ij})$$

# Ruijsenaars-Schneider models

## Rational model

$$H = c^2 \sum_{i=1}^N \cosh \frac{p_i}{c} \prod_{i \neq j}^N \sqrt{1 + \frac{\kappa^2}{c^2 q_{ij}^2}}$$

## Hyperbolic model

$$H = c^2 \sum_{i=1}^N \cosh \frac{p_i}{c} \prod_{i \neq j}^N \sqrt{1 + \frac{\kappa^2}{c^2 \sinh^2 q_{ij}^2}}$$

## Elliptic model

$$H = c^2 \sum_{i=1}^N \cosh \frac{p_i}{c} \prod_{i \neq j}^N \sqrt{\lambda + \mu \wp(q_{ij})}$$

$$\{p_i, q_j\} = \delta_{ij}$$

Expanding in the limit  $c \rightarrow \infty$  the corresponding Hamiltonians of the CMS models are recovered


What about spin models?

# Outline

- *Spin RS models: equations of motion*
- *Heisenberg double*
- *Oscillator manifold*
- *Poisson-Lie group action on a product manifold*
- *Reduction*
- *Superintegrability*
- *Conclusions and future directions*

*with Enrico Olivucci, arXiv:1906.02619*

# Equations of motion of the spin RS model



$$V(z) = \zeta(z) - \zeta(z + \gamma)$$

$$\dot{q}_i = f_{ii}$$

$$\dot{\mathbf{a}}_{i\alpha} = \sum_{j \neq i}^N V(q_{ij}) (\mathbf{a}_{j\alpha} - \mathbf{a}_{i\alpha})$$

$$\dot{\mathbf{c}}_{i\alpha} = \sum_{j \neq i}^N \left( V(q_{ij}) f_{ij} \mathbf{c}_{i\alpha} - V(q_{ji}) f_{ji} \mathbf{c}_{j\alpha} \right)$$

*Krichever & Zabrodin, 1995*

$$f_{ij} = \sum_{\alpha=1}^{\ell} \mathbf{a}_{i\alpha} \mathbf{c}_{\alpha j}$$

$$\sum_{\alpha=1}^{\ell} \mathbf{a}_{i\alpha} = 1 \quad \forall \alpha$$

*collective spin variables*

# Hamiltonian structure in the rational case

$$L_{ij} = \frac{f_{ij}}{q_{ij} + \gamma}$$

$$\{q_i, q_j\} = 0, \quad \{q_i, \mathbf{a}_{i\alpha}\} = 0, \quad \{q_i, \mathbf{c}_{j\alpha}\} = \delta_{ij} \mathbf{c}_{j\alpha},$$

$$\{\mathbf{a}_{i\alpha}, \mathbf{a}_{j\beta}\} = \frac{\delta_{i \neq j}}{q_{ij}} (\mathbf{a}_{i\alpha} \mathbf{a}_{j\beta} + \mathbf{a}_{i\beta} \mathbf{a}_{j\alpha} - \mathbf{a}_{i\alpha} \mathbf{a}_{i\beta} - \mathbf{a}_{j\alpha} \mathbf{a}_{j\beta})$$

$$\{\mathbf{a}_{i\alpha}, \mathbf{c}_{\beta j}\} = \mathbf{a}_{i\alpha} L_{ij} - \delta_{\alpha\beta} L_{ij} - \frac{\delta_{i \neq j}}{q_{ij}} (\mathbf{a}_{i\alpha} - \mathbf{a}_{j\alpha}) \mathbf{c}_{\beta j},$$

$$\{\mathbf{c}_{\alpha i}, \mathbf{c}_{\beta j}\} = \frac{\delta_{i \neq j}}{q_{ij}} (\mathbf{c}_{i\alpha} \mathbf{c}_{\beta j} + \mathbf{c}_{\beta i} \mathbf{c}_{\alpha j}) - \mathbf{c}_{\alpha i} L_{ij} + \mathbf{c}_{\beta j} L_{ji}$$

G.A. & Frolov, 1997

Hamiltonian reduction

$$\mathcal{M} = T^*G \times \Sigma, \quad \Sigma = \underbrace{\mathcal{O} \times \mathcal{O} \times \dots \times \mathcal{O}}_{\ell}$$

$\mathcal{O}$  - coadjoint orbit of minimal dimension


$$G : \mathcal{M} \rightarrow \mathcal{M} \quad \Longrightarrow \quad \mu : \mathcal{M} \rightarrow \mathfrak{g}^*$$

$$\mathcal{P} = \mu^{-1}(\gamma \mathbb{1}) / G$$

quasi – Hamiltonian reduction

*Chalykh & Fairon, 2018*

quasi-Poisson (Van der Bergh's bracket)



Jordan quiver/ $G$

On the other hand, there is a deformation hierarchy of initial phase spaces

$$T^*G \longrightarrow D_+(G)$$

Heisenberg double



*Gorsky & Nekrasov, 1994*

*G.A. & Frolov, 1996*

....

*Feher & Klimcik, 2009*

...

$$\mathcal{M} = D_+(G) \times ???$$



What should be there in the spin case?

# Heisenberg double

$(\mathfrak{g}, \mathfrak{g}^*)$  - factorisable Lie bialgebra,  $\mathfrak{g}^* \simeq \mathfrak{g}$

$$\mathcal{D} = \mathfrak{g} \oplus \mathfrak{g} \longleftarrow \text{double}$$

$$(X, X) \subset \mathcal{D}, \quad \forall X \in \mathfrak{g}$$

$$(X_+, X_-) = (\hat{\varepsilon}_+ X, \hat{\varepsilon}_- X) \subset \mathcal{D}, \quad \forall X \in \mathfrak{g}^* \simeq \mathfrak{g}$$

$\hat{\varepsilon}_\pm = \hat{\varepsilon} \pm \frac{1}{2}\mathbb{1}$  are two linear operators,  $\hat{\varepsilon}_\pm : \mathfrak{g}^* \rightarrow \mathfrak{g}_\pm \subset \mathfrak{g}$   
 $\varepsilon \in \mathfrak{g} \wedge \mathfrak{g}$  split solution of mCYBE

$$D = G \times G \longleftarrow \text{double Lie group}$$

$$G^* \simeq (u_+, u_-) \subset D$$

diffeomorphism  $\sigma : G^* \simeq G$

$$\sigma(u_+, u_-) = u_+ u_-^{-1} = u$$



# Heisenberg double

$D_+(G)$

$A, B \in G = \text{GL}_N(\mathbb{C})$

$$\begin{aligned}\frac{1}{\varkappa}\{A_1, A_2\} &= -\tau_- A_1 A_2 - A_1 A_2 \tau_+ + A_1 \tau_- A_2 + A_2 \tau_+ A_1, \\ \frac{1}{\varkappa}\{B_1, B_2\} &= -\tau_- B_1 B_2 - B_1 B_2 \tau_+ + B_1 \tau_- B_2 + B_2 \tau_+ B_1, \\ \frac{1}{\varkappa}\{A_1, B_2\} &= -\tau_- A_1 B_2 - A_1 B_2 \tau_- + A_1 \tau_- B_2 + B_2 \tau_+ A_1, \\ \frac{1}{\varkappa}\{B_1, A_2\} &= -\tau_+ B_1 A_2 - B_1 A_2 \tau_+ + B_1 \tau_- A_2 + A_2 \tau_+ B_1.\end{aligned}$$

$$\tau_{\pm} = \pm \frac{1}{2} \sum_{i=1}^N E_{ii} \otimes E_{ii} \pm \sum_{i \leq j}^N E_{ij} \otimes E_{ji}$$

$$\tau_+ - \tau_- = C_{12} = \sum_{i,j=1}^N E_{ij} \otimes E_{ji}$$

$$\tau = \frac{1}{2}(\tau_+ + \tau_-)$$

Poisson action of a Poisson-Lie group  $G$

$$A \rightarrow hAh^{-1}, \quad B \rightarrow hBh^{-1}, \quad h \in G$$

The Poisson-Lie structure of  $G$  is given in terms of the Sklyanin bracket

$$\{h_1, h_2\} = -\varkappa [\varkappa_{\pm}, h_1 h_2], \quad h \in G.$$

The non-abelian moment map for this action  $(\mathcal{M}_+, \mathcal{M}_-)$

$$m = m_+ m_-^{-1} \in G \quad \longrightarrow \quad \mathcal{M} = BA^{-1}B^{-1}A$$

$$\frac{1}{\varkappa} \{m_1, m_2\} = -\varkappa_+ m_1 m_2 - m_1 m_2 \varkappa_- + m_1 \varkappa_- m_2 + m_2 \varkappa_+ m_1$$

 Semenov-Tian-Shansky bracket

Involutive family  $\{H_k, H_m\} = 0$

$$H_k = \text{Tr}(BA^{-1})^k = \text{Tr}(A^{-1}B)^k, \quad k \in \mathbb{Z}$$

$$\underline{\Sigma_{N,\ell}} : \quad a_{i\alpha} \equiv (a)_{i\alpha}, \quad b_{\alpha j} \equiv (b)_{\alpha j} \quad i = 1, \dots, N, \quad \alpha = 1, \dots, \ell$$

$$\{a_1, a_2\}_{\pm} = \varkappa (\tau a_1 a_2 \mp a_1 a_2 \rho),$$

$$\{b_1, b_2\}_{\pm} = \varkappa (b_1 b_2 \tau \mp \rho b_1 b_2),$$

$$\{a_1, b_2\}_{\pm} = \varkappa (-b_2 \tau_+ a_1 \pm a_1 \rho \mp b_2) - C_{12}^{\text{rec}},$$

$$\{b_1, a_2\}_{\pm} = \varkappa (-b_1 \tau_- a_2 \pm a_2 \rho \pm b_1) + C_{21}^{\text{rec}}.$$

$$C_{12}^{\text{rec}} = \sum_{i=1}^N \sum_{\alpha=1}^{\ell} E_{i\alpha} \otimes E_{\alpha i}$$

$$\rho_{\pm} = \pm \frac{1}{2} \sum_{\alpha=1}^{\ell} E_{\alpha\alpha} \otimes E_{\alpha\alpha} \pm \sum_{\alpha \lesssim \beta}^{\ell} E_{\alpha\beta} \otimes E_{\beta\alpha}$$

$$\rho_+ - \rho_- = C_{12}^{\text{s}} = \sum_{\alpha, \beta=1}^{\ell} E_{\alpha\beta} \otimes E_{\beta\alpha}$$

$$\rho = \frac{1}{2}(\rho_+ + \rho_-)$$

$$\varkappa = 0$$

$$\{a_{i\alpha}, b_{\beta j}\} = -\delta_{ij} \delta_{\alpha\beta}$$

$N\ell$  pairs of canonically conjugate variables

# Oscillator manifold

$$\omega = \mathbb{1} + \varkappa ab$$

Define the following action of the Poisson-Lie group  $G$  on oscillators

$$\delta_X a_{i\alpha} = (\text{Ad}_\omega^* X a)_{i\alpha} \quad \delta_X b_{\alpha i} = -(b \text{Ad}_\omega^* X)_{\alpha i}, \quad X \in \mathfrak{g}$$

$\text{Ad}_g^* X$  for  $g \equiv (g_+, g_-) \in G^*$  is the dressing transformation

★ This action is Poisson

★ If  $\omega = \omega_+ \omega_-^{-1}$  then  $(\omega_+^{-1}, \omega_-^{-1}) \in G^*$  is the moment map

$$n = \omega_+^{-1} \omega_- \in G$$

$$\frac{1}{\varkappa} \{n_1, n_2\} = -\varkappa_+ n_1 n_2 - n_1 n_2 \varkappa_- + n_1 \varkappa_- n_2 + n_2 \varkappa_+ n_1$$

← Semenov-Tian-Shansky bracket

# Poisson-Lie action on a product manifold

Let  $\mathcal{M}_1$  and  $\mathcal{M}_2$  be two Poisson manifolds with brackets  $\{\cdot, \cdot\}_{\mathcal{M}_1}$  and  $\{\cdot, \cdot\}_{\mathcal{M}_2}$


$$m_i : \mathcal{M}_i \rightarrow G^*$$

$$\mathcal{M} = \mathcal{M}_1 \times \mathcal{M}_2$$

$$m = m_1 m_2 \quad \longrightarrow \quad G : \mathcal{M} \rightarrow \mathcal{M}$$

$$\xi_X f = \langle X, \{m, f\}_{\mathcal{M}} m^{-1} \rangle, \quad f \in \text{Fun}(\mathcal{M})$$

$$X \rightarrow \xi_X \quad \text{Lie algebra homomorphism}$$


$$\mathcal{M} = D_+(G) \times \Sigma_{N,\ell}^{\pm}$$

# Moment map equation

$$\mathcal{M}, \mathcal{N} \in G$$

Product in  $G^*$

$$\mathcal{M} \star \mathcal{N} = q \mathbb{1}$$

Moment map for  
the action of  $G$   
on the double

Moment map for  
the action of  $G$   
on the oscillator manifold

$$\mathcal{M} = q \omega_+ \omega_-^{-1} = q \omega$$


$$BA^{-1}B^{-1}A = q(\mathbb{1} + \varkappa ab)$$

# Reduction

$$\mathcal{P} = \{\text{Solutions of } BA^{-1}B^{-1}A = q(\mathbb{1} + \varkappa ab)\}/G$$

$$\delta_X a_{i\alpha} = (\text{Ad}_{\omega \star m^{-1}}^* X a)_{i\alpha} \quad \delta_X b_{\alpha i} = -(b \text{Ad}_{\omega \star m^{-1}}^* X)_{\alpha i}, \quad X \in \mathfrak{g},$$

$$\omega \star m^{-1} = \omega_+ m_+^{-1} m_- \omega_-^{-1} \equiv q^{-1} \mathbb{1}$$



$$a_{i\alpha} \longrightarrow (h a)_{i\alpha} \quad b_{\alpha i} \longrightarrow (b h^{-1})_{\alpha i}, \quad h = e^X \in G$$

Construction of  $G$ -invariants becomes elementary !

# Reduction

$$A = TQT^{-1}, \quad B = UP^{-1}T^{-1}$$

G.A. & Frolov, 1996

diagonal

$$\sum_{j=1}^N T_{ij} = \sum_{j=1}^N U_{ij} = 1, \quad \forall i = 1, \dots, N$$

Frobenius

$$t \text{ diagonal} \rightarrow t_{ij} = \delta_{ij} \sum_{\alpha=1}^{\ell} (T^{-1}a)_{i\alpha}$$

$$\underline{L = t^{-1}T^{-1}UP^{-1}tQ^{-1},}$$

Lax matrix

$$\underline{\mathbf{a} = t^{-1}T^{-1}a,}$$

$$\underline{\mathbf{c} = bA^{-1}BTt}$$

Invariant spins

$$L - qQ^{-1}LQ = q\kappa \mathbf{ac}.$$

$\implies$

$$L = q\kappa \sum_{i,j=1}^N \frac{Q_i}{Q_i - qQ_j} (\mathbf{ac})_{ij} E_{ij}$$



# Poisson algebra

$$Z = Q^{-1}LQ$$

$$\{Q_i, \mathbf{a}_{j\alpha}\} = 0, \quad \{Q_i, \mathbf{c}_{\alpha j}\} = \delta_{ij} \mathbf{c}_{\alpha j} Q_j$$

$$\{\mathbf{a}_1, \mathbf{a}_2\}_{\pm} = \varkappa[(r^{\bullet} \mp Y) \mathbf{a}_1 \mathbf{a}_2 \mp \mathbf{a}_1 \mathbf{a}_2 \rho \mp \mathbf{a}_1 X_{21} \mathbf{a}_2 \pm \mathbf{a}_2 X_{12} \mathbf{a}_1],$$

$$\{\mathbf{a}_1, \mathbf{c}_2\}_{\pm} = \varkappa[\mathbf{c}_2(r_{12}^* \pm Y) \mathbf{a}_1 \pm \mathbf{a}_1 \rho \mp \mathbf{c}_2 \pm \mathbf{a}_1 \mathbf{c}_2 X_{21} \mp X_{12}^{\mp} \mathbf{a}_1 \mathbf{c}_2] + K_{21} \mathbf{a}_1 Z_2 - C_{12}^{\text{rec}} Z_2,$$

$$\{\mathbf{c}_1, \mathbf{a}_2\}_{\pm} = \varkappa[\mathbf{c}_1(-r_{21}^* \pm Y) \mathbf{a}_2 \pm \mathbf{a}_2 \rho \pm \mathbf{c}_1 \mp \mathbf{a}_2 \mathbf{c}_1 X_{12} \pm X_{21}^{\mp} \mathbf{a}_2 \mathbf{c}_1] - K_{12} \mathbf{a}_2 Z_1 + C_{21}^{\text{rec}} Z_1,$$

$$\{\mathbf{c}_1, \mathbf{c}_2\}_{\pm} = \varkappa[\mathbf{c}_1 \mathbf{c}_2 (r^{\circ} \mp Y) \mp \rho \mathbf{c}_1 \mathbf{c}_2 \pm \mathbf{c}_1 X_{12}^{\mp} \mathbf{c}_2 \mp \mathbf{c}_2 X_{21}^{\mp} \mathbf{c}_1] + \mathbf{c}_2 K_{12} Z_1 - \mathbf{c}_1 K_{21} Z_2,$$

$$X_{12} = \sum_{i\beta\sigma\delta} (\mathbf{a}_1 \rho)_{i\beta\sigma\delta} E_{ii} \otimes E_{\sigma\delta},$$

$$X_{12}^{\pm} = \sum_{i\beta\sigma\delta} (\mathbf{a}_1 \rho^{\pm})_{i\beta\sigma\delta} E_{ii} \otimes E_{\sigma\delta},$$

$$K_{12} = \sum_{i\sigma} E_{\sigma i} \otimes E_{ii},$$

$$Y_{12} = \sum_{i\beta k\delta} (\mathbf{a}_1 \mathbf{a}_2 \rho)_{i\beta k\delta} E_{ii} \otimes E_{kk}.$$

$$r^{\bullet} = \frac{1}{2} \sum_{i,j=1}^N \frac{Q_i + Q_j}{Q_i - Q_j} (E_{ii} - E_{ij}) \otimes (E_{jj} - E_{ji}),$$

$$r^* = \frac{1}{2} \sum_{i,j=1}^N \frac{Q_i + Q_j}{Q_i - Q_j} (E_{ij} - E_{ii}) \otimes E_{jj}, \quad r^{\circ} = \frac{1}{2} \sum_{i,j=1}^N \frac{Q_i + Q_j}{Q_i - Q_j} (E_{ii} \otimes E_{jj} - E_{ij} \otimes E_{ji}).$$

# Poisson algebra

$$\frac{1}{\kappa} \{L_1, L_2\}_{\pm} = (r_{12} \mp Y)L_1L_2 - L_1L_2(\underline{r}_{12} \pm Y) + L_1(\bar{r}_{21} \pm Y)L_2 - L_2(\bar{r}_{12} \mp Y)L_1$$

$$r = \sum_{i \neq j}^N \left( \frac{Q_j}{Q_{ij}} E_{ii} - \frac{Q_i}{Q_{ij}} E_{ij} \right) \otimes (E_{jj} - E_{ji}),$$

$$\bar{r} = \sum_{i \neq j}^N \frac{Q_i}{Q_{ij}} (E_{ii} - E_{ij}) \otimes E_{jj}, \quad \underline{r} = \sum_{i \neq j}^N \frac{Q_i}{Q_{ij}} (E_{ij} \otimes E_{ji} - E_{ii} \otimes E_{jj});$$

$$Y = \sum_{i\beta j\delta} (\mathbf{a}_1 \mathbf{a}_2 \rho)_{i\beta j\delta} E_{ii} \otimes E_{jj}$$

The  $L$ -algebra is not the same as in the spin case!

$$\frac{1}{\kappa} \{L_1, L_2\} = r_{12}L_1L_2 - L_1L_2\underline{r}_{12} + L_1\bar{r}_{21}L_2 - L_2\bar{r}_{12}L_1$$

# Poisson algebra

$$H_m = \text{Tr}(BA^{-1})^m \quad \longleftarrow \text{commutative family}$$

$$J_n^+ = \text{Tr}[S(BA^{-1})^n], \quad J_n^- = \text{Tr}[S(A^{-1}B)^n]$$

$$\{H_m, I_n\} = \{H_m, J_n\} = 0$$

$$J_n^{+\alpha\beta} = \text{Tr}[S^{\alpha\beta}(BA^{-1})^n], \quad J_n^{-\alpha\beta} = \text{Tr}[S^{\alpha\beta}(A^{-1}B)^n]$$

$$(S^{\alpha\beta})_{ij} = a_{i\alpha} b_{\beta j}$$

$$J_n^{+\alpha\beta} = \text{Tr}[S^{\alpha\beta} Q^{-1} L^{-1} Q L^n], \quad J_n^{-\alpha\beta} = \text{Tr}[S^{\alpha\beta} Q^{-1} L^{n-1} Q]$$

$$(S^{\alpha\beta})_{ij} = \mathbf{a}_{i\alpha} \mathbf{c}_{\beta j}$$


$$J_0^{+\alpha\beta} = J_0^{-\alpha\beta} = \text{Tr} S^{\alpha\beta}$$

# Poisson algebra

$$\begin{aligned}
 \frac{1}{\kappa} \{J_n^{\alpha\beta}, J_m^{\gamma\delta}\} &= \frac{1}{\kappa} (\delta^{\beta\gamma} J_{n+m}^{\alpha\delta} - \delta^{\alpha\delta} J_{n+m}^{\gamma\beta}) \\
 &\pm \left[ \rho_{\alpha\mu, \gamma\nu} J_n^{\mu\beta} J_m^{\nu\delta} + J_n^{\alpha\mu} J_m^{\gamma\nu} \rho_{\mu\beta, \nu\delta} - J_m^{\gamma\nu} \rho_{\pm\alpha\mu, \nu\delta} J_n^{\mu\beta} - J_n^{\alpha\mu} \rho_{\mp\mu\beta, \gamma\nu} J_m^{\nu\delta} \right] \\
 &\pm \left[ -\frac{1}{2} (J_n^{\alpha\delta} J_m^{\gamma\beta} - J_m^{\alpha\delta} J_n^{\gamma\beta}) + \sum_{p=0}^m (J_{n+m-p}^{\alpha\delta} J_p^{\gamma\beta} - J_{m-p}^{\alpha\delta} J_{n+p}^{\gamma\beta}) \right] \\
 &+ \frac{1 \mp 1}{2} (J_{n+m}^{\alpha\delta} J_0^{\gamma\beta} - J_0^{\alpha\delta} J_{n+m}^{\gamma\beta}).
 \end{aligned}$$

$J_n^{\pm\alpha\beta}$

$\Sigma_{N, \ell}^{\pm}$




# Poisson algebra of the spin group

$$\{J_0^{\alpha\beta}, J_0^{\gamma\delta}\} = \delta^{\beta\gamma} J_0^{\alpha\delta} - \delta^{\alpha\delta} J_0^{\gamma\beta} \\ \pm \kappa \left[ \rho_{\alpha\mu, \nu\rho} J_0^{\mu\beta} J_0^{\nu\delta} + J_0^{\alpha\mu} J_0^{\gamma\nu} \rho_{\mu\beta, \nu\delta} - J_0^{\gamma\nu} \rho_{\pm\alpha\mu, \nu\delta} J_0^{\mu\beta} - J_0^{\alpha\mu} \rho_{\mp\mu\beta, \gamma\nu} J_0^{\nu\delta} \right]$$

$$\omega^{\mu\nu} = \delta^{\mu\nu} + \kappa J_0^{\alpha\beta}$$

$$\{\omega_1, \omega_2\}_{\pm} = \pm(\rho \omega_1 \omega_2 + \omega_1 \omega_2 \rho - \omega_2 \rho_{\pm} \omega_1 - \omega_1 \rho_{\mp} \omega_2)$$



Semenov-Tian-Shansky bracket

$\omega$   $\longleftarrow$  moment map for the Poisson action of the spin Poisson-Lie group  $S = \text{GL}_{\ell}(\mathbb{C})$

$$a_{i\alpha} \longrightarrow (ag)_{i\alpha}, \quad b_{\alpha i} \longrightarrow (g^{-1}b)_{\alpha i}, \quad g \in S$$

$$\{g_1, g_2\} = \pm \kappa [\rho, g_1 g_2]$$

# Poisson algebra

$$J(\lambda) = \sum_{n=0}^{\infty} J_n^+ \lambda^{-n-1}$$

$$\begin{aligned} \{J_1(\lambda), J_2(\mu)\}_{\pm} &= \frac{1}{\lambda - \mu} [C_{12}^s, J_1(\lambda) + J_2(\mu)] \\ &\pm \varkappa \left[ \rho_{\pm}(\lambda, \mu) J_1(\lambda) J_2(\mu) + J_1(\lambda) J_2(\mu) \rho_{\mp}(\lambda, \mu) - J_2(\mu) \rho_{\pm} J_1(\lambda) - J_1(\lambda) \rho_{\mp} J_2(\mu) \right] \end{aligned}$$

$$\rho_{\pm}(\lambda, \mu) = \rho \pm \frac{1}{2} \frac{\lambda + \mu}{\lambda - \mu} C_{12}^s = \frac{\lambda \rho_{\pm} \mp \mu \rho_{\mp}}{\lambda - \mu}$$

# Solving equation of motion

$$\underline{H_1} \longrightarrow \dot{A} = -B, \quad \dot{B} = -BA^{-1}B, \quad \dot{a} = 0 = \dot{b}$$

$BA^{-1} = I$  is an integral of motion and also  $a = \text{const}$ ,  $b = \text{const}$

$$A(\tau) = e^{-I\tau} A(0), \quad B(\tau) = I e^{-I\tau} A(0)$$

$$\underline{\text{Initial data}} \quad A(0) \equiv Q, \quad a_{i\alpha}(0) \equiv a_{i\alpha}, \quad \sum a_{i\alpha} = 1 \quad \forall i$$

$$I = L(0)$$

$$e^{-L(0)\tau} Q = T(\tau) Q(\tau) T(\tau)^{-1}$$

 Frobenius,  $T(\tau) = \mathbb{1}$

$$\mathbf{a}_{i\alpha}(\tau) = \frac{T(\tau)_{ij}^{-1} a_{j\alpha}}{\sum_{\beta} T(\tau)_{ij}^{-1} a_{j\beta}} = T(\tau)_{ij}^{-1} a_{j\alpha}$$

# Conclusions

The Hamiltonian structure of the hyperbolic spin RS model is found from the Poisson reduction of  $D_+(G) \times \Sigma_{N,\ell}^\pm$

- ★ Elliptic version?
- ★ Quantum model?