

Radiatively Induced Breaking of Conformal Symmetry in the Standard Model

Andrej Arbuzov

JINR, Dubna

In collaboration with: V.N. Pervushin, R.G. Nazmitdinov,
A.E. Pavlov, and A.F. Zakharov

arXiv:1411.5124 [hep-ph]

2nd July 2015, 9th APCTP-BLTP Workshop, Almaty,

- Motivation
- Energy scales in fundamental interactions
- Naturalness problem of SM
- Quark condensate
- Radiatively induced symmetry breaking
- Spontaneous conformal symmetry breaking in SM
- Conclusions

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM
- **No** any experimental signal of **new physics** up to about 1 TeV

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM
- **No** any experimental signal of **new physics** up to about 1 TeV
- Intrinsic **problems** of the Standard Model, Standard Cosmology, and General Relativity

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM
- **No** any experimental signal of **new physics** up to about 1 TeV
- Intrinsic **problems** of the Standard Model, Standard Cosmology, and General Relativity
- **The most serious problems are related to EXPLICIT scale invariance breaking in these theories**

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM
- **No** any experimental signal of **new physics** up to about 1 TeV
- Intrinsic **problems** of the Standard Model, Standard Cosmology, and General Relativity
- **The most serious problems are related to EXPLICIT scale invariance breaking in these theories**
- We **have to** look for alternatives . . .

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM
- **No** any experimental signal of **new physics** up to about 1 TeV
- Intrinsic **problems** of the Standard Model, Standard Cosmology, and General Relativity
- **The most serious problems are related to EXPLICIT scale invariance breaking in these theories**
- We **have to** look for alternatives . . .
- **Symmetry** principles to be exploited

Motivation (I)

- Recent discovery of the Higgs boson with mass $125.09 \pm 0.21 \pm 0.11$ GeV, which is **consistent with the SM**
- Still a **(too) good** description of all HEP observations within the SM
- **No** any experimental signal of **new physics** up to about 1 TeV
- Intrinsic **problems** of the Standard Model, Standard Cosmology, and General Relativity
- **The most serious problems are related to EXPLICIT scale invariance breaking in these theories**
- We **have to** look for alternatives . . .
- **Symmetry** principles to be exploited
- **Correspondence** to SM should be preserved

Motivation (II)

Higgs boson with $M_H \approx 125$ GeV makes the SM **stable** up to the Planck energy scale, i.e. 10^{19} GeV.

New physics is not required?


F. Bezrukov, M.Y. Kalmykov, B.A. Kniehl, M. Shaposhnikov, JHEP'2012

S. Alekhin, A. Djouadi and S. Moch, Phys. Lett. B'2012

A.V. Bednyakov, A.F. Pikelner and V.N. Velizhanin, Nucl. Phys. B'2013, 2014; Phys. Lett. B'2014.

...

SM at the Planck scale?



A.V. Bednyakov, A.F. Pikelner et al., Nucl. Phys. B'2013, 2014;
Phys. Lett. B'2014, 2015

Motivation (III)

At the EW scale we have a remarkable empirical relation

$$v = \sqrt{M_H^2 + M_W^2 + M_Z^2 + m_t^2}$$

for today PDG values we have a perfect agreement within experimental errors

$$246.22 = 246 \pm 1 \text{ GeV}$$

Obviously, there should be some tight clear relation between the top quark mass and the Higgs boson one (or the EW scale)

Note also

$$2 \frac{M_h^2}{m_t^2} = 1.05 \approx 1 \approx 2 \frac{m_t^2}{v^2} \equiv y_t^2 = 0.99$$

Motivation (IV)

We will try to apply the mechanism of the **chiral symmetry breaking** to the SM

The Nobel Prize in Physics 2008 (one half) was awarded to Yoichiro **Nambu** "for the discovery of the mechanism of spontaneous broken symmetry in subatomic physics".

The prize in 2013 was awarded to Francois **Englert** and Peter **Higgs** "for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles . . ."

Mechanisms of Spontaneous Symmetry Breaking (SSB) in SM and QCD are similar but of **different types**

Scale invariance breaking

The observed world is obviously not Scale Invariant (SI)

But many physical laws **are SI**, see e.g. Newtonian mechanics (w/o gravity) and Maxwell equations

There is **only one term** (the Higgs tachyon mass) in the SM Lagrangian, which **explicitly** breaks SI

then we have **dimensional transmutation** in QCD

and an **explicitly** dimensionful coupling constant in Gravity

All those make **real troubles** for the fundamental theory

Does the Higgs boson really give masses to everything that we see?

not really

Λ -term and dark matter in Cosmology?

the proton mass?

neutrino masses?

the Higgs mass itself?

We still do not understand the origin of masses

and of fundamental physical energy scales in general

Higgs boson in SM (I)

Remind the Standard Model mechanism:

$$V_{\text{Higgs}}(\phi) = \lambda(\phi^\dagger\phi)^2 + \mu^2\phi^\dagger\phi$$

Due to **spontaneous symmetry breaking** (SSB) of $O(4)$ symmetry if $\mu^2 < 0$, one component of the complex scalar doublet field $\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$ acquires a non-zero vacuum expectation value

$$\langle \phi^0 \rangle = v/\sqrt{2}$$

The **vacuum stability** condition $\lambda > 0$ is always assumed

Higgs boson in SM (II)

The $O(4)$ symmetry of the Higgs field is broken spontaneously down to $O(3)$, while the EW gauge symmetry is broken only **fictitiously** [L.Faddeev & A.Niemi]. The gauge and custodial symmetries protect EW observables from large quantum corrections, in particular

$$\Delta m_{W,Z}^2 \sim \alpha m_{W,Z}^2 \ln \frac{\Lambda^2}{m_{W,Z}^2}$$

But the Higgs boson mass is not protected by any symmetry, it gets huge corrections

$$\Delta m_H^2 \sim \Lambda^2$$

That is known as the **naturalness** or **fine tuning** or **hierarchy** problem of SM. That is because M_W and M_Z have the **pure** SSB origin, while M_H is related to the **tachyon** mass term ($\mu^2 < 0$) which breaks the **conformal symmetry** of SM **explicitly**

Naturalness problem (I)

There are two **general** ways to solve the naturalness problem:

I. **Cancel out** the huge radiative corrections

- either due to some (super)symmetry
- or due to fine tuning (anthropic principle)

II. Make Λ **small**, i.e. $\Lambda \lesssim 1$ TeV with some new physics motivation

- but LHC and others do not see anything new at this scale


Naturalness problem (II)

Let us look at some details of the problem.

In the SM, the Λ^2 divergent terms cancel out everywhere except the corrections to the Higgs mass

They appear as scalar Passarino-Veltman integrals

$$A_0(m^2, \Lambda^2) = \int_{\Lambda} \frac{d^4 k}{i\pi^2} \frac{1}{k^2 - m^2 + i\epsilon} = \Lambda^2 - m^2 \ln \frac{\Lambda^2}{m^2} + \mathcal{O}(\Lambda^{-2})$$




N.B. That is the so-called **tadpole** Feynman diagram

Naturalness problem (III)

Three types of diagrams contribute:

- Higgs boson loop
- EW boson loop
- top-quark loop


EW and Higgs boson loops:



N.B. Actually longitudinal components of EW bosons, i.e. goldstones, are relevant

Naturalness problem (IV)

Top quark loop



$$\int_{\Lambda_t} \frac{d^4 k}{i\pi^2} \frac{\text{Tr}(\hat{k} + m_t)((\hat{p} - \hat{k}) + m_t)}{(k^2 - m_t^2)((p - k)^2 - m_t^2)} \rightarrow 4 \int_{\Lambda_t} \frac{d^4 k}{i\pi^2} \frac{1}{k^2 - m_t^2} + \mathcal{O}(m_t^2)$$
$$= 4A_0(m_t^2, \Lambda_t^2) + \mathcal{O}(m_t^2)$$

Naturalness problem (V)

Combined in the lowest approximation (if Λ_i are the same)

$$M_H^2 = (M_H^0)^2 + \frac{3\Lambda^2}{8\pi^2 v^2} \left[M_H^2 + 2M_W^2 + M_Z^2 - 4m_t^2 \right]$$

It is **unnatural** to have $\Lambda \gg M_H$.

The most **natural** option would be $\Lambda \sim M_H$, e.g. everything is defined by the EW scale. But this is not the case of the SM.


Obviously, the problem is caused by the **explicit** breaking of the conformal symmetry in the SM

Quark condensate (I)

By definition **formally**

$$\langle \bar{q} q \rangle = -N_C \int_{\Lambda_q} \frac{d^4 k}{i(2\pi)^4} \frac{\text{Tr}(\hat{k} + m_q)}{k^2 - m^2 + i\epsilon} \sim -4N_C m_q A_0(m_q^2, \Lambda_q^2)$$

In particular the **top quark condensate** gives $\langle \bar{t} t \rangle / m_t$ contribution to ΔM_H^2 . This statement concerns as formal definitions as well as **observables**.




N.B. $\langle \bar{q} q \rangle \equiv 0$ if $m_q = 0$

Quark condensate (II)

Light quark condensate is “measured”: $\sqrt[3]{\langle \bar{q} q \rangle} \simeq -250 \text{ MeV}$

The “measurement” itself is possible due to **nonperturbative** effects at low energies

In perturbative QCD the condensate can not be accessed just due to the Furry theorem:



I.e. the condensate can be finite, but its contribution is exactly zero

Quark condensate (III)

We **do not know** exactly how does appear the low-energy QCD scale, but we see

$$-\sqrt[3]{\langle \bar{q} q \rangle} \sim M_q \sim \Lambda_{\text{QCD}}$$

where M_q is the constituent light quark mass

$$\text{Or } \langle \bar{q} q \rangle \sim -M_q \times \Lambda_{\text{QCD}}^2$$

Very likely that the Λ_{QCD} scale comes from **outside** QCD. The QCD dynamics just helps it to **propagate** into M_q and $\langle \bar{q} q \rangle$.

It is very likely that **radiatively induced dimensional transmutation** is realized in QCD. It means a **SOFT** breaking of conformal symmetry there.

Coleman-Weinberg mechanism (I)

S. Coleman & E. Weinberg 1973

Semi-classical **conformal-invariant** $V = \lambda\phi_c^4/4!$ is transformed by quantum loop corrections into

$$V_{\text{eff}} = \frac{\lambda}{4!}\phi_c^4 + \frac{\lambda^2\phi_c^4}{256\pi^2} \left(\ln \frac{\phi_c^2}{M^2} - \frac{25}{6} \right)$$

where M is a scale, which should be introduced to avoid infrared divergences.

Minimization of the effective potential leads to $\langle\phi\rangle \neq 0$ and consequently to $m_\phi \neq 0$

Coleman-Weinberg mechanism (II)

Let us apply the C-W procedure for the case of **scalar+fermion**:


$$V_{\text{cl}} = \lambda \phi_c^4 / 4! + y \phi_c \bar{f} f$$

Scalar and fermion loops give:

$$\Delta V_{\text{sc}} = \frac{1}{2} \int \frac{d^4 k}{(2\pi^4)} \ln \left(1 + \frac{\lambda \phi_c^2}{2k^2} \right) \rightarrow \frac{\lambda \Lambda^2}{256\pi^2} \phi_c^2 + \frac{\lambda^2 \phi_c^4}{256\pi^2} \left(\ln \frac{\lambda \phi_c^2}{2\Lambda^2} - \frac{1}{2} \right)$$

$$\Delta V_f = -4N_C \int \frac{d^4 k}{(2\pi^4)} \ln \left(1 + \frac{ym_f \phi_c}{k^2 - m_f^2} \right) \rightarrow -4N_C \frac{ym_f \Lambda_f^2}{16\pi^2} \phi_c$$
$$-4N_C \frac{y^2 m_f^2 \phi_c^2}{32\pi^2} \left(\ln \frac{ym_f \phi_c}{\Lambda_f^2} - \frac{1}{2} \right) + \dots$$

N.B. The first term in ΔV_f is the fermion tadpole



Coleman-Weinberg mechanism (II)

Let us apply the C-W procedure for the case of **scalar+fermion**:


$$V_{\text{cl}} = \lambda \phi_c^4 / 4! + y \phi_c \bar{f} f$$

Scalar and fermion loops give: **Renormalize** $\rightarrow 0!$

$$\Delta V_{\text{sc}} = \frac{1}{2} \int \frac{d^4 k}{(2\pi^4)} \ln \left(1 + \frac{\lambda \phi_c^2}{2k^2} \right) \rightarrow \frac{\lambda \Lambda^2}{256\pi^2} \phi_c^2 + \frac{\lambda^2 \phi_c^4}{256\pi^2} \left(\ln \frac{\lambda \phi_c^2}{2\Lambda^2} - \frac{1}{2} \right)$$

$$\Delta V_f = -4N_C \int \frac{d^4 k}{(2\pi^4)} \ln \left(1 + \frac{ym_f \phi_c}{k^2 - m_f^2} \right) \rightarrow -4N_C \frac{ym_f \Lambda_f^2}{16\pi^2} \phi_c$$
$$-4N_C \frac{y^2 m_f^2 \phi_c^2}{32\pi^2} \left(\ln \frac{ym_f \phi_c}{\Lambda_f^2} - \frac{1}{2} \right) + \dots$$

N.B. The first term in ΔV_f is fermion tadpole



Coleman-Weinberg mechanism (II)

Let us apply the C-W procedure for the case of **scalar+fermion**:

$$V_{\text{cl}} = \lambda \phi_c^4 / 4! + y \phi_c \bar{f} f$$


Scalar and fermion loops give: **Renormalize! $\rightarrow 0$**

$$\Delta V_{\text{sc}} = \frac{1}{2} \int \frac{d^4 k}{(2\pi^4)} \ln \left(1 + \frac{\lambda \phi_c^2}{2k^2} \right) \rightarrow \frac{\lambda \Lambda^2}{256\pi^2} \phi_c^2 + \frac{\lambda^2 \phi_c^4}{256\pi^2} \left(\ln \frac{\lambda \phi_c^2}{2\Lambda^2} - \frac{1}{2} \right)$$

$$\Delta V_f = -4N_C \int \frac{d^4 k}{(2\pi^4)} \ln \left(1 + \frac{y m_f \phi_c}{k^2 - m_f^2} \right) \rightarrow -4N_C \frac{y m_f \Lambda_f^2}{16\pi^2} \phi_c$$

$$-4N_C \frac{y^2 m_f^2 \phi_c^2}{32\pi^2} \left(\ln \frac{y m_f \phi_c}{\Lambda_f^2} - \frac{1}{2} \right) + \dots \quad \text{Renormalize} \rightarrow ?$$

N.B. The first term in ΔV_f is the fermion tadpole



Coleman-Weinberg mechanism (III)

Conformal-invariant **unbroken** phase (classical only):

$$m_\phi = m_f \equiv 0, \quad \langle \phi \rangle \equiv 0, \quad \langle \bar{f} f \rangle \equiv 0$$

In the **softly** broken phase for $\lambda \sim 1$ and $y \sim 1$:

$$m_\phi \sim m_f \sim M, \quad \langle \phi \rangle \sim M, \quad \langle \bar{f} f \rangle \sim -M^3$$

like in QCD, but **non-perturbativity is not required**

Let us look for a **stable solution** in the broken phase

SCSB for Higgs (I)

The dominant terms of Higgs interactions (for $\mu \equiv 0$) are

$$L_{\text{int}} = -\frac{\lambda}{4}\phi^4 - \frac{y_t}{\sqrt{2}}\phi \bar{t}t$$

C.-W. mechanism gives the leading **effective potential** in the form

$$V_{\text{cond}}(\phi) = \frac{\lambda}{4}\phi^4 + \frac{y_t}{\sqrt{2}}\langle \bar{t}t \rangle \phi$$

The extremum condition

$$\left. \frac{dV_{\text{cond}}}{d\phi} \right|_{\phi=v} = 0 \quad \longrightarrow \quad v^3 = -\frac{y_t}{\sqrt{2}}\langle \bar{t}t \rangle$$

The Yukawa coupling $y_t \approx 0.99$ is known from $m_t = v \cdot y_t / \sqrt{2}$

The potential takes the form

$$V_{\text{cond}}(\phi)|_{\phi=v+H} = V_{\text{cond}}(v) + \frac{3\lambda v^2}{2}H^2 + \lambda v H^3 + \frac{\lambda}{4}H^4$$

SCSB for Higgs (II)

So the Higgs mass is

$$M_H^2 = 3\lambda v^2 = -\frac{3y_t \langle \bar{t} t \rangle}{\sqrt{2} v} = -\frac{3m_t \langle \bar{t} t \rangle}{v^2}$$

N.B. The difference from the SM is in the value of λ :

$$\lambda = \frac{2}{3}\lambda_{\text{SM}}$$

Top quark condensate (I)

To get $M_H = 126$ GeV we need $\langle \bar{t} t \rangle = (-123 \text{ GeV})^3$

It is just a **natural value** according to the **naturalness principle**

There are no any (other) phenomenological restrictions on $\langle \bar{t} t \rangle$

Having non-zero top quark condensate does **NOT** lead to top quark bound states in our case

Top quark condensate (II)

We have an idea, how to evaluate a quark condensate using a suppression form factor which should take into account **physical vacuum** properties:

$$\begin{aligned}\langle 0|\bar{q}q|0\rangle &\rightarrow_{\text{phys}} \langle 0|\bar{q}q|0\rangle_{\text{phys}} \\ &= -4m_t^3 N_c \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\sqrt{\mathbf{p}^2+1}} F(\mathbf{p}) \approx 0.39 \\ F(\mathbf{p}) &= \left[1 + \exp\left(\sqrt{\mathbf{p}^2+1} - 1\right) \right]^{-1}, \quad \mathbf{p} \equiv \frac{\vec{p}}{m_t}\end{aligned}$$

That gives $\langle \bar{t}t \rangle \approx (-126 \text{ GeV})^3$

N.B. $F(\mathbf{p})$ is a typical distribution function with an “effective temperature” $T \simeq m_t$. A similar distribution function is known to regularize the Casimir energy.

Naturalness (once more)

W. Bardeen (1995): radiative stability of the Higgs boson mass, i.e. resolution of the naturalness problem, can be ensured by the classical scale invariance

The constructed semi-classical solution is **stable** at least around the EW scale

For $\lambda \sim 1$ and $y_t \approx 1$ it is **natural** to have

$$m_t \sim M_H \sim v \sim \sqrt[3]{-\langle \bar{t} t \rangle}$$

Coleman-Weinberg: we **have to** introduce a finite scale but not into the Lagrangian. It can be a property of the quantum **physical vacuum**. How does the scale defines the observables depends on the model.

Conclusions

1. We proposed a simple modification of the SM based on the Nambu condensate mechanism. The difference from SM is only in 1.5 times lower value of the Higgs self-coupling λ
2. Here M_H and m_t are mutually related and define together EW scale
3. Our estimate of the top quark condensate value looks natural
4. The suggested mechanism automatically protects M_H from running away, since renormalization happens at the EW scale
5. The picture resembles the EW **bootstrap** suggested by Nambu and Bardeen et al. (1989). But their approaches were not based on the conformal symmetry. They just tried to cancel out the quadratic divergences.
6. Similar relations are used also in modern **technicolor** models, but the Higgs boson is composite there