

Identity method – a new tool for studying chemical fluctuations

M. Maćkowiak for the NA49 Collaboration^{1,*}

¹*Faculty of Physics, Warsaw University of Technology, Poland*

Event-by-event fluctuations of the chemical composition of the hadronic system produced in nuclear collisions are believed to be sensitive to properties of the transition between confined and deconfined strongly interacting matter.

In this paper a new technique for the study of chemical fluctuation, the identity method, is introduced and its features are discussed. The method is tested using data on central PbPb collisions at 40 AGeV registered by the NA49 experiment at the CERN SPS.

1. INTRODUCTION

The most interesting features of the phase diagram of strongly interacting matter are the Critical Point (CP) and the 1st order phase transition line. Event-by-event fluctuations of the chemical composition of the hadronic system produced in nuclear collisions are believed to be sensitive to both of them. The first relevant measurements were performed by the NA49 experiment at the CERN SPS. A systematic scan in beam energy and system size was recently started by the NA61 collaboration. Furthermore, additional insight is expected from the RHIC beam energy scan program.

There are several measures used to quantify chemical fluctuations, among them: σ_{dyn} [1–3] used by NA49 and ν_{dyn} [4] used by STAR. Both are related as $\sigma_{\text{dyn}}^2 \approx \nu_{\text{dyn}}$ and share the same disadvantage, namely, they depend on volume (number of wounded nucleons) and volume fluctuations in thermodynamical (wounded nucleon) models. Another measure Φ_x [5, 6] was used by NA49 to characterize transverse momentum [7, 8], electric charge [9], and azimuthal angle fluctuations [10]. It is free of the mentioned disadvantage of σ_{dyn} , ν_{dyn} . However, all these measures of chemical fluctuations are affected by non-perfect particle identification. This is illustrated in Fig. 1a, where the spectrum of the specific energy loss

* Electronic address: majam@if.pw.edu.pl

(dE/dx) of particles measured by the NA49 Time Projection Chambers [11] is shown for a selected phase-space bin. The dE/dx signal depends on the particle mass and together with the particle charge measurement is used to identify particles. It is seen that the dE/dx distributions of different particle species partly overlap and thus unique particle identification is not possible.

The identity method adapts the Φ_x measure to take into account a non-unique particle identification, while keeping the advantages of Φ_x .

2. IDENTITY METHOD

Let us assume that particles are identified according to their measured mass. The measured mass spectra of all particles and of particles of type h in the analyzed event sample are denoted as ρ and ρ_h , respectively. The spectra are normalized to the corresponding mean multiplicities per event, namely:

$$\int dm \rho(m) = \langle N \rangle , \quad \int dm \rho_h(m) = \langle N_h \rangle . \quad (1)$$

Furthermore, we define a single particle variable called the particle identity as:

$$w_h(m) \stackrel{\text{def}}{=} \frac{\rho_h(m)}{\rho(m)} . \quad (2)$$

The fluctuation measure Ψ_{wh} is then introduced in a way similar (the roots over the two components are absent) to the Φ measure. First, a single particle variable z is defined as:

$$z \stackrel{\text{def}}{=} w_h - \bar{w}_h , \quad (3)$$

where the bar denotes the inclusive mean and thus $\bar{w} = \langle N_h \rangle / \langle N \rangle$. Second, an event variable Z , which is the multiparticle analog of z , is calculated as:

$$Z \stackrel{\text{def}}{=} \sum_{i=1}^N (w_h(m_i) - \bar{w}_h) , \quad (4)$$

where N is the multiplicity and i is the particle index in an event.

Finally, the fluctuation measure Ψ_{wh} is defined as:

$$\Psi_{wh} \stackrel{\text{def}}{=} \frac{\langle Z^2 \rangle}{\langle N \rangle} - \bar{z}^2 . \quad (5)$$

For further analysis one denotes two possible values of Ψ_{wh} :

- Ψ_{res} which is the value of Ψ_{wh} for the experimental mass resolution case,
- Ψ_{corr} which is the value of Ψ_{wh} for the perfect mass resolution case.

In order to correct for the non-unique particle identification we calculate the variance per particle due to random identification for the experimental mass resolution case:

$$\text{Var}_{\text{res}} = \frac{1}{\langle N \rangle} \int_0^\infty dm \rho(m) \cdot w_h(m)(1 - w_h(m)) . \quad (6)$$

It is easy to show that for unique particle identification (the perfect mass resolution case, $w_h(m) = \delta(m - m_h)$) the result is $\text{Var}_{\text{res}} \stackrel{\text{def}}{=} \text{Var}_A = 0$, whereas for no mass resolution ($w_h(m) = \text{const}$) one obtains $\text{Var}_{\text{res}} \stackrel{\text{def}}{=} \text{Var}_B = \frac{\langle N_h \rangle}{\langle N \rangle} (1 - \langle N_h \rangle / \langle N \rangle)$. For the experimental data analysis the integral in Eq. (6) is replaced by a sum over all particles.

The following key relation can be proven ¹:

$$\Psi_{\text{corr}} = \Psi_{\text{res}} (1 - \text{Var}_{\text{res}} / \text{Var}_B)^{-2} . \quad (7)$$

It shows that the measured fluctuations Ψ_{res} can be corrected for the effect of non-unique particle identification in a model independent way. This is because the correction factor, $(1 - \text{Var}_{\text{res}} / \text{Var}_B)^{-2}$, depends only on the experimental resolution and mean particle multiplicities.

Eq. 7 was checked by numerous Monte Carlo simulations with different types of correlations, mass resolution functions and mean particle multiplicities. The results of these simulations are shown in Fig. 1b.

3. THE IDENTITY METHOD TEST USING NA49 DATA

In the analysis of experimental data we use the particle energy loss dE/dx as the measure of the mass m . For optimal identification the dE/dx spectra from the NA49 TPCs are determined in bins of total and transverse momentum, azimuthal angle as well as for both electric charges separately [1–3]. In each bin four Gauss functions (for electrons, pions, kaons and protons) are fitted. An example of such a fit is displayed in Fig. 1a. The fitted functions are then used as the ρ_h and ρ functions of the identity method. The further analysis steps are as follows:

¹ M. Gaździcki, K. Grebieszko, M. Maćkowiak, and S. Mrówczyński publication in preparation

- using mean particle multiplicities the variance $\text{Var}_B = \langle N_h \rangle / \langle N \rangle (1 - \langle N_h \rangle / \langle N \rangle)$ is obtained;
- for each particle its identity is calculated

$$w_{hi}(\langle dE/dx \rangle_i | q, p_{\text{tot}}, p_T, \phi) = \frac{\rho_h(\langle dE/dx \rangle_i | q, p_{\text{tot}}, p_T, \phi)}{\rho(\langle dE/dx \rangle_i | q, p_{\text{tot}}, p_T, \phi)}, \quad (8)$$

- using the experimental dE/dx resolution functions, ρ_h and ρ (M is the total number of particles used in the analysis) the variance

$$\text{Var}_{\text{res}} = \frac{1}{M} \sum_1^M w_{hi}(\langle dE/dx \rangle_i | q, p_{\text{tot}}, p_T, \phi) (1 - w_{hi}(\langle dE/dx \rangle_i | q, p_{\text{tot}}, p_T, \phi)), \quad (9)$$

is computed;

- using the identity values, w_{hi} , Ψ_{res} is calculated and
- the corrected value of Ψ_{corr} is obtained using Eq. (7).

As a first test of the identity method proton fluctuations were studied in PbPb collisions at 40 AGeV energy. Positively and negatively charged particles with total momentum up to 40 GeV/ c and transverse momentum up to 2 GeV/ c were used for the analysis. The total number of analyzed events was 4000. The mean multiplicities are $\langle N \rangle = 165.40$, and $\langle N_p \rangle = 42.16$. The obtained value of $\Psi_{\text{res}} \cdot 1000 = -17.4 \pm 3.5$. The corresponding correction factor for non-unique particle identification was calculated to be ≈ 1.2 . The value corrected for the finite resolution is $\Psi_{\text{corr}} \cdot 1000 = -22.3 \pm 4.4$. The analysis of PbPb collisions at all NA49 energies is in progress.

-
1. S. V. Afanasev et al. (NA49 Collab.), Phys. Rev. Lett. **86**, 1965 (2001).
 2. C. Alt et al. (NA49 Collab.), Phys. Rev. C **79**, 044910 (2009).
 3. D. Kresan (for the CBM and NA49 Collabs.), PoS CPOD 2009, 031 (2009); arXiv: 0908.2875.
 4. B. I. Abelev et al. (STAR Collab.), Phys. Rev. Lett. **103**, 092301 (2009).
 5. M. Gazdzicki, S. Mrowczynski, Z. Phys. C **54**, 127 (1992).
 6. M. Gazdzicki, Eur. Phys. J. C **8**, 131 (1999).
 7. T. Anticic et al. (NA49 Collab.), Phys. Rev. C **70**, 034902 (2004).

8. T. Anticic et al. (NA49 Collab.), Phys. Rev. C **79**, 044904 (2009).
9. C. Alt et al. (NA49 Collab.), Phys. Rev. C **70**, 064903 (2004).
10. T. Cetner, K. Grebieszko, S. Mrówczyński, arXiv: 1011.1631
11. S. Afanasev et al. (NA49 Collab.), Nucl. Instrum. Meth. A **430**, 210-244 (1999).




Figure 1. (a): Distribution of specific energy loss measured in the NA49 TPCs for positively charged particles is a bin $p_{\text{tot}} \in (4.4\text{--}5.3)$ GeV/ c , $p_T \in (0.0\text{--}0.2)$ GeV/ c and $\phi \in (0.75\pi\text{--}\pi)$. The fitted Gauss functions are shown by solid curves. (b): The ratio $\Psi_{\text{res}}/\Psi_{\text{corr}}$ versus $\text{Var}_{\text{res}}/\text{Var}_{\text{B}}$ calculated within several Monte Carlo simulations with different parameters of experimental resolution, particle multiplicities and fluctuations. The results agree with the analytical dependence given by Eq. (7).

FIGURE CAPTIONS

Fig. 1: (a): Distribution of specific energy loss measured in the NA49 TPCs for positively charged particles is a bin $p_{\text{tot}} \in (4.4-5.3)$ GeV/ c , $p_T \in (0.0-0.2)$ GeV/ c and $\phi \in (0.75\pi-\pi)$. The fitted Gauss functions are shown by solid curves. (b): The ratio $\Psi_{\text{res}}/\Psi_{\text{corr}}$ versus $\text{Var}_{\text{res}}/\text{Var}_{\text{B}}$ calculated within several Monte Carlo simulations with different parameters of experimental resolution, particle multiplicities and fluctuations. The results agree with the analytical dependence given by Eq. (7).