

Thermodynamics of hadron matter in PNJL model

A. V. Friesen

BLTP, JINR

The modern sketch of HIC

Experiment

The picture of the heavy ion collision's evolution






Figure 1: arXiv (nucl-th): 1304.3634

The QCD phase diagram



- chiral symmetry restoration (constituent quarks \rightarrow current quarks);
- deconfinement;


The QCD phase diagram



- chiral symmetry restoration (constituent quarks \rightarrow current quarks);
- deconfinement;

Do they coincide?

The QCD phase diagram




- chiral symmetry restoration (constituent quarks \rightarrow current quarks);
- deconfinement;

Do they coincide?

Lattice QCD

Hands S. Contemp. Phys. 42, 209 [2001], $T_c = 0.17$ GeV (SU(2))



The Nambu-Jona-Lasinio model

The Lagrangian

$$\mathcal{L}_{\text{NJL}} = \bar{q} (i\not{\partial} - \hat{m}_0 - \gamma_0\mu) q + G_s \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5\vec{\tau}q)^2 \right],$$

G_s the effective coupling strength,

\bar{q} и q - quark fields

$\hat{m}_0 = \text{diag} (m_u^0, m_d^0)$, $m_u^0 = m_d^0$ - the current quark masses, $\vec{\tau}$ - Pauli matrices SU(2).

M. K. Volkov, *Ann. Phys.* 157,282 (1989); *Sov. J. Part and Nuclei* 17, 433 (1986) S. P. Klevansky, *Rev. Mod. Phys.* 64, 649 (1992).

We can:

The Nambu-Jona-Lasinio model

The Lagrangian

$$\mathcal{L}_{\text{NJL}} = \bar{q} (i\not{\partial} - \hat{m}_0 - \gamma_0\mu) q + G_s \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5\vec{\tau}q)^2 \right],$$

G_s the effective coupling strength,

\bar{q} и q - quark fields

$\hat{m}_0 = \text{diag} (m_u^0, m_d^0)$, $m_u^0 = m_d^0$ - the current quark masses, $\vec{\tau}$ - Pauli matrices SU(2).

M. K. Volkov, *Ann. Phys.* 157,282 (1989); *Sov. J. Part and Nuclei* 17, 433 (1986) S. P. Klevansky, *Rev. Mod. Phys.* 64, 649 (1992).

We can:

- explain spontaneous chiral symmetry broken as $m_q = m_0 + \langle \bar{q}q \rangle$;

The Nambu-Jona-Lasinio model

The Lagrangian

$$\mathcal{L}_{\text{NJL}} = \bar{q} (i\cancel{\partial} - \hat{m}_0 - \gamma_0 \mu) q + G_s \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5 \vec{\tau}q)^2 \right],$$

G_s the effective coupling strength,

\bar{q} и q - quark fields

$\hat{m}_0 = \text{diag} (m_u^0, m_d^0)$, $m_u^0 = m_d^0$ - the current quark masses, $\vec{\tau}$ - Pauli matrices SU(2).

M. K. Volkov, *Ann. Phys.* 157,282 (1989); *Sov. J. Part and Nuclei* 17, 433 (1986) S. P. Klevansky, *Rev. Mod. Phys.* 64, 649 (1992).

We can:

- explain spontaneous chiral symmetry broken as $m_q = m_0 + \langle \bar{q}q \rangle$;
- describe chiral phase transitions.

The Nambu-Jona-Lasinio model

The Lagrangian

$$\mathcal{L}_{\text{NJL}} = \bar{q} (i\not{\partial} - \hat{m}_0 - \gamma_0\mu) q + G_s \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5\vec{\tau}q)^2 \right],$$

G_s the effective coupling strength,

\bar{q} и q - quark fields

$\hat{m}_0 = \text{diag} (m_u^0, m_d^0)$, $m_u^0 = m_d^0$ - the current quark masses, $\vec{\tau}$ - Pauli matrices SU(2).

M. K. Volkov, *Ann. Phys.* 157,282 (1989); *Sov. J. Part and Nuclei* 17, 433 (1986) S. P. Klevansky, *Rev. Mod. Phys.* 64, 649 (1992).

We can:

- explain spontaneous chiral symmetry broken as $m_q = m_0 + \langle \bar{q}q \rangle$;
- describe chiral phase transitions.
- describe light quarks and mesons properties,

The mean - field approximation

We can introduce the partition function

$$\mathcal{Z}[\bar{q}, q] = \int \mathcal{D}\bar{q} \mathcal{D}q \exp \left\{ \int_0^\beta d\tau \int_v d^3x [\mathcal{L}_{\text{NJL}}] \right\}. \quad (1)$$

Then, using the mean-field approximation procedure, we get

$$\mathcal{Z}_{\text{MF}}[\bar{q}, q] = \exp \left\{ - \int_0^\beta d\tau \int_v d^3x \frac{\sigma_{\text{MF}}'^2}{4G} + \text{Tr} \ln S_{\text{MF}}^{-1}[m] \right\}. \quad (2)$$

And then

$$\Omega_{\text{NJL}}(T, \mu) = -\frac{T}{V} \ln Z_{\text{MF}}(\bar{q}, q). \quad (3)$$

The grand potential

$$\Omega_{\text{NJL}} = G_s \langle \bar{q}q \rangle^2 - 2N_c N_f \int \frac{d^3p}{(2\pi)^3} E_p - 2N_c N_f T \int \frac{d^3p}{(2\pi)^3} [\ln N^+(E_p) + \ln N^-(E_p)] \quad (4)$$

where $N^+(E_p) = 1 + e^{-\beta(E_p - \mu)}$, $N^-(E_p) = 1 + e^{-\beta(E_p + \mu)}$
 $E_p = \sqrt{p^2 + m^2}$ - quark energy and $\beta = 1/T$.

The model parameters

The model has free parameters:

- m_0 - current quark mass,
- Λ - three-momentum cut-off,
- G_s - the effective coupling strength

To fix the parameters we use the experimental data:

- The pion decay constant $f_\pi = 0.092$ GeV,
- The pion mass $M_\pi = 0.139$ GeV
- The quark condensate $\langle \bar{q}q \rangle = (-0.25 \text{ GeV})^3$

	m_0 [MeV]	Λ [GeV]	G_s [GeV] ⁻²	f_π [GeV]	m_π [GeV]	m [GeV]
Set A	5.5	0.639	5.227	0.092	0.139	0.319
Set B	5.6	0.646	5.56	0.099	0.141	0.394

Table 1: The NJL parameters.




Figure 2: The NJL phase diagram.

Set A: $T_c = 0.186$ GeV ,

$T_{\text{CEP}} (0.05, 0.3165)$

Set B: $T_c = 0.2265$ GeV,

$T_{\text{CEP}} (0.08, 0.3425)$

A. V. Friesen, Yu. L. Kalinovsky,

Phys. Part. Nucl. Lett. 6, 737

(2015)

The critical temperatures:

- $T_{\text{Mott}} (M_\pi = 2m_q)$
- T_c - crossover line

$$\max \frac{\partial \langle q\bar{q} \rangle}{\partial T}$$
- 1st-order transition -

$$\max \frac{\partial^2 \Omega}{\partial \mu^2} \Big|_{T=\text{const}}$$
- T_{CEP}

The NJL model:

- can reproduce chiral phase transition;
- shows crossover phase transition at low density and high temperature;
- shows 1st order transition at low temperature and high density;
- is local theory and cannot describe confinement/deconfinement properties.

The Polyakov-loop extended Nambu-Jona-Lasinio model

$$\mathcal{L}_{\text{PNJL}} = \bar{q} (i\gamma_\mu D^\mu - \hat{m}_0 - \gamma_0 \mu) q + G_s \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5 \vec{\tau}q)^2 \right] - \mathcal{U}(\Phi, \bar{\Phi}; T)$$

C. Ratti, M. Thaler, W. Weise, PRD 73, 014019 (2006)

$q = (q_u, q_d)$ quark fields,

$\hat{m}_0 = \text{diag}(m_u^0, m_d^0)$ -current quark masses, $m_u^0 = m_d^0 = m_0$

$D^\mu = \partial^\mu - iA^\mu$ - covariant derivative,

$A^\mu(x) = g\mathcal{A}_a^\mu \frac{\lambda_a}{2}$, \mathcal{A}_a^μ the gauge field SU(3),

$A^\mu = \delta_0^\mu A^0 = -i\delta_4^\mu A_4$,

λ_a - Gell-Mann matrices,


G_s - scalar coupling strength.

The Polyakov field Φ is determined as: $\Phi[A] = \frac{1}{N_c} \text{Tr}_c L(\vec{x})$,

$$L(\vec{x}) = \mathcal{P} \exp \left[i \int_0^\beta d\tau A_4(\vec{x}, \tau) \right],$$

$$\langle l(\vec{x}) \rangle = e^{-\beta \Delta F_Q(\vec{x})}.$$

The effective potential



M. Panero, PRL 103, 232001 (2009)

G. Boyd et. al, NPB 469, 419 (1996)

The effective potential

Polynomial fit:

$$\frac{\mathcal{U}(\Phi, \bar{\Phi}; T)}{T^4} = -\frac{b_2(T)}{2} \bar{\Phi}\Phi - \frac{b_3}{6} (\Phi^3 + \bar{\Phi}^3) + \frac{b_4}{4} (\bar{\Phi}\Phi)^2,$$
$$b_2(T) = a_0 + a_1 \left(\frac{T_0}{T}\right) + a_2 \left(\frac{T_0}{T}\right)^2 + a_3 \left(\frac{T_0}{T}\right)^3.$$

The effective potential

Polynomial fit:



$$\frac{\mathcal{U}(\Phi, \bar{\Phi}; T)}{T^4} = -\frac{b_2(T)}{2} \bar{\Phi}\Phi - \frac{b_3}{6} (\Phi^3 + \bar{\Phi}^3) + \frac{b_4}{4} (\bar{\Phi}\Phi)^2,$$
$$b_2(T) = a_0 + a_1 \left(\frac{T_0}{T}\right) + a_2 \left(\frac{T_0}{T}\right)^2 + a_3 \left(\frac{T_0}{T}\right)^3.$$

Logarithmic fit:

$$\frac{\mathcal{U}(\Phi, \bar{\Phi}; T)}{T^4} = -\frac{1}{2}a(T) \bar{\Phi}\Phi + b(T) \ln [1 - 6\bar{\Phi}\Phi + 4(\bar{\Phi}^3 + \Phi^3) - 3(\bar{\Phi}\Phi)^2],$$
$$a(T) = \tilde{a}_0 + \tilde{a}_1 \left(\frac{T_0}{T}\right) + \tilde{a}_2 \left(\frac{T_0}{T}\right)^2, \quad b(T) = \tilde{b}_3 \left(\frac{T_0}{T}\right)^3.$$


Breaking of Z_3 symmetry

$T = 0.05 \text{ GeV}$




Breaking of Z_3 symmetry

$T = 0.05 \text{ GeV}$




$T = T_0 = 0.27 \text{ GeV}$




Breaking of Z_3 symmetry


$T = 0.05 \text{ GeV}$




$T = T_0 = 0.27 \text{ GeV}$



$T = 3T_0 = 0.81 \text{ GeV}$



The effective potential parametrization



M. Panero, PRL 103, 232001 (2009)

G. Boyd et. al, NPB 469, 419 (1996)

- $\Phi \rightarrow 1$, $p/T^4 \rightarrow 1.75$, where $T \rightarrow \infty$
- $\Rightarrow \tilde{a}_0 = 3.51$ for logarithmic fit
 $1.75 = a_0/2 + b_3/3 - b_4/4$ for polynomial fit
- $\frac{\partial \mathcal{U}(\Phi, \bar{\Phi}, T)}{\partial \Phi} \Big|_{\mu=0} = 0$ ($\Phi = \bar{\Phi}$ at $\mu = 0$)
 \Rightarrow the mean square method $\Rightarrow a_i, b_i$

A. V. Friesen et al, IJMP A27, 1250013 (2012)

Parameters

	\tilde{a}_0	\tilde{a}_1	\tilde{a}_2	\tilde{b}_3	a_0	a_1	a_2	a_3	b_3	b_4
Set A	3.51	-2.47	15.2	-1.75	6.75	-1.95	2.625	-7.44	0.75	7.5
Set B	3.51	-5.121	20.99	-2.09	6.47	-4.62	7.95	-9.09	1.03	7.32




Figure 3: Effective potential as function Φ for different temperatures.




Figure 4: Polyakov loop field Φ for (a) polynomial and (b) logarithmic effective potentials. The solid (dashed) curve corresponds to the sets B (A) parameters. (Lattice data from [Karsch F, Laermann E, Peikert A Phys. Lett. B 478 447 (2000)].)

The mean-field approximation

- The PNJL grand potential ($N_f = 2$):

$$\Omega(\Phi, \bar{\Phi}, m, T, \mu) = \mathcal{U}(\Phi, \bar{\Phi}; T) + G \langle \bar{q}q \rangle^2 - 2N_c N_f \int \frac{d^3p}{(2\pi)^3} E_p - 2N_f T \int \frac{d^3p}{(2\pi)^3} [\ln N_{\Phi}^+(E_p) + \ln N_{\bar{\Phi}}^-(E_p)],$$


where $N_{\Phi}^{\pm}(E_p) = \left[1 + 3 \left(\Phi + \bar{\Phi} e^{-\beta E_p^{\pm}} \right) e^{-\beta E_p^{\pm}} + e^{-3\beta E_p^{\pm}} \right]$

and $E_p = \sqrt{p^2 + m^2}$ - quark energy; $E_p^{\pm} = E_p \mp \mu$.

- the equations of motion


$$\frac{\partial \Omega_{\text{MF}}}{\partial \sigma_{\text{MF}}} = 0, \quad \frac{\partial \Omega_{\text{MF}}}{\partial \Phi} = 0, \quad \frac{\partial \Omega_{\text{MF}}}{\partial \bar{\Phi}} = 0.$$

Symmetries restoration and breaking




Crossover transition

$$\left. \frac{\partial \langle \bar{q}q \rangle}{\partial T} \right|_{\mu=\text{const}}$$





1st order transition Three solves of equation \rightarrow the quark susceptibility:

$$\frac{\chi_q(T, \mu)}{T^2} = \frac{\partial^2(p/T^4)}{\partial(\mu/T)^2} = \frac{\partial}{\partial(\mu/T)} (\rho/T^3).$$



Phase diagram of PNJL model

Parameters: $m_0, \Lambda, G_s, a_i, b_i, T_0 = 0.27 \Gamma \Theta B$



	T_c [GeV]	T_{CEP}
NJL _{3D} ^A	0.186	(0.05, 0.3165)
PNJL _{new} ^{pol}	0.2395	(0.118, 0.3166)
PNJL _{old} ^{pol}	0.253	(0.11, 0.3192)
PNJL _{new} ^{log}	0.23	(0.10, 0.3175)
PNJL _{old} ^{log}	0.234	(0.09, 0.322)

PNJL with vector interaction


Introduction of vector interaction into model

$$\mathcal{L}_{\text{PNJL}} = \bar{q} (i\gamma_\mu D^\mu - \hat{m}_0 - \gamma_0 \mu) q + G_s \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5 \vec{\tau}q)^2 \right] - G_v (\bar{q}\gamma_\nu q)^2 - \mathcal{U}(\Phi, \bar{\Phi}; T)$$

leads to re-normalization of chemical potential:

$$\tilde{\mu} = \mu - 4G_v N_c N_f \int_\Lambda \frac{d^3p}{(2\pi)^3} \frac{m}{E_p} [f_\Phi^+ + f_\Phi^-].$$

$$T_0 = 0.19 \text{ GeV}$$



Extended PNJL



It is possible to introduce a phenomenological dependence of $G_s(\Phi)$ and $G_v(\Phi)$:

$$\begin{aligned}\tilde{G}_s(\Phi) &= G_s[1 - \alpha_1 \Phi\bar{\Phi} - \alpha_2(\Phi^3 + \bar{\Phi}^3)], \\ \tilde{G}_v(\Phi) &= G_v[1 - \alpha_1 \Phi\bar{\Phi} - \alpha_2(\Phi^3 + \bar{\Phi}^3)],\end{aligned}$$


with $\alpha_1 = \alpha_2 = 0.2$.

Y. Sakai et al PRD 82, 076003 (2010)

P. de Forcrand, O. Philipsen NPB 642, 290(2002)



A. V. Friesen et al. IJMP A30 1550089 (2015)




- Masayuki A, Koichi Y Nucl. Phys. A 504 668 (1989);
Scavenius O et al. Phys. Rev. C 64 045202 (2001); nucl-th/0007030
Ejiri S et al. Theor. Phys. Suppl. 153 118 (2003)
Gavai R V, Gupta S Phys. Rev. D 71 114014 (2005)
Fodor Z, Katz S D JHEP (03) 014 (2002)
Fodor Z, Katz S D JHEP (04) 050 (2004)

Crossover curvature

It was suggested that critical curves for all physical quantities (chiral condensate, quark susceptibility, strange quark susceptibility, Polyakov loop) must meet at one point, which is the CEP.



$$\frac{T_c(\mu)}{T_c(0)} = 1 - k \left(\frac{\mu}{T_c(\mu)} \right)^2.$$



Thermodynamics of PNJL model

Pressure

$$\frac{p}{T^4} = \frac{p(T, \mu, m) - p(0, 0, m)}{T^4}.$$









Figure 5: Pressure as a function of the chemical potential (left panel). Pressure ΔP (solid, dashed and dashed-dotted lines) as a function of the quark density for different temperatures indicated near the curves. The dotted curve depicts the spinodal domain boundary (right panel).

A. V. Friesen, Yu. L. Kalinovsky, V. D. Toneev Phys. Usp. 59 (4) 367 (2016)

The quark density

$$n_q = - \left(\frac{\partial \Omega}{\partial \mu} \right)_T.$$



In PNJL model we can

- describe the confinement properties & describe the chiral symmetry;
- check how additional interactions (vector interaction and extended couplings) effect on phase diagram;

Thank you for attention