



Mini-Workshop on "Lattice and Functional Techniques for Exploration of Phase Structure and Transport Properties in Quantum Chromodynamics", Dubna, July 10 - 14, 2017

**Topology  
in hot QCD with a dynamical charm  
(and axion physics)**



M.P. Lombardo

*in collaboration with*


A. Trunin, F. Burger, E.-M. Ilgenfritz, M. Muller-Preussker<sup>†</sup>

## Temperatures:

$$150 \text{ MeV} < T < 500 \text{ MeV}$$

..and beyond

Quark Gluon Plasma:  
E.-M. Ilgenfritz's talk.



## Time from Big Bang




## Temperatures


$$150 \text{ MeV} < T < 500 \text{ MeV}$$

..and beyond

Temperature and Time from Big Bang are linked by the Equation of State



Quark Gluon Plasma:  
E.-M. Ilgenfritz's talk.



# The Equation of State of the Quark Gluon Plasma paves the way to Cosmology

Cold Dark Matter candidates might have been created after the inflation


Several CDM candidates are highly speculative - but one, **the axion**, is

**Theoretically well motivated in QCD**

**Amenable to quantitative estimates once QCD topological properties are known:**

## Post-inflationary axions

$$m_a(T) = \sqrt{\chi(T)} / f_a$$



## The two faces of QCD topology



Window to Axions

Friday's talk

Property of Quark Gluon Plasma

Today's talk

# QCD topology, long standing focus of strong interaction:

- learning about the structure of the (s)QGP
- fundamental symmetries, strongCP problem  $\rightarrow$  axions
- hampered by technical difficulties

## Recent developments:

- methodological progress: gradient flow, chiral fermions
- first results for dynamical fermions at high temperature:

Trunin *et al.* **J.Phys.Conf.Ser. 668 (2016) no.1, 012123**

Bonati *et al.* **JHEP 1603 (2016) 155**

Borsany *et al.* **Nature 539 (2016) no.7627, 69-71**

Petreczky *et al.* **Phys.Lett. B762 (2016) 498-505**

Burger *et al.* **Nucl. Phys. A, in press**

Taniguchi *et al.* **Phys.Rev. D95 (2017) no.5, 054502**

This talk

+ work in progress

# Outline

- Lattice setup -

- Results:

$\chi_{top}$  - *Gluonic operator and gradient flow*


$\chi_{top}$  - *Fermionic operator*

-Comments/outlook

# Our setup at a glance

Talk by E.M. Ilgenfritz


# Hot QCD and Nf 2+1+1 twisted mass Wilson fermions






# Why $N_f = 2 + 1 + 1$ ? Why Wilson twisted?

## QCD Symmetries, lattice and the real world



c,b,t do not participate in the chiral dynamics around the critical temperature. Lattice simulations **around  $T_c$**  are then performed with up,down, strange quarks –  $N_f = 2+1$



	$SU(N) \times SU(N)$	$U_A(1)$
Staggered	Remnant $U(1)$	Broken
Wilson	Broken	Broken
Domain Wall	Exact (for $L \rightarrow \infty$ )	Exact (for $L \rightarrow \infty$ )
Overlap	Exact	Exact
Wilson twisted	As good as staggered	Broken

**NB: Doubling**

**Good compromise**



# Why $N_f = 2 + 1 + 1$ ?

$T_c$

340 – 380 MeV  
RHIC AuAu  
200 GeV

420-480 MeV  
LHC  
2.76 TeV


500- 600MeV  
LHC hot spots  
2.76 TeV



1 GeV  
LHC  
7 TeV

$\approx 200\text{MeV}$

## Quark Gluon Plasma @ Colliders



Analytic studies suggest that a dynamical charm becomes relevant above 400 MeV, well within the reach of LHC

Laine Schroeder 2006

Fixed  
varying  
scale

For each lattice spacing we explore a range of temperatures 150MeV — 500 MeV by varying  $N_t$

We repeat this for three different lattice spacings following ETMC T=0 simulations.

Four pion masses

Advantages: we rely on the setup of ETMC T=0 simulations. Scale is set once for all.

Disadvantages: mismatch of temperatures - need interpolation before taking the continuum limit

Number of flavours	$m_{\pi^\pm}$
	210
$N_f = 2 + 1 + 1$	260
	370
	470
$N_f = 2$	360
	430

## Setup

$T = 0$ (ETMC) nomenclature	$\beta$	$a$ [fm] [6]	$N_\sigma^3$	$N_\tau$	$T$ [MeV]	# confs.				
A60.24	1.90	0.0936(38)	$24^3$	5	422(17)	585				
				6	351(14)	1370				
				7	301(12)	341				
				8	263(11)	970				
				9	234(10)	577				
				10	211(9)	525				
			$32^3$	11	192(8)	227				
				12	176(7)	1052				
				13	162(7)	294				
				14	151(6)	1988				
				B55.32	1.95	0.0823(37)	$32^3$	5	479(22)	595
								6	400(18)	345
								7	342(15)	327
								8	300(13)	233
9	266(12)	453								
10	240(11)	295								
11	218(10)	667								
12	200(9)	1102								
13	184(8)	308								
14	171(8)	1304								
D45.32	2.10	0.0646(26)	$32^3$	15	160(7)	456				
				16	150(7)	823				
				6	509(20)	403				
				7	436(18)	412				
				8	382(15)	416				
				10	305(12)	420				
			$40^3$	12	255(10)	380				
				14	218(9)	793				
				16	191(8)	626				
				18	170(7)	599				
$48^3$	20	153(6)	582							

## Results I

Gluonic (butterfly) operator  
+  
Gradient Flow Method



$\mathcal{Q}$

Massimo D'Elia's talk

$$\mathcal{L}_\theta = \frac{1}{4} F_{\mu\nu}^a(x) F_{\mu\nu}^a(x) - i\theta \frac{g^2}{64\pi^2} \epsilon_{\mu\nu\rho\sigma} F_{\mu\nu}^a(x) F_{\rho\sigma}^a(x),$$

$$\exp[-VF(\theta)] = \int [dA] \exp\left(-\int d^4x \mathcal{L}_\theta\right)$$

# Gradient flow

Lüscher, Lüscher Weisz

Evolve the link variables in a fictitious flow time:

$$\dot{V}_{x,\mu}(t) = -g_0^2 \left[ \partial_{x,\mu} S_{\text{Wilson}}(V(t)) \right] V_{x,\mu}(t);$$

Monitor  $\langle E \rangle = \frac{1}{2N_\tau N_\sigma^3} \sum_{x,\mu,\nu} \text{Tr}[F_{\mu\nu}(x) F^{\mu\nu}(x)]$  as a function of  $t$

Stop flowing when  $t^2 \langle E \rangle \big|_{t=t_0} = 0.3$

Observables  $\langle O(t) \rangle$  renormalized at  $\mu = 1/\sqrt{8t}$



Continuum limit of  $\langle O(t) \rangle$  is independent on the chosen reference value


Caveat: note comments by Kanaya et al.

# Distribution of the topological charge $P(Q)$


cluster around integers as cooling proceeds


(results for  $a = 0.06$  fm)

T=153 MeV



T=253 MeV



$P(Q)$  $Q$ 

$$P(Q) \text{ and } F(\theta)$$

Taylor expansion, and cumulants of the topological charge distribution

$$e^{-F(\theta)} = \langle e^{i\theta Q} \rangle$$

$$P_\nu = \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} e^{-i\theta\nu} e^{-F(\theta)} \quad Q = \nu$$

$$C_n = (-1)^{n+1} \frac{1}{V} \frac{d^{2n}}{d\theta^{2n}} F(\theta) \Big|_{\theta=0} \equiv \langle Q^{2n} \rangle_{conn}$$

$$F(\theta) = V \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\theta^{2n}}{(2n)!} C_n$$

$$P_\nu = \frac{e^{-\frac{\nu^2}{2\sigma^2}}}{\sqrt{2\pi\sigma^2}} \left[ 1 + \frac{1}{4!} \frac{\tau}{\sigma^2} \text{He}_4(\nu/\sigma) \right]$$

$\sigma^2 = VC_1$  and  $\tau = C_2/C_1$   $P(Q)$  is Gaussian for  $V \rightarrow \infty$

$F(\theta)$  is 'hidden' in  $P(Q)$ 's cumulants




In practice only the first two cumulants are accessible:

$$F(\theta, T) = 1/2 \chi(T) \theta^2 s(\theta, T)$$

$$s(\theta, T) = 1 + b_2(T) \theta^2 + \dots$$

$$b_2 = - \frac{\langle Q^4 \rangle - 3 \langle Q^2 \rangle^2}{12 \langle Q^2 \rangle}$$



Taylor coefficients of  $F(\theta, T)$


DIGA — at very high temperature — predicts

$$F(\theta, T) - F(0, T) = \chi(T)(1 - \cos(\theta)) \longrightarrow b_2 = -1/12$$





# Flowing towards the plateau

$$t^2 \langle E \rangle |_{t=t_x, x=0-6} = (0.3, 0.66, 0.1, 0.15, 0.2, 0.25, 0.45)$$





# On finer lattices, plateau is almost reached:

Gradient method coincides with cooling



# Results for the topological susceptibility for $M_\pi = 270$ MeV




Continuum limit:

- in principle independent on flow limit
- we need to interpolate results at fixed scale to match T

$$\chi(T, m_\pi) = \lim_{a \rightarrow 0} \chi^{1/4}(T, a, m_\pi, t_x)$$

$$\chi^{1/4}(T, a, m_\pi, t_x) = \chi^{1/4}(T, m_\pi) + a^2 k(T, t_x)$$




# Continuum results for $m_\pi = 370$ MeV



$$\chi(T)^{0.25} \simeq aT^{-0.26} \simeq T, T > 200 \text{ MeV}$$




# Detailed analysis for $T > 200$ MeV (use approx. linearity) - **1**

(In)dependence of continuum limit on flow's limit: 0.3 OK



# Detailed analysis for $T > 200$ MeV


Interpolation ok.









A mass rescaling appears to work nicely

**Bonati et al. 2016**



However: there is  
no mass  
dependence..






# Possible explanation : strong scaling violations

Comparison with BNL results



numerical data courtesy *S. Sharma*



Indication of corrections to  $a^2$  scaling on our two coarser lattices

# Comparison with BNL results (contn'd)

numerical data courtesy *S. Sharma*




Consistent trend for other temperatures: on our finer lattice the corrections to  $a^2$  scaling seem moderate

# Instanton potential - cumulants' ratio b2


DIGA predicts

$$F(\theta, T) - F(0, T) = \chi(T)(1 - \cos(\theta)) \longrightarrow b_2 = -1/12$$



$$b_2 = -1/12$$

DIGA limit for  $T > 350$  MeV



Consistent with Bonati et al.

Results II

Fermionic operator

$$n_L - n_R = Q_{top}$$


$$m \int d^4x \bar{\psi} \gamma_5 \psi = Q_{top}$$

# Topological and chiral susceptibility

Kogut, Lagaë, Sinclair 1999

HotQCD, 2012

$$\chi_{top} = \langle Q_{top}^2 \rangle / V = m_l^2 \chi_{5,disc}$$




$$\chi_\pi - \chi_\delta = \chi_{disc} = \chi_{5,disc}, \quad \text{for } T \geq T_c, m_l \rightarrow 0.$$

$$\chi_{top} = \langle Q_{top}^2 \rangle / V = m_l^2 \chi_{disc}$$


$$T > T_{U(1)_A} \simeq T_c$$

# Chiral susceptibility


Pion mass 370 MeV




Pion mass 470 MeV




Within errors, no discernable spacing dependence








# Results for physical pion mass




# Comparison with BNL results including fermionic results

numerical data courtesy *S. Sharma*

199 < T < 210 MeV




333 < T < 350 MeV




*dotted lines to guide the eye*

Effective exponent :  $d(T) = -T \frac{d}{dT} \ln \chi^{0.25}(T)$

$$\chi^{0.25}(T) = aT^{-d(T)}$$



Possibly consistent with instant-dyon?  
Shuryak 2017





Faster decrease before DIGA sets in

Effective exponent :

$$\chi_{top}^{1/4} = aT^{-d(T)}$$


Same DIGA onset seen in  $b_2 \approx 350$  MeV




Effective exponent  $d(T)$ :

$$\chi_{top}^{1/4} = aT^{-d(T)}$$

Comparisons with other results :





# Summary and open points I

## *-Gluonic operator with gradient flow method:*

Strong lattice artifacts for  $a > 0.06$  fm. The results for  $a = 0.06$  compare well with BNL results, where  $a^2$  corrections are still visible. No reliable continuum limit for the topological susceptibility.

$b_2$  is approaching the DIGA value for  $T > 300$  MeV on all the lattices, possibly due to a cancellation of lattice artifacts

## *-Fermionic operator:*

Residual lattice artifacts below statistical errors, allowing a continuum limit estimate. The results for  $T > 300$  are broadly consistent with others once rescaled to the physical pion mass, and confirm the DIGA behavior

We observe a faster decrease closer to  $T_c$ , in agreement with recent instanton-dyons predictions. This feature has not been seen in other studies

## Summary and open points II

### *-What next for Topology and QGP phenomenology*

- All in all, there is an emerging evidence that the QGP behaves as a DIGA for  $T > 300$  MeV, but such evidence only comes from the exponent and  $b_2$ . Can this agreement be accidental?

The behavior around  $T_c$  is still under scrutiny, and should be clarified to better understand the approach to DIGA, and the nature of the medium produced at the LHC.

### *-What next for the lattice*

Twisted mass Wilson fermions seem to perform well for topology: very little spacing effects for the fermionic operator, access to the cumulants even on coarse lattices.

Needless to say, simulations for smaller masses, and finer lattices would be most useful, and in view of the positive features of these fermions very worthwhile. The disconnected susceptibilities should be measured as well.

- *Dirac Spectrum analysis, FRG calculations (exp. for  $U(1)$  axial symmetry?)???*