

Temperature dependence of QGP shear viscosity within lattice SU(3) – gluodynamics

N.Yu. Astrakhantsev, V.V. Braguta, A.Yu. Kotov




Hadronic Matter under Extreme Conditions 2016

1 November, 2016

Outline

- ▶ Introduction
- ▶ Details of the calculation
- ▶ Fitting of the data
- ▶ Backus-Gilbert method
- ▶ Conclusion

Introduction



- ▶ Elliptic flow from STAR experiment (Nucl. Phys. A 757, 102 (2005))

$$\frac{dN}{d\phi} \sim (1 + 2v_1 \cos(\phi) + 2v_2 \cos^2(\phi)), \text{ } \phi - \text{scattering angle}$$

- ▶ Quark-gluon plasma is close to ideal liquid ($\frac{\eta}{s} = (1 - 3)\frac{1}{4\pi}$)

M. Luzum and P. Romatschke, Phys. Rev. C 78, 034915 (2008)

Other works

SU(3) gluodynamics:

- ▶ Karsch, F. et al., Phys.Rev. D35 (1987)
- ▶ A. Nakamura, S. Sakai, Phys. Rev. Lett. 94, 072305 (2005)
- ▶ H. B. Meyer, Phys.Rev. D76 (2007) 101701
- ▶ H. B. Meyer, Nucl.Phys. A830 (2009) 641C-648C

Results:

- ▶ $\frac{\eta}{s} = 0.134 \pm 0.033$ ($T/T_c = 1.65, 8 \times 28^3$)
- ▶ $\frac{\eta}{s} = 0.102 \pm 0.056$ ($T/T_c = 1.24, 8 \times 28^3$)
- ▶ $\frac{\eta}{s} = 0.20 \pm 0.03$ ($T/T_c = 1.58, 16 \times 48^3$)
- ▶ $\frac{\eta}{s} = 0.26 \pm 0.03$ ($T/T_c = 2.32, 16 \times 48^3$)

SU(2) gluodynamics:

- ▶ $\frac{\eta}{s} = 0.134 \pm 0.057$ ($T/T_c = 1.2, 16 \times 32^3$)

N.Yu. Astrakhantsev, V.V. Braguta, A.Yu. Kotov, JHEP 1509 (2015) 082

Lattice calculation of shear viscosity

Lattice calculation of shear viscosity

The first step:

Measurement of the correlation function:

$$C(t) = \langle T_{12}(t) T_{12}(0) \rangle$$

Lattice calculation of shear viscosity

The first step:

Measurement of the correlation function:

$$C(t) = \langle T_{12}(t) T_{12}(0) \rangle$$

The second step:

Calculation of the spectral function $\rho(\omega)$:


$$C(t) = \int_0^\infty d\omega \rho(\omega) \frac{\text{ch}\left(\frac{\omega}{2T} - \omega t\right)}{\text{sh}\left(\frac{\omega}{2T}\right)}$$

$$\eta = \pi \lim_{\omega \rightarrow 0} \frac{\rho(\omega)}{\omega}$$

Details of the calculation

- ▶ SU(3) gluodynamics
- ▶ Two-level algorithm
- ▶ Lattice size $32^3 \times 16$
- ▶ Temperatures $T/T_c = 0.9, 0.925, 0.95, 1.0, 1.2, 1.35, 1.5$
- ▶ Accuracy $\sim 2 - 3\%$ at $t = \frac{1}{2T}$
- ▶ $\langle T_{12}(x) T_{12}(y) \rangle \sim (\langle T_{11}(x) T_{11}(y) \rangle - \langle T_{11}(x) T_{22}(y) \rangle)$
- ▶ Clover discretization for the $\hat{F}_{\mu\nu}$

Correlation functions



Spectral function

$$C(t) = \int_0^\infty d\omega \rho(\omega) \frac{ch\left(\frac{\omega}{2T} - \omega t\right)}{sh\left(\frac{\omega}{2T}\right)}$$

Properties of the spectral function:

- ▶ $\rho(\omega) \geq 0, \rho(-\omega) = -\rho(\omega)$
- ▶ Asymptotic freedom: $\rho(\omega)|_{\omega \rightarrow \infty}^{NLO} = \frac{1}{10} \frac{d_A}{(4\pi)^2} \omega^4 \left(1 - \frac{5N_c \alpha_s}{9\pi}\right)$
 $\sim 90\%$ of the total contribution $t = 1/(2T)$
- ▶ Hydrodynamics: $\rho(\omega)|_{\omega \rightarrow 0} = \frac{\eta}{\pi} \omega$

Spectral function

$$C(t) = \int_0^\infty d\omega \rho(\omega) \frac{ch\left(\frac{\omega}{2T} - \omega t\right)}{sh\left(\frac{\omega}{2T}\right)}$$


Properties of the spectral function:

- ▶ $\rho(\omega) \geq 0, \rho(-\omega) = -\rho(\omega)$
- ▶ Asymptotic freedom: $\rho(\omega)|_{\omega \rightarrow \infty}^{NLO} = \frac{1}{10} \frac{d_A}{(4\pi)^2} \omega^4 \left(1 - \frac{5N_c \alpha_s}{9\pi}\right)$
 $\sim 90\%$ of the total contribution $t = 1/(2T)$
- ▶ Hydrodynamics: $\rho(\omega)|_{\omega \rightarrow 0} = \frac{\eta}{\pi} \omega$

Ansatz for the spectral function (QCD sum rules motivation)

$$\rho(\omega) = \frac{\eta}{\pi} \omega \theta(\omega_0 - \omega) + A \rho_{lat}(\omega) \theta(\omega - \omega_0)$$


Lattice spectral function



Results (fitting procedure)

$$\rho(\omega) = \frac{\eta}{\pi}\omega\theta(\omega_0 - \omega) + A\rho_{lat}(\omega)\theta(\omega - \omega_0)$$

$$\chi^2/dof \sim 1, \ A \sim 1, \ \omega_0/T \sim 7 - 8$$



Properties of the spectral function

- ▶ Hydrodynamical approximation works well up to $\omega < \pi T \sim 1\text{GeV}$ (H.B. Meyer, arXiv:0809.5202)

Properties of the spectral function

- ▶ Hydrodynamical approximation works well up to $\omega < \pi T \sim 1 \text{ GeV}$ (H.B. Meyer, arXiv:0809.5202)
- ▶ Asymptotic freedom works well from $\omega > 3 \text{ GeV}$

Properties of the spectral function

- ▶ Hydrodynamical approximation works well up to $\omega < \pi T \sim 1 \text{ GeV}$ (H.B. Meyer, arXiv:0809.5202)
- ▶ Asymptotic freedom works well from $\omega > 3 \text{ GeV}$
- ▶ Poor knowledge of the spectral function in the region $\omega \in (1, 3) \text{ GeV}$
⇒ Main source of uncertainty in the fitting procedure

Backus-Gilbert method for the spectral function

- ▶ Problem: find $\rho(\omega)$ from the integral equation

$$C(x_i) = \int_0^\infty d\omega \rho(\omega) K(x_i, \omega), \quad K(x_i, \omega) = \frac{ch\left(\frac{\omega}{2T} - \omega x_i\right)}{sh\left(\frac{\omega}{2T}\right)}$$

- ▶ Define an estimator $\tilde{\rho}(\bar{\omega})$ ($\delta(\bar{\omega}, \omega)$ - resolution function):

$$\tilde{\rho}(\bar{\omega}) = \int_0^\infty d\omega \hat{\delta}(\bar{\omega}, \omega) \rho(\omega)$$

- ▶ Let us expand $\delta(\bar{\omega}, \omega)$ as

$$\delta(\bar{\omega}, \omega) = \sum_i b_i(\bar{\omega}) K(x_i, \omega) \quad \tilde{\rho}(\bar{\omega}) = \sum_i b_i(\bar{\omega}) C(x_i)$$

- ▶ Goal: minimize the width of the resolution function


$$b_i(\bar{\omega}) = \frac{\sum_j W_{ij}^{-1} R_j}{\sum_{ij} R_i W_{ij}^{-1} R_j},$$

$$W_{ij} = \int d\omega K(x_i, \omega) (\omega - \bar{\omega})^2 K(x_j, \omega), R_i = \int d\omega K(x_i, \omega)$$

- ▶ Regularization by the covariance matrix S_{ij} :

$$W_{ij} \rightarrow \lambda W_{ij} + (1 - \lambda) S_{ij}, \quad 0 < \lambda < 1$$

Resolution function $\delta(0, \omega)$ ($T/T_c = 1$, $\lambda = 0.001$)




- ▶ Width of the resolution function $\omega/T \sim 4$
- ▶ Hydrodynamical approximation works up to $\omega/T < \pi$
- ▶ Problem: large contribution from ultraviolet tail ($\sim 50\%$)


Solution:

- Take ultraviolet contribution in the form:

$$\rho_{ultr} = A \rho_{lat}(\omega) \theta(\omega - \omega_0)$$


- Determine the value of the A from the $C(t)$ at small t
- Subtract ultraviolet contribution and obtain η/s as a function of ω_0






- ▶ For $T/T_c = 1$. $\omega_0/T = 7.33 \pm 0.47$
- ▶ $\eta/s = 0.22 \pm 0.04$

Preliminary results



Conclusion



- ▶ We calculated η/s for $T/T_c \in (0.9, 1.5)$
- ▶ Applied fitting procedure and Backus-Gilbert method for extracting SF
- ▶ η/s is close to N=4 SYM and in agreement with experiment