

НЕПЕРЕНОРМИРУЕМЫЕ КАК ПЕРЕНОРМИРУЕМЫЕ ...

**Д.КАЗАКОВ
БЛТФ ОИЯИ**

Motivation:

- The Standard Model is renormalizable
- Gravity is not renormalizable

Non-renormalizable theories are not accepted due to:

- UV divergences are not under control - infinite number of new types of divergences
- The amplitudes increase with energy (in PT) and violate unitarity

Suggestion (novel approach to NR interactions):

- To replace the multiplicative renormalization procedure by a new one, where the renormalization constant Z is replaced by an operator \hat{Z} , which depends on kinematics
- To sum up the leading asymptotics in all orders of PT (generalized RG) and to study the high-energy behavior

Renormalization

Consider 2->2 scattering amplitude on shell

$$\Gamma_4(s, t, u) = \Gamma_4^{tree} \bar{\Gamma}_4(s, t, u)$$

$$\bar{\Gamma}_4 = 1 + \lambda \dots + \lambda^2 \dots$$

Renormalization (dimensional regularization)

$$\bar{\Gamma}_4 = Z_4(\lambda) \bar{\Gamma}_4^{bare} |_{\lambda_{bare} \rightarrow \lambda Z_4},$$

$$\lambda_{bare} = \mu^\epsilon Z_4(\lambda) \lambda$$

BPHZ R-operation

$$RG = (1 - K)R'G$$

$$Z = 1 - \sum_i KR'G_i$$

In NR theories $Z \rightarrow \hat{Z}$ \hat{Z} is a function (polynomial) of s,t,u acting as an operator

Example (taken from D=8 YM theory)

Exactly follows the BPHZ R-operation

$$\hat{Z} = 1 + g^2 \frac{st}{\epsilon}$$

$$g^2 s t \square \implies g^2 \left(s \triangle + t \nabla \right)$$

Either s or t are to be inserted into the loop and integrated

$$\phi_D^4$$

$$D = 4, 6, 8, 10$$

$$[\lambda] = 2 - D/2$$

2->2 scattering amplitude on shell


$$m = 0$$

$$s + t + u = 0$$

$$\Gamma_4(s, t, u) = \lambda(1 + \Gamma_s(s, t, u) + \Gamma_t(s, t, u) + \Gamma_u(s, t, u))$$

PT:

$$\Gamma_s = \sum_{n=1}^{\infty} (-z)^n S_n, \quad \Gamma_t = \sum_{n=1}^{\infty} (-z)^n T_n, \quad \Gamma_u = \sum_{n=1}^{\infty} (-z)^n U_n, \quad z \equiv \frac{\lambda}{\epsilon}$$



PT expansion (only s-channel is shown)

BPHZ R-operation

$$\mathcal{R}'G_n = \frac{A_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^n} + \frac{A_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^n} + \dots + \frac{A_1^{(n)} (\mu^2)^\epsilon}{\epsilon^n} + \text{lower pole terms}$$

$A_k^{(n)} (\mu^2)^{k\epsilon}$ terms appear after subtraction of (n-k) loop counter terms

Statement: $R'G_n$ is local, i.e. terms like $\log^k \mu^2 / \epsilon^m$ should cancel for any k and m


Consequence: $A_n^{(n)} = (-1)^{n+1} \frac{A_1^{(n)}}{n}$

$$KR'G_n = \sum_{k=1}^n \left(\frac{A_k^{(n)}}{\epsilon^n} \right) \equiv \frac{A_n^{(n)'}}{\epsilon^n} \quad A_n^{(n)'} = (-1)^{n+1} A_n^{(n)} = \frac{A_1^{(n)}}{n}.$$

$A_1^{(n)}$ is the contribution to the leading pole in n-loops from the diagrams appearing in due course of R-operation after subtraction of (n-1) loop counter terms

The leading divergences are governed by 1 loop diagrams!

Recurrence Relations for the Leading Poles



$$\begin{aligned}
 nS_n(s, t, u) &= \frac{s^{D/2-2}}{\Gamma(D/2-1)} \int_0^1 dx [x(1-x)]^{D/2-2} (S_{n-1}(s, t', u') + T_{n-1}(s, t', u') + U_{n-1}(s, t', u')) \\
 &+ \frac{1}{2} \frac{s^{D/2-2}}{\Gamma(D/2-1)} \int_0^1 dx [x(1-x)]^{D/2-2} \sum_{k=1}^{n-2} \sum_{p=0}^{(D/2-2)k} \sum_{l=0}^p \frac{1}{p!(p+D/2-2)!} \times \\
 &\times \frac{d^p}{dt'^l du'^{p-l}} (S_k + T_k + U_k) \frac{d^p}{dt'^l du'^{p-l}} (S_{n-k-1} + T_{n-k-1} + U_{n-k-1}) s^p [x(1-x)]^p t^l u^{p-l}
 \end{aligned}$$

$$t' = -xs, u' = -(1-x)s$$

Solution of Recurrence Relations

Starting point:




$$S_1 = \frac{1}{2} \frac{\Gamma(D/2 - 1)}{\Gamma(D - 2)} s^{D/2-2}, \quad \text{etc}$$



Use the recurrence relation

$$S_2 = \frac{1}{4} \frac{\Gamma(D/2 - 1)}{\Gamma(D - 2)} \left[\frac{\Gamma(D/2 - 1)}{\Gamma(D - 2)} + 2(-)^{D/2} \frac{\Gamma(D - 3)}{\Gamma(3D/2 - 4)} \right] s^{D-4}, \quad \text{etc}$$



Notice the difference with renormalizable theory: S_1 depends on kinematics!

To get S_2 one has to integrate T_1 and U_1 over the loop

This is exactly what we do in writing the recurrence relation

$$\frac{s^{D/2-2}}{\Gamma(D/2 - 1)} \int_0^1 dx [x(1-x)]^{D/2-2} (S_{n-1}(s, t', u') + T_{n-1}(s, t', u') + U_{n-1}(s, t', u'))$$

Differential Equation

Summing up the recurrence relation $\sum_{n=2}^{\infty} (-z)^n$ one gets the diff equation

$$\begin{aligned}
 -\frac{d\Gamma_s(s, t, u)}{dz} &= \frac{1}{2} \frac{\Gamma(D/2 - 1)}{\Gamma(D - 2)} s^{D/2-2} & \Gamma_s(z = 0) &= 0 \\
 + \frac{s^{D/2-2}}{\Gamma(D/2 - 1)} \int_0^1 dx [x(1-x)]^{D/2-2} & [\Gamma_s(s, t', u') + \Gamma_t(s, t', u') + \Gamma_u(s, t', u')] & \Big|_{\substack{t' = -xs, \\ u' = -(1-x)s}} & \\
 + \frac{1}{2} \frac{s^{D/2-2}}{\Gamma(D/2 - 1)} \int_0^1 dx [x(1-x)]^{D/2-2} & \sum_{p=0}^{\infty} \sum_{l=0}^p \frac{1}{p!(p + D/2 - 2)!} \times \\
 \times \left(\frac{d^p}{dt'^l du'^{p-l}} (\Gamma_s + \Gamma_t + \Gamma_u) \Big|_{\substack{t' = -xs, \\ u' = -(1-x)s}} \right)^2 & s^p [x(1-x)]^p t^l u^{p-l}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d\Gamma_s(s, t, u)}{d \log \mu^2} &= -\frac{\lambda}{2} \frac{s^{D/2-2}}{\Gamma(D/2 - 1)} \int_0^1 dx [x(1-x)]^{D/2-2} \sum_{p=0}^{\infty} \sum_{l=0}^p \frac{1}{p!(p + D/2 - 2)!} \times \\
 & \times \left(\frac{d^p \bar{\Gamma}_4(s, t', u')}{dt'^l du'^{p-l}} \Big|_{\substack{t' = -xs, \\ u' = -(1-x)s}} \right)^2 s^p [x(1-x)]^p t^l u^{p-l}
 \end{aligned}$$

$$\Gamma_s(\log \mu^2 = 0) = 0$$

Solution of RG Equation

$$D = 4$$

$$s \sim t \sim u \sim E^2$$

$$\frac{d\bar{\Gamma}_4}{d \log \mu^2} = -\lambda \frac{3}{2} \bar{\Gamma}_4^2, \quad \bar{\Gamma}_4(\log \mu^2 = 0) = 1 \quad \rightarrow \quad \bar{\Gamma}_4 = \frac{1}{1 + \frac{3}{2} \lambda \log(\mu^2 / E^2)}$$

General Solution for any D

$$\bar{\Gamma}_4(s, t, u) = \mathcal{P} \frac{1}{1 + \frac{1}{2} \frac{\Gamma(D/2-1)}{\Gamma(D-2)} \lambda (s^{D/2-2} + t^{D/2-2} + u^{D/2-2}) \log(\mu^2 / E^2)}$$

\mathcal{P} is the symbol of ordering in a sense of recurrence relation

To get S_2 for instance one has to take $s^{D/2-2}(s^{D/2-2} + t^{D/2-2} + u^{D/2-2})$ terms

and integrate s, t and u over the s-loop. This is exactly given by

$$\frac{s^{D/2-2}}{\Gamma(D/2-1)} \int_0^1 dx [x(1-x)]^{D/2-2} (S_{n-1}(s, t', u') + T_{n-1}(s, t', u') + U_{n-1}(s, t', u'))$$

High Energy Behaviour of the scattering amplitude in ϕ_D^4 theory

$$\Gamma_4(s, t, u) = \mathcal{P} \frac{\lambda}{1 + \frac{1}{2} \frac{\Gamma(D/2-1)}{\Gamma(D-2)} \lambda (s^{D/2-2} + t^{D/2-2} + u^{D/2-2}) \log(\mu^2/E^2)}$$

$$s \sim t \sim u \sim E^2$$

$D = 4$ $3/2 > 0$ As a result one has a Landau pole as $E \rightarrow \infty$

$D = 6$ $s + t + u = 0$ All the leading divergences (logs) cancel in all loops

One can explicitly check that S_2 given above vanishes

$D = 8$ $s^2 + t^2 + u^2 > 0$

$D = 10$ $s^3 + t^3 + u^3 = 3stu > 0$ $s > 0, t, u < 0$

Conclusion: ϕ_D^4 has a Landau pole as $E \rightarrow \infty$