

Мемориальная конференция

памяти академика

Андрея Алексеевича Славнова

21–22 декабря 2022 г., Москва



# QUANTUM CORRECTIONS TO EFFECTIVE POTENTIAL IN ARBITRARY SCALAR THEORIES

**D.KAZAKOV**

In collaboration with R. Iakhibbaev and D. Tolkachev



**BLTP JINR**

22 December 2022



# Effective Potential in Scalar Theory in D=4

Generating functional for Green functions

$$Z(J) = \int \mathcal{D}\phi \exp \left( i \int d^4x \mathcal{L}(\phi, d\phi) + J\phi \right)$$

$$W(J) = -i \log Z(J) \quad \text{IPI generating functional}$$

Effective action

$$\Gamma(\phi) = W(J) - \int d^4x J(x)\phi(x) \quad \text{Legendre transformation}$$

$$e^{i\Gamma(\hat{\phi})} = \int \mathcal{D}\phi e^{i(S[\hat{\phi}+\phi] - \phi S'[\hat{\phi}])}$$

Shifted Classical action

$$S[\hat{\phi} + \phi] = S[\hat{\phi}] + \phi S'[\hat{\phi}] + \frac{1}{2} \phi^2 S''[\hat{\phi}] + \frac{1}{3!} \phi^3 S'''[\hat{\phi}] + \dots$$

↑  
Classical external field

↑  
Field dependent mass


↑  
Interaction vertex

# Effective Potential in Scalar Theory in D=4

General scalar field theory in D=4

$$\mathcal{L} = \frac{1}{2}(\partial_\mu\phi)^2 - gV_0(\phi)$$

Vacuum diagrams



$$v_2(\phi) \equiv \frac{d^2V_0(\phi)}{d\phi^2}$$

$$v_n \equiv d^nV_0/d\phi^n$$

One loop

$$v_2^2 \frac{1}{\epsilon}$$

$$\phi^4$$

$$v_2 \sim \phi^2$$

$$\phi^4 \frac{1}{\epsilon}$$

Two loops

$$\phi^6$$

$$v_2 \sim \phi^4$$

$$\phi^8 \frac{1}{\epsilon}$$

$$(v_2^2 v_4 + v_3^2 v_2)/\epsilon^2$$

$$\phi^4$$

$$v_4 \sim 1, v_3 \sim \phi$$

$$\phi^4/\epsilon^2$$

$$\phi^6$$

$$v_4 \sim \phi^2, v_3 \sim \phi^3$$

$$\phi^{10}/\epsilon^2$$

# Divergences and Log $\phi$ behaviour



$$Diag \sim \frac{1}{\epsilon} v_2^2 \left( \frac{\mu^2}{m^2} \right)^\epsilon \rightarrow v_2^2 \left( \frac{1}{\epsilon} - \log \frac{m^2}{\mu^2} \right), \quad m^2 = gv_2^2$$

$$\Delta V_1 = \frac{g^2}{16\pi^2} v_2^2 \log \frac{gv_2}{\mu^2}$$


$$\phi^4 \quad \Delta V_1 = \frac{g^2}{16\pi^2} \phi^4 \log \frac{g\phi^2}{\mu^2}$$

$$\phi^6 \quad \Delta V_1 = \frac{g^2}{16\pi^2} \phi^8 \log \frac{g\phi^4}{\mu^2}$$

# BPHZ R-operation


$$\mathcal{R}'G_n = \frac{A_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^n} + \frac{A_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^n} + \dots + \frac{A_1^{(n)} (\mu^2)^\epsilon}{\epsilon^n}$$

lower pole terms



$A_k^{(n)} (\mu^2)^{k\epsilon}$  terms appear after subtraction of (n-k) loop counter terms


Statement:  $R'G_n$  is local, i.e. terms like  $\log^k \mu^2 / \epsilon^m$  should cancel for any k and m

Consequence:  $A_n^{(n)} = (-1)^{n+1} \frac{A_1^{(n)}}{n}$  

The leading divergences are governed by 1 loop diagrams!

# Two loop example

 $\phi^4_4$ 



$$= \left( \frac{A_2^{(2)}}{\epsilon^2} + \frac{A_1^{(2)}}{\epsilon} \right) \left( \frac{\mu^2}{s} \right)^{2\epsilon}$$

$$\mathcal{R}' \left[ \text{triangle with bubble} \right] = \left[ \text{triangle with bubble} \right] - \left[ \text{fish diagram} \right] \cdot \left[ \text{tadpole} \right] = \left( \frac{A_2^{(2)}}{\epsilon^2} + \frac{A_1^{(2)}}{\epsilon} \right) \left( \frac{\mu^2}{s} \right)^{2\epsilon} - \frac{A_1^{(1)}}{\epsilon} \left( \frac{\mu^2}{s} \right)^\epsilon \frac{A_1^{(1)}}{\epsilon}$$

$$= \frac{A_2^{(2)}}{\epsilon^2} - \frac{(A_1^{(1)})^2}{\epsilon^2} + \underbrace{2 \frac{A_2^{(2)}}{\epsilon} \log(\mu^2/s) - \frac{(A_1^{(1)})^2}{\epsilon} \log(\mu^2/s)}_{\text{non-local terms to be cancelled}} = -\frac{1}{2} \frac{(A_1^{(1)})^2}{\epsilon^2} + \dots$$

non-local terms to be cancelled

Leading divergence is given by the one-loop term

$$A_2^{(2)} = \frac{1}{2} (A_1^{(1)})^2$$

- These statements are universal and are valid in non-renormalizable theories as well.
- The only difference is that the counter term  $A_1^{(1)}$  depends on kinematics and has to be integrated through the remaining one-loop graph.
- As a result  $A_2^{(2)}$  is not the square of  $A_1^{(1)}$  anymore but is the integrated square .
- This last statement is the general feature of any QFT irrespective of renormalizability

# Divergences and Log $\phi$ behaviour



$$Diag \sim \frac{1}{\epsilon} v_2^2 \left( \frac{\mu^2}{m^2} \right)^\epsilon \rightarrow v_2^2 \left( \frac{1}{\epsilon} - \log \frac{m^2}{\mu^2} \right), \quad m^2 = gv_2^2$$

The leading divergences



The leading logs


- In non-renormalizable theories divergences cannot be absorbed into the renormalization of the couplings and fields.
- If they are subtracted some way one is left with infinite arbitrariness.
- Coefficients of the leading divergences (logs) do not depend on this arbitrariness !

The aim is to calculate the leading divergences  $\sim \frac{1}{\epsilon^n}$  in n-th order of PT

# Recurrence relations for the leading poles

Kazakov, Iakhibbaev, Tolkachev 22

Action of  $R'$ -operation on divergent diagram



$$nS_n = \frac{1}{2}v_2 D_2 S_{n-1} + \frac{1}{4} \sum_{k=1}^{n-2} D_2 S_k D_2 S_{n-1-k}, \quad n \geq 2 \quad S_1 = \frac{1}{4}v_2^2$$

$$nS_n = \frac{1}{4} \sum_{k=0}^{n-1} D_2 S_k D_2 S_{n-1-k}, \quad n \geq 1, \quad S_0 = V_0$$



## RG pole equation for arbitrary potential

$$\Sigma(z, \phi) = \sum_{n=0}^{\infty} (-z)^n S_n(\phi) \quad z = \frac{g}{\epsilon}$$

RG pole equation

$$\frac{d\Sigma}{dz} = -\frac{1}{4} (D_2 \Sigma)^2 \quad \Sigma(0, \phi) = V_0(\phi)$$

This a non-linear partial differential equation!

Effective potential

$$V_{eff}(g, \phi) = g \Sigma(z, \phi) \Big|_{z \rightarrow -\frac{g}{16\pi^2} \log gv_2/\mu^2} \quad v_2(\phi) \equiv \frac{d^2 V_0(\phi)}{d\phi^2}$$

# Example I: Power like Potential

$$gV_0(\phi) = g \frac{\phi^p}{p!} \quad y = g\phi^{p-4} \quad \Sigma(z, \phi) = \frac{\phi^p}{p!} f(z\phi^{p-4})$$


$$f'(y) = -\frac{1}{4p!} [p(p-1)f(y) + (p-4)(3p-5)yf'(y) + (p-4)^2y^2f''(y)]^2$$

$$f(0) = 1, f'(0) = -\frac{1}{4} \frac{p(p-1)}{(p-2)!}$$

**p=4**

$$f'(y) = -\frac{3}{2}f(y)^2 \quad f(y) = \frac{1}{1 + \frac{3}{2}y}$$

$$V_{eff}(\phi) = \frac{g\phi^4/4!}{1 - \frac{3}{2} \frac{g}{16\pi^2} \log\left(\frac{g\phi^2}{2\mu^2}\right)}$$



# Example I: Power like Potential

$p > 4$


$$gV_0(\phi) = g \frac{\phi^p}{p!}$$

$$f(y) = u(y)/y$$


$$yu'(y) - u(y) = -\frac{1}{4p!} [12u(y) + (p-4)(p+3)yu'(y) + (p-4)^2y^2u''(y)]^2$$

$$u(\pm 0) = 0, u'(\pm 0) = \pm 1$$

Discontinuity at  $y=0$



$y = 0$




$y = 0$


$$y \rightarrow -\frac{g}{16\pi^2} \phi^{p-4} \log \frac{g\phi^{p-2}}{\mu^2/(p-2)!}$$

# Example I: Power like Potential

$p=5$



$p=6$



- Finite gap instead of an infinite barrier as for  $p=4$
- Metastability of the quantum state
- No new minima

# Applicability of approximation

Validity of PT

$$\frac{g}{16\pi^2} \phi^{p-4} < 1$$

Validity of LL approximation

$$\log \frac{g\phi^{p-2}}{(p-2)!\mu^2} > 1$$

Possible simultaneously for small coupling  $g$  and temporal field  $\phi$

Singular point  $\frac{g\phi^{p-2}}{(p-2)!\mu^2} = 1$  is within validity region

$$p=6 \quad \phi^2 < \frac{16\pi^2}{g} \implies \log \frac{g\phi^4}{4!\mu^2} < \log \frac{(16\pi^2)^2}{24g\mu^2} > 1$$

$$g\mu^2 < 1$$


$$y \rightarrow -\frac{g}{16\pi^2} \phi^{p-4} \log \frac{g\phi^{p-2}}{\mu^2/(p-2)!}$$

# Example II: Exponential Potential

$$gV_0 = ge^{|\phi/m|}$$

$$y = g/m^4 e^{\phi/m}$$

$$f'(y) = -\frac{1}{4} (y^2 f''(y) + 3yf'(y) + f(y))^2$$



$$y \rightarrow \frac{g}{16\pi^2 m^4} e^{|\phi/m|} \log \left[ \frac{g}{16\pi^2 m^2 \mu^2} e^{|\phi/m|} \right]$$

- Finite gap
- Metastability of the quantum state
- No new minima

Applicability  $\frac{g}{16\pi^2 m^4} e^{|\phi/m|} < 1$

$$\log \left[ \frac{g}{m^2 \mu^2} e^{|\phi/m|} \right] > 1$$



# Example III: Inflation Potential

$$gV_0 = g \tanh^2(\phi/m)$$

$$\frac{d\Sigma}{dz} = -\frac{1}{4} \left( \frac{d^2\Sigma}{d\phi^2} \right)^2 \quad \Sigma\left(\frac{z}{m^4}, \frac{\phi}{m}\right) \quad \Sigma|_{\phi \rightarrow \infty} \rightarrow 1 \quad z = \frac{g}{\epsilon}$$

$$\Sigma'|_{\phi \rightarrow \infty} \rightarrow 0$$

$$V_{eff}(g, \phi) = g\Sigma(z, \phi)|_{z \rightarrow -\frac{g}{16\pi^2} \log gv_2/\mu^2} \cdot \quad v_2(\phi) \equiv \frac{d^2V_0(\phi)}{d\phi^2}$$



- Peak at the origin
- Additional minima

# Conclusion on Effective potential

- The effective potential in the LL approximation obeys the RG master equation which is a partial non-linear differential equation
  - In some cases this equation is simplified to the ordinary differential one and can be solved at least numerically. I
  - In all the cases that we studied the obtained ordinary differential equations obey the solution with a discontinuity.
  - The effective potential has a metastable minima at the origin and no other minima exists.
- 
- The main message is that under certain assumptions while studying the CW mechanism one may not be restricted by the renormalizable potentials but consider much wider possibilities. We provided the method of such analysis.
  - This might be useful for cosmological applications where they are usually not limited by renormalizability since gravity makes it non-renormalizable anyway.