A Feasibility Experiment at RHIC to Measure the Analyzing Power for Drell-Yan Production (A_NDY)



L. Nogach, IHEP (Protvino)
for A_NDY

DSPIN-2011, September 20-24

AnDY

"Large Rapidity Drell -Yan Production at RHIC"

Letter of Intent submitted 24 May 2010:

http://www.bnl.gov/npp/docs/pac0610/Crawford_LoI.100524.v1.pdf

PAC presentation:

http://www.bnl.gov/npp/docs/pac0610/aschenauer_DY-collider_june10.pdf

Proposal to 2011 PAC:

http://www.bnl.gov/npp/docs/pac0611/DY_pro_110516_final.2.pdf

E.C. Aschenauer, A. Bazilevsky, L.C. Bland, K. Drees, C. Folz, Y. Makdisi, A. Ogawa, P. Pile, T.G. Throwe
Brookhaven National Laboratory
H.J. Crawford, J.M. Engelage, E.G. Judd
University of California, Berkeley / Space Sciences Laboratory
C.W. Perkins
University of California, Berkeley / Space Sciences Laboratory / Stony Brook University
A. Derevshchikov, N. Minaev, D. Morozov, L. Nogach
Institute for High Energy Physics, Protvino
G. Igo
University of California, Los Angeles
M. Grosse Perdekamp
University of Illinois
M.X. Liu
Los Alamos National Laboratory
H. Avakian
Thomas Jefferson National Accelerator Facility
E.J. Brash
Christopher Newport University and TJNAF
C.F. Perdrisat
College of William and Mary
V. Punjabi
Norfolk State University
X. Li
Shandong University, China
M. Planinic, G. Simatovic
University of Zagreb, Croatia
A. Vossen
Indiana University
G. Schnell, C. van Hulse
University of the Basque Country and IKERBASQUE, Spain
A. Shahinyan, S. Abrahamyan
Yerevan Physics Institute

RHIC as a polarized proton collider



Transverse single spin asymmetry measured at RHIC



B.I.Abelev (STAR), Phys. Rev. Lett. 101 (2008)

U. D'Alesio, F. Murgia,
Phys. Rev. D 70, 074009 (2004)

C. Kouvaris, J. Qiu, W. Vogelsang, F. Yuan,
Phys. Rev. D 74, 114013 (2006)

Significant spin effects are found at x_F>0.3; the x-range does not overlap with SIDIS

Theory prediction for Drell-Yan process

- Sivers function: [illegible]
- In QCD

DIS: attractive



Drell-Yan: repulsive



As a result:

$$\mathrm{Sivers}|_{\mathrm{DIS}} = -\mathrm{Sivers}|_{\mathrm{DY}}$$

Test of this relation is a test of TMD factorization

Theory prediction for DY – new development

- SIDIS and RHIC pion production do not overlap in momentum fraction (x)
- Attempts to describe both result in a sign "mismatch" conclusion [Kang, Qiu, Vogelsang, Yuan, PRD83 (2011) 094001]



Combined analysis of SIDIS and inclusive pion production leads to the conclusion that the u-quark Sivers function has a node at x~0.4

Essential to test predicted sign change for DY in same kinematics as SIDIS

A. Prokudin, Z.B. Kang, "Opportunities for Drell-Yan Physics at RHIC" workshop (May, 2011)

Motivation for a feasibility experiment at RHIC

- Forward upgrades of STAR and PHENIX are major undertakings and would benefit from a feasibility demonstration of forward DY production.
- Forward DY production is of interest for more than just the analyzing power, e.g. most robust observable to low-x parton distributions for intercomparison to a future electron-ion collider.
- A_NDY can run in parallel with RHIC W program.

A_NDY goals

- To establish that large-x_F low-mass dileptons from the DY process can be discriminated from background in $\sqrt{s}$=500 GeV p↑+p collisions.
- To provide sufficient statistical precision for the analyzing power for DY production to test the theoretical prediction of a sign change compared to transverse single spin asymmetries for semi inclusive deep inelastic scattering.

Requirements for DY

- Luminosity (Lint ~150/pb required)

- Background Reduction
  - electron/hadron discrimination
  - Charged/neutral discrimination and photon conversion background
  - Open heavy flavor (c, b) production
  - Is a magnet plus tracking required for forward DY?

Collision energy dependence of DY production



Comments…

- RHIC pp luminosity largest at √s=500 GeV
- partonic luminosities increase with √s
- net result is that DY grows with √s
- in any case, largest √s probes lowest x

⇒ Consider large-x_F DY at √s=500 GeV

$$\text{large } x_F \Rightarrow x_1 \approx x_F \text{ and } x_2 \approx \frac{M^2}{x_F s}$$

⇒ Forward DY production probes valence region for "beam" and $x_2 \approx 10^{-4}$ for "target" for √s=500 GeV (M>4 GeV/c^2)

Schematic of detector considered

Original plan for Run-11



Equipment in place:

- **Hcal is existing 2x9x12 modules from E864 (NIM406,227)**
- **BBC and ZDC**

Goal:

- **establish impact of 3 IR operation on PheniX and Star luminosity**
- **calibrate HCAL**
  - **absolute Energy scale with ρ, Φ, K_s**
  - **gains with cosmics**
- **measure the hadronic background to bench mark MCs further**

Schematic of detector considered

Run-12 configuration



- HCal is existing 9x12 modules from E864
- ECal is modeled as only $(3.8cm)^2$x(45cm) lead glass
- Preshower would require construction

PHOBOS split dipole expected to be in place, but not used

<u>Goal</u>: to acquire data sample to observe $J/\psi \rightarrow e+e-$, $\Upsilon \rightarrow e+e-$ and dilepton continuum between these signals as a benchmark for DY feasibility

Schematic of detector considered
Run-13 configuration



- HCal is existing 9x12 modules from E864
- ECal is modeled as only $(3.8cm)^2$x(45cm) lead glass
- Preshower would require construction
- PHOBOS split-dipole magnetic field in GEANT model
- Fiber tracker stations and MWPC require construction

Uses PHOBOS split dipole for charge sign

<u>Goal</u>: to acquire 2nd data sample with tracking to quantify the role of charge sign discrimination in suppressing backgrounds

Background estimate



Comments:

- Conversion photons significantly reduced by $\pi^0 \rightarrow \gamma\gamma$ veto
- Preshower thickness tuned, although perhaps is not so critical given photon veto
- Linearly decreasing dN/df (fast-simulation model for hadronic response of ECal) estimates smaller hadronic background $\Rightarrow$ increased sophistication needed for reliable estimates, although other model uncertainties could easily dominate
- Open heavy flavor backgrounds also estimated and found small due to large rapidity

Run-11 implementation



Schematic of detector for Run-11

Polarized proton collisions at √s=500 GeV from February to April 2011



- Beam-beam counter (BBC) for minimum-bias trigger and luminosity measurement from PHOBOS [NIM A474 (2001) 38]
- Zero-degree calorimeter and shower maximum detector for luminosity measurement and local polarimetry (ZDC/ZDC-SMD, not shown)
- Hadron calorimeter (HCal) are modules of 9x12 lead-scintilating fiber cells from AGS-E864 [NIM406, 227]
- Small ECal (~120 lead glass cells, $(4cm)^2$x40cm) loaned from BigCal at JLab
- Newly constructed preshower detector (two planes, 2.5 & 10 cm)

Impact of IP2 collisions



IP2 collisions have begun <3 hours after physics ON with minimal impact on IP6, IP8. Adequate luminosity for A_NDY (10 pb^{-1}/week for $\sqrt{s}$=500 GeV polarized proton collisions) is projected for subsequent runs.

Integrated luminosity in Run-11



- Presently, no accidentals correction (found to be large in some stores for ZDC). All rates are scaled by bunch-crossing scalers, following 20110330 access and subsequent timing adjustments.
- Effective BBC cross section has been revisited through full simulations and now is in better agreement with vernier scan measurements (fill=15457)
- Jet analyzing power measurement sets goal of 10/pb for run11

Many thanks to C-A for IP2 collisions

Calibration of Hadron Calorimeter

ADC spectra from 9x12 cells of HCal module for cosmic-ray triggers



- Relative gains – with cosmic ray muons
- LED system to monitor stability

Calibration of Hadron Calorimeter

Absolute energy scale - based on $\pi^0 \rightarrow \gamma\gamma$ reconstruction



- Require: (1) 1-tower clusters; (2) E>1.8 GeV; (3) |x|>50 cm to avoid ECal shadow; (4) >1 clusters to form pairs; (5) E_{pair}>5 GeV; and (6) z_{pair}<0.5.
- Apply to 20M minimum-bias events from run11 data
- Apply to 20M PYTHIA events subjected to BBC charge sum trigger emulation (no vertex cut)
- Data and simulations are both absolutely normalized, so PYTHIA is expected to provide a good basis for QCD backgrounds to DY.
- Hadronic corrections expected to be small. Mass reconstructions to demonstrate this are underway.

Jet Trigger



Hadron calorimeter is quiet ~107ns before jet event

- Jet trigger sums HCal response excluding outer two perimeters (rather than just two columns closest to beam)
- Definition is consistent with objective of having jet thrust axis centered in hadron calorimeter modules
- HCal energy scale is now determined
- >750M jet-triggered events acquired during RHIC run11

Hadron calorimeter is quiet again ~107ns after jet event

Towards Forward Jets





- Good agreement between data and PYTHIA/GEANT simulation for summed HCal response excluding outer two perimeters of cells ⇒ QCD backgrounds can be modeled
- Good agreement between data and simulation for jet shape
- Next up: forward jet analyzing power

Forward Jet Energy Scale





- Jet energy scale determined by association analysis of simulation
- Required to add ECal energy deposition to masked summed HCal response
- Small rescaling (<10%) of HCal energy scale is applied, likely to account for hadronic corrections

Status of A_NDY

- Calorimetry required for experiment is at BNL. It is on loan from JLab until July 2014.
- Layout for second stage (no magnet) is complete, with specification of azimuthally complete HCal, ECal and segmented preshower detector.
- Existing split dipole does require modification to match acceptance of final calorimetry plan.
- Staging of apparatus awaits funding review.
- At present, first attempt at transverse spin DY measurement will be in RHIC run13.



Projected sensitivity of A_NDY for L_{int}=100 pb^{-1} for $p^{\uparrow}p^{\uparrow}$ collisions at $\sqrt{s}$=500 GeV requiring γ^* M>4 GeV/c^2, p_T<2 GeV/c and $0.1<x_F<0.3$

Measurements of forward pion and jet production with small ECal might be possible in run12 at $\sqrt{s}$=200/500 GeV.

Backup

Run?? implementation



Proposed configuration of the A_NDY ECal as seen from IP2. The inner calorimeter is a 32-row × 32-column matrix of $(3.8\ \text{cm})^2 \times 45$ cm lead-glass detectors with a central 6-row × 4-column hole for the vacuum pipe for the beams. It is surrounded by $(40\ \text{mm})^2 \times 40$ cm lead-glass detectors. The calorimeter separates into two halves that can be translated to larger |x|.



y-z view at x=0 (top) and x-z view at y=0 (bottom) showing the relationship of the model=6 ECal and HCal. These schematic views show only the lead-glass of ECal and the active portion of HCal.

Previous Work on Low-Mass DY at a Collider

p+p DY at ISR, √s=53,63 GeV
Phys. Lett. B91 (1980) 475

STUDY OF MASSIVE ELECTRON PAIR PRODUCTION AT THE CERN INTERSECTING STORAGE RINGS

C. KOURKOUMELIS and L.K. RESVANIS
University of Athens, Athens, Greece

T.A. FILIPPAS and E. FOKITIS
National Technical University, Athens, Greece

A.M. CNOPS, J.H. COBB [1], R. HOGUE, S. IWATA [2], R.B. PALMER, D.C. RAHM, P. REHAK and I. STUMER
Brookhaven National Laboratory [3], Upton, NY, USA

C.W. FABJAN, T. FIELDS [4], D. LISSAUER [5], I. MANNELLI [6], P. MOUZOURAKIS, K. NAKAMURA [7], A. NAPPI [6], W. STRUCZINSKI [8] and W.J. WILLIS
CERN, Geneva, Switzerland

M. GOLDBERG, N. HORWITZ and G.C. MONETI
Syracuse University [9], Syracuse, NY, USA

and

A.J. LANKFORD [10]
Yale University, New Haven, CT, USA

Received 18 February 1980



Fig. 1. The cross section $(d^2\sigma/dm\,dy)_{y=0}$ versus mass for the data at $\sqrt{s}$ = 53 and 63 GeV combined. The curve is a result of the fit to the continuum displayed in fig. 2. The inset show the mass acceptance for "1977" and "1978" triggers and geometrical configurations calculated for isotropic decay distributions and production uniform in rapidity with p_T dependence $d\sigma/dp_T^2 \sim \exp(-bp_T)$, where b = 1.4 GeV^{-1}. The mass acceptance changes by ± 15% when the helicity decay distribution follows $dN/d\cos\theta = 1 + \alpha\cos^2\theta$ when $\alpha = \pm 1$, where θ is measured in the s-channel helicity frame.

Comments (note: large x_F at collider breaks new ground)...

- e+e- low-mass DY done at ISR and by UA2 [see review J.Phys. G19 (1993) D1]
- UA2 [PLB275 (1992) 202] did not use magnet / CCOR did [PLB79 (1979) 398]
- most fixed target experiments do μ+μ- DY

Pair mass from bare EMcal



arXiv:0906.2332



arXiv:0907.4396

- pair mass backgrounds well modeled
- J/ψ→e+e- observation with bare EMcal at $\langle x_F \rangle \sim 0.67$ emboldens DY consideration

Impact of Collisions at IP2

The anatomy of initiating collisions at IP2



Early after a RHIC store is set up, beams are colliding at IP6 (STAR) and IP8 (PHENIX). Beams are transversely separated at IP2 (A_NDY). The arrow indicates when collisions begin at IP2



After the beam intensity decays to a threshold (here, 1.3 $\times 10^{11}$ ions/bunch), collisions begin at IP2. There is loss of beam in the Yellow ring.



The beam loss is monitored. The spikes in the Blue ring are due to insertions of carbon ribbons for measuring the beam polarization. Beam-beam tune shift causes loss of ions in Yellow when collisions begin at IP2. This loss typically decays with time, as shown.



Luminosity at IP6 (STAR) and IP8 (PHENIX) is mostly constant when collisions are initiated at IP2 (A_NDY).

HCal events



- Cosmic ray trigger is essentially the same as jet trigger, except that the threshold on the summed calorimeter response is set at 5 pC (20 counts)
- This is a trigger that will work without beam. We have other cosmic-ray triggers that will work with beam, when commissioned, for continuous monitoring.
- The tracks test noise, patterns, etc.

Run-11 A_N(Jet)

- Sivers effect only (no Collins effect contribution)
- Need A_N(Jet) measurements before DY
- With ~10/pb & P=50%, A_NDY run11 could measure A_N(Jet)



arXiv:1103.1591 jet A_N measurements are required to clarify signs of quark/gluon correlators related to Sivers functions.



Non-zero jet analyzing power essentially a prerequisite before proceeding to Drell Yan