

Models of neutrino mass, mixing and CP violation

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys. G: Nucl. Part. Phys. 42 123001

(<http://iopscience.iop.org/0954-3899/42/12/123001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 159.93.14.8

This content was downloaded on 12/11/2015 at 13:06

Please note that [terms and conditions apply](#).

Topical Review

Models of neutrino mass, mixing and CP violation

Stephen F King

Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, UK

E-mail: king@soton.ac.uk

Received 4 March 2015

Accepted for publication 2 October 2015

Published 9 November 2015



CrossMark

Abstract

In this topical review we argue that neutrino mass and mixing data motivates extending the Standard Model (SM) to include a non-Abelian discrete flavour symmetry in order to accurately predict the large leptonic mixing angles and CP violation. We begin with an overview of the SM puzzles, followed by a description of some classic lepton mixing patterns. Lepton mixing may be regarded as a deviation from tri-bimaximal mixing, with charged lepton corrections leading to solar mixing sum rules, or tri-maximal lepton mixing leading to atmospheric mixing rules. We survey neutrino mass models, using a roadmap based on the open questions in neutrino physics. We then focus on the seesaw mechanism with right-handed neutrinos, where sequential dominance (SD) can account for large lepton mixing angles and CP violation, with precise predictions emerging from constrained SD (CSD). We define the flavour problem and discuss progress towards a theory of flavour using GUTs and discrete family symmetry. We classify models as direct, semidirect or indirect, according to the relation between the Klein symmetry of the mass matrices and the discrete family symmetry, in all cases focussing on spontaneous CP violation. Finally we give two examples of realistic and highly predictive indirect models with CSD, namely an A to Z of flavour with Pati–Salam and a fairly complete $A_4 \times SU(5)$ SUSY GUT of flavour, where both models have interesting implications for leptogenesis.

Keywords: Stephen F King, editorial board, neutrino, supersymmetry, beyond the Standard Model

(Some figures may appear in colour only in the online journal)

1. Introduction

The Nobel Prize in Physics for 2015 has just been awarded to Takaaki Kajita (Super-Kamiokande Collaboration, University of Tokyo, Japan) and to Arthur B McDonald (Sudbury Neutrino Observatory Collaboration, SNO, Canada) ‘for the discovery of neutrino oscillations, which shows that neutrinos have mass’ and ‘for their key contributions to the experiments which demonstrated that neutrinos change identities. This metamorphosis requires that neutrinos have mass. The discovery has changed our understanding of the innermost workings of matter and can prove crucial to our view of the Universe.’ In 1998 Takaaki Kajita presented to the world the discovery that neutrinos produced in the atmosphere switch between two identities on their way to Earth. Arthur McDonald subsequently led the Canadian collaboration which demonstrated that neutrinos from the Sun do not disappear on their way to Earth, but change identity by the time of arrival to the SNO detector. Since then there have been many developments in neutrino physics. This topical review will focus on the most recent developments since 2012.

The year 2012 was important in physics for two quite different reasons: the discovery of the Higgs boson and the measurement of the neutrino reactor angle. While the Higgs boson discovery supports the electroweak symmetry breaking sector of the standard model (SM), the reactor angle marks the completion of the \mathcal{CP} conserving part of the leptonic mixing matrix. While the Higgs discovery made a big splash in network TV headlines across the world, the reactor angle measurement only made a small plop in physics blogs and scientific journals, even though a year earlier neutrinos had been globally reported to *travel faster than light (sic)*. However truth is stranger than fiction, since although neutrinos are not superluminal, their mass and mixing requires physics beyond the SM, making them ‘ghostly beacons of new physics’ [1]. By contrast, the Higgs discovery serves only to confirm the SM, with its properties being exactly as predicted to increasing levels of accuracy.

While all physicists agree that the measured reactor angle opens up the prospect of measuring \mathcal{CP} violation in neutrino experiments in the foreseeable future, the theoretical significance of the reactor angle discovery splits the community. Consider the early history of neutrino model building, from 1998 onwards, summarized in the reviews [2–5]. Following the large atmospheric and solar mixing discoveries in 2002, many of the models in [3] involved sequential dominance (SD) [6, 7], which predicts a normal neutrino mass hierarchy $m_1 \ll m_2 \ll m_3$ and a large reactor angle $\theta_{13} \lesssim m_2/m_3$. Models incorporating SD were constructed with $SU(3)$ family symmetry providing an explanation of maximal atmospheric mixing via vacuum alignment [8]. There then followed the age of tri-bimaximal (TB) mixing with an explosion of models involving a zero reactor angle, enforced by discrete family symmetry [9], as reviewed in [10, 11]. A polar opposite approach called Anarchy was also put forward early on [12], the idea being that lepton mixing is determined randomly as if God plays dice with the neutrino mass matrix. According to Anarchy the reactor angle is on the same footing as the atmospheric and solar angles, and hence was generally expected to be large. Following the measurement of a large reactor angle in 2012, many people have jumped to the conclusion that Anarchy is preferred to discrete family symmetry. However, unlike that other dice-throwing theory (quantum mechanics), Anarchy is intrinsically untestable. Instead one is led to rather sterile arguments about whether Anarchy is statistically better than models with family symmetry [13]. By contrast discrete family symmetry models are highly testable, indeed many of them were excluded by the measurement of the reactor angle. Moreover, the model builders have been hard at work, showing how discrete family symmetry models could be modified to account for the observed reactor angle, as discussed in recent reviews [14, 15]. Before concluding that model builders are serial revisionists, it is worth remembering that SD

predicted a large reactor angle $\theta_{13} \lesssim m_2/m_3$, a decade before it was measured [7], although understanding why this bound is saturated requires further input. This topical review, then, summarizes the model building developments since 2012, including the latest progress in spontaneous \mathcal{CP} violation and versions of CSD which explain why $\theta_{13} \sim m_2/m_3$, as well as grand unified theories (GUTs) with discrete flavour symmetry which incorporate these ideas.

We have mentioned that the discovery of neutrino mass and mixing in 1998, unlike the Higgs boson discovery, requires new physics beyond the SM. To understand why, it is enough to realize that the origin of neutrino mass remains unknown. Although many types of new physics beyond the SM allow neutrino mass, since no BSM physics has been found, many people favour adding just right-handed (RH) neutrinos and nothing else. If we do this then immediately we face the questions of how many RH neutrinos and what are their Majorana masses, which may range from zero to the Planck scale? Alternatively, we may just add effective operators with some cut-off mass scales (proposed by Weinberg) but if we do this we soon learn that some of these operators must be associated with a mass scale below the Planck scale, and probably also below the scale of grand unification, so the SM must break down at some scale. It is clear then that the origin of neutrino mass requires the first (and so far only) new physics beyond the SM of particle physics. What is the nature of the new physics? Although the origin of tiny neutrino mass is unknown, it could imply some sort of seesaw mechanism at a high mass scale, or maybe new particles associated with loop models of neutrino mass, or perhaps R-parity violating supersymmetry (SUSY)—maybe even extra dimensions? Any example of such new physics would have implications for the unification of matter, forces and flavour GUTs, and the extra seven parameters (or nine parameters if neutrinos are Majorana) associated with neutrino mass and mixing makes the flavour problem of the SM less ignorable since any version of an extended SM must now have around 30 parameters in total, surely too many for a satisfactory theory of particle physics? The presence of such a large number of parameters associated with flavour, as well as the phenomenon of large lepton mixing, very unlike that witnessed in the quark sector, has stimulated attempts to address the flavour problem based on family symmetry. Since atmospheric and solar mixing continues to display the TB form, indicative of two and three-fold permutation symmetries, this continues to motivate the use of discrete non-Abelian family symmetry. An important point to realize is that discrete non-Abelian family symmetry does not imply zero reactor angle, as we discuss at length in this review.

In the realm of Cosmology, neutrinos could be responsible for our very existence, since leptogenesis is now the leading candidate for the origin of matter–antimatter asymmetry [16] (the SM gives too small a value for such asymmetry). Neutrino mass tends to wash out galaxy structures, since eV neutrinos represent a hot dark matter component. On the other hand warm dark matter, for example from keV sterile neutrinos, could be responsible for all the dark matter in the Universe [17]. More speculatively, neutrinos could play a role in inflating the Universe from Planck scale size to its present size via sneutrino inflation [18]. And it remains an intriguing possibility that neutrino mass is somehow related to dark energy since the scales happen to be the same order of magnitude (see e.g. [19] and references therein).

Turning to neutrino phenomenology, the observed pattern of neutrino masses and lepton mixing is quite remarkable. Neutrinos have tiny masses (for all three neutrinos, much less than the electron mass) which are not very hierarchical. Such masses break the separate lepton numbers L_e, L_μ, L_τ , but may or may not preserve the total lepton number $L = L_e + L_\mu + L_\tau$ (depending on whether neutrinos are Dirac or Majorana). Neutrinos certainly mix a lot (unlike the quarks). As mentioned already, neutrino mass implies at least seven new parameters as compared to the minimal SM: three neutrino masses, three lepton mixing angles, one \mathcal{CP} violating phase. If the three neutrino masses are Majorana in nature, there are two further \mathcal{CP}




Figure 1. Neutrino mixing angles (assuming zero \mathcal{CP} violation) may be represented as Euler angles relating the charged lepton mass basis states (ν_e, ν_μ, ν_τ) to the mass eigenstate basis states (ν_1, ν_2, ν_3).

violating phases. As discussed above, the origin of neutrino mass is unknown. For example, heavy RH neutrino exchange could be responsible for the Weinberg operators. Such heavy RH Majorana masses play a crucial role in generating the matter–antimatter asymmetry via leptogenesis [16], mentioned above. Alternatively, the origin of neutrino mass (and matter–antimatter asymmetry) could be something completely different.

The ground breaking neutrino oscillation milestones may be summarized as (for original experimental references see e.g. [20]):

- 1998 Atmospheric ν_μ disappear, implying large θ_{23} (SuperKamiokande).
- 2002 Solar ν_e disappear, implying large θ_{12} (SuperKamiokande, following the classic Homestake and Gallium experiments).
- 2002 Solar ν_e converted to ν_μ and ν_τ (Sudbury Neutrino Observatory).
- 2004 Reactor $\bar{\nu}_e$ seen to disappear and reappear (KamLAND).
- 2004 Accelerator ν_μ first seen to disappear (K2K).
- 2006 Accelerator ν_μ disappearance studied in detail (MINOS).
- 2010 Accelerator ν_μ converted to an observed ν_τ (OPERA).
- 2011 Accelerator ν_μ converted to ν_e giving a hint for θ_{13} (T2K, MINOS).
- 2012 Reactor $\bar{\nu}_e$ disappear, θ_{13} accurately measured (Daya Bay, RENO).

The fast pace of neutrino physics is well illustrated by the reactor angle which was unmeasured before 2012 but is now measured to incredible accuracy: $\theta_{13} \approx 8.4^\circ \pm 0.2^\circ$ (see [21] and references therein). The other angles are determined from global fits [22–24] to be: $\theta_{12} \approx 34^\circ \pm 1^\circ$ and $\theta_{23} \approx 45^\circ \pm 3^\circ$, and first hints of the \mathcal{CP} -violating (CPV) phase $\delta \sim -\pi/2$ have been reported, however with a large error $\pm\pi/3$. The meaning of the angles is given in figure 1. Two possible mass squared orderings are possible as explained in figure 2. The above quoted angles are extracted from the global fits which are displayed in figure 3 for the normal ordering (NO) case.

Despite the great pace of progress in neutrino physics, there are still several unanswered experimental questions, as follows:

- Is the atmospheric neutrino angle θ_{23} in the first or second octant?
- Do neutrino mass squared eigenvalues have a NO or inverted ordering (IO)?
- What is the value of the lightest neutrino mass?
- Are neutrinos Dirac or Majorana?
- Is \mathcal{CP} violated in the leptonic sector and if so by how much?




Figure 2. The probability that a particular neutrino mass state ν_i with mass m_i contains a particular charged lepton mass basis state (ν_e, ν_μ, ν_τ) is represented by colours. The left and right panels of the figure are referred to as normal or inverted mass squared ordering, respectively, referred to as NO or IO. The value of the lightest neutrino mass is presently unknown.




Figure 3. Global fits [22–24] to the lepton mixing angles for the case of normal neutrino mass squared ordering. The green dots are the best fit points, the red (blue) areas indicate the one (three) sigma ranges. The dashed lines indicate tri-bimaximal-Cabibbo (TBC) mixing [26], namely the values: $s_{12}^2 = 1/3, s_{23}^2 = 1/2 s_{13}^2 = \theta_C^2/2$. The Fogli *et al* fits do not include the latest Daya Bay results, whereas the other two do. This is a modified version of a figure provided privately by Stefano Morisi.

What is the \mathcal{CP} violating phase δ ? Is the current hint $\delta \sim -\pi/2$ going to hold up? It is common but incorrect to refer to the mass squared ordering question as the ‘neutrino mass hierarchy’. However the ‘ordering’ question is separate from whether neutrinos are hierarchical in nature or approximately degenerate, which is to do with the lightest neutrino mass. Many neutrino experiments are underway or in the planning stages to address these questions such as T2K, NO ν A, Daya Bay, JUNO, RENO, KATRIN, LBNE/DUNE and many neutrinoless double beta decay experiments running and planned [25].

The layout of the remainder of this topical review is as follows. In section 2 we give an overview of the SM puzzles, followed in section 3 by a description of some classic lepton mixing patterns. Lepton mixing may be regarded as a deviation of TB mixing, as discussed in section 4, with charged lepton corrections leading to solar mixing sum rules, while tri-maximal lepton mixing leads to atmospheric mixing rules. Motivated by the open questions in neutrino physics, in section 5 we discuss a roadmap of the origin of neutrino mass, focussing

on the seesaw mechanism with RH neutrinos, leading to SD, whose constrained form can lead to predictions for mixing angles. In section 6 we survey the flavour problem and its possible resolution based on symmetry, using both GUTs and family symmetry, then classify flavour models as direct, semidirect or indirect, according to the relation between the Klein symmetry of the mass matrices and the underlying discrete family symmetry, focussing on the possibility of spontaneous \mathcal{CP} violation. In section 7 we give two examples of realistic and highly predictive indirect models, namely an A to Z of flavour with Pati–Salam (PS) and a fairly complete $A_4 \times SU(5)$ SUSY GUT of flavour, where both models incorporate constrained sequential dominance (CSD) and have interesting implications for leptogenesis. In section 8 we speculate about the possible F-theory origin of SUSY GUTs with discrete family symmetry. Section 9 concludes this topical review.

2. The SM puzzles

Even though the SM is essentially complete, following the Higgs boson discovery, we are far from satisfied since it offers no solutions to the cosmological puzzles of matter–antimatter asymmetry, dark matter and dark energy. It therefore cannot be the final answer. In addition it leaves three unresolved puzzles in its wake:

The origin of mass—the origin of the Higgs vacuum expectation value (VEV), its stability under radiative corrections, and the solution to the hierarchy problem (most urgent problem of LHC).

The quest for unification—the question of whether the three known forces of the SM may be related into a GUT, and whether such a theory could also include a unification with gravity.

The problem of flavour—the problem of the undetermined fermion masses and mixing angles (including neutrino masses and lepton mixing angles) together with the \mathcal{CP} violating phases, in conjunction with the observed smallness of flavour changing neutral currents and very small strong \mathcal{CP} violation. In particular the unknown origin of the extremely small neutrino masses for all three families may offer a clue as to what lies beyond the SM.

The differences between quark and lepton mixing may also offer clues concerning the flavour problem. Certainly the flavour problem has now become much richer, following the discovery of neutrino mass and mixing, so we shall discuss more about this in section 6. We now digress slightly to discuss an alternative point of view that frequently is voiced. Namely, all that is required for neutrino masses is to add two or three RH neutrinos with zero Majorana mass due to a conserved $B - L$, and Yukawa couplings for all neutrino families of about 10^{-11} and that no new physics beyond this is required. However this conservative point of view involves a new mystery, namely why the third family Yukawa couplings are of order unity for the top quark, and not very small for the bottom quark and τ lepton, but are of order 10^{-11} for the third family of neutrinos. The seesaw mechanism [27], i.e. large third family neutrino Yukawa couplings, with physical neutrino masses suppressed by heavy RH Majorana masses with $B - L$ broken at a high scale, provides an elegant solution to this mystery, and opens the door to leptogenesis. However, the seesaw mechanism by itself does not account for the observed large lepton mixing, so we need to go further.

It has been one of the long standing goals of theories of particle physics beyond the SM to predict quark and lepton masses and mixings. With the discovery of neutrino mass and mixing, this quest has received a massive impetus. Indeed, perhaps the greatest advance in particle physics over the past decade has been the discovery of neutrino mass and mixing involving large mixing. The largeness of the lepton mixing angles contrasts with the

smallness of the quark mixing angles, and this observation, together with the smallness of neutrino masses, provides new and tantalizing clues in the search for the origin of quark and lepton flavour. For example, it is intriguing that the smallest lepton mixing may be related to the largest quark mixing, $|U_{e3}| \approx \theta_C/\sqrt{2}$ where θ_C is the Cabibbo angle, although this relation is in tension with the latest Daya Bay results. The quest to understand the origin of the three families of quarks and leptons and their pattern of masses and mixing parameters is called the flavour puzzle, and motivates the introduction of family symmetry. In particular, as we shall see, lepton mixing provides a motivation for discrete family symmetry, which will form the central part of this review. As we shall also see, such theories demand a high precision knowledge of the lepton mixing angles, beyond that currently achieved.

The PDG [28] advocates CKM and the PMNS mixing matrices being parameterized by:

$$\begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{pmatrix}, \quad (1)$$

where $\delta \equiv \delta_{\mathcal{CP}}$ is the \mathcal{CP} violating phase in each sector (quark and lepton) and $s_{13} = \sin \theta_{13}$, etc with (very) different angles for quarks and leptons. In the quark sector the mixing angles are all small, with

$$s_{12} = \lambda, \quad s_{23} \sim \lambda^2, \quad s_{13} \sim \lambda^3, \quad (2)$$

where $\lambda = 0.226 \pm 0.001$ is the Wolfenstein parameter [28]. The \mathcal{CP} violating phase in the quark sector is roughly $\delta \sim (\pi/2)/\sqrt{2}$. The large lepton mixing, discussed in further in the next section, must arise in conjunction with the mechanism responsible for the smallness of neutrino mass, which however is unknown. In the case of Majorana neutrinos, the PMNS matrix also involves the phase matrix [28]: $\text{diag}(1, e^{i\frac{\alpha_{21}}{2}}, e^{i\frac{\alpha_{31}}{2}})$ which post-multiplies the above matrix. It is a puzzle why the quark mixing angles are so small while the lepton mixing angles are so large.

3. Patterns of lepton mixing

The origin of neutrino mass is unknown, as discussed above, meaning that there is not a unique electroweak description, as for the quarks. There are basically two possibilities, either neutrinos are Dirac or Majorana. Here we shall exclusively focus on the Majorana case. Majorana neutrino masses must be generated in such a way that, below the electroweak breaking scale, we should obtain the leptonic Lagrangian

$$\mathcal{L}^{\text{lepton}} = -\nu_d Y_{ij}^e \bar{\nu}_L^i e_R^j - \frac{1}{2} m_{ij}^{\nu_e} \bar{\nu}_{eL}^i \nu_{eL}^j + \text{h.c.} \quad (3)$$

The resulting matrices are diagonalized by unitary transformations

$$U_{eL} Y^e U_{eR}^\dagger = \begin{pmatrix} y_e & 0 & 0 \\ 0 & y_\mu & 0 \\ 0 & 0 & y_\tau \end{pmatrix}, \quad U_{\nu_{eL}} m^{\nu_e} U_{\nu_{eL}}^T = \begin{pmatrix} m_1 & 0 & 0 \\ 0 & m_2 & 0 \\ 0 & 0 & m_3 \end{pmatrix}. \quad (4)$$

¹ Interestingly, in the original KM parametrization, the \mathcal{CP} violating phase is roughly maximal $\delta \sim \pi/2$, as is the angle $\alpha \sim \pi/2$ in the standard unitarity triangle representing the orthogonality of the first and third columns of the CKM matrix [28].

The couplings to W^- are given by $-\frac{g}{\sqrt{2}}\bar{e}_L^i\gamma^\mu W_\mu^- \nu_{eL}^i$, hence the charged currents in terms of mass states are

$$\mathcal{L}_{\text{lepton}}^{\text{CC}} = -\frac{g}{\sqrt{2}}(\bar{e}_L \quad \bar{\mu}_L \quad \bar{\tau}_L)U_{\text{PMNS}}\gamma^\mu W_\mu^- \begin{pmatrix} \nu_{1L} \\ \nu_{2L} \\ \nu_{3L} \end{pmatrix} + \text{h.c.}, \quad (5)$$

where we identify the PMNS matrix as

$$U_{\text{PMNS}} = U_{eL} U_{\nu eL}^\dagger. \quad (6)$$

Now only three of the six phases can be removed since each of the three charged lepton mass terms such as $m_e \bar{e}_L e_R$, etc, is left unchanged by a phase rotation $e_L \rightarrow e^{i\phi_e} e_L$ and $e_R \rightarrow e^{i\phi_e} e_R$, etc, where the three phases ϕ_e , etc, are chosen to leave three physical (irremovable) phases in U_{PMNS} . There is no such phase freedom in the Majorana mass terms $-\frac{1}{2}m_i \bar{\nu}_{iL}^c \nu_{iL}^c$ where m_i are real and positive.

We already discussed the PDG parametrization of the PMNS matrix U_{PMNS} in equation (1). We now discuss three simple ansätze for U_{PMNS} which have been proposed. Although each of them involves a zero reactor angle and is hence excluded, they will serve to motivate approaches which involve a non-zero reactor angle.

3.1. Bimaximal (BM) mixing

An early suggested pattern of lepton mixing is known as BM mixing with $s_{13}^2 = 0$ and $s_{12}^2 = s_{23}^2 = 1/2$ which could originate from the discrete group S_4 (see later). It has a maximal solar mixing angle [29], and is given by a matrix of the form

$$U_{\text{BM}} = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (7)$$

3.2. TB mixing

A second pattern of lepton mixing which came to dominate the model building community until the measurement of the reactor angle is the TB mixing matrix [30]. This has been associated with models based on the flavour symmetries A_4 and S_4 (see later). Like BM mixing it predicts $s_{13}^2 = 0$ and $s_{23}^2 = 1/2$ but differs in that it predicts a solar mixing angle given by $s_{12} = 1/\sqrt{3}$, i.e. $\theta_{12} \approx 35.3^\circ$. The mixing matrix is given explicitly by

$$U_{\text{TB}} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (8)$$

3.3. Golden ratio (GR) mixing

Another pattern of lepton mixing which was viable until the reactor angle measurement associates the GR $\varphi = \frac{1+\sqrt{5}}{2}$ with the solar mixing angle. The original GR mixing pattern is related to the flavour symmetry A_5 [31]. As above, it predicts $s_{13}^2 = 0$ and $s_{23}^2 = 1/2$ but

differs by having a solar mixing angle given by $t_{12}' = 1/\varphi$, i.e. $\theta_{12}' \approx 31.7^\circ$, resulting in the mixing matrix

$$U_{\text{GR}} = \begin{pmatrix} \frac{\varphi}{\sqrt{2+\varphi}} & \frac{1}{\sqrt{2+\varphi}} & 0 \\ -\frac{1}{\sqrt{4+2\varphi}} & \frac{\varphi}{\sqrt{4+2\varphi}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{4+2\varphi}} & -\frac{\varphi}{\sqrt{4+2\varphi}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (9)$$

4. Deviations from TB mixing

4.1. Deviation parameters

After the measurement of the reactor angle, TB mixing is excluded. However, TB mixing still remains a reasonable approximation to lepton mixing for the solar and atmospheric angles. It therefore makes sense to expand the angles about their TB values [32, 33]:

$$\sin \theta_{12} = \frac{1}{\sqrt{3}}(1 + s), \quad (10)$$

$$\sin \theta_{23} = \frac{1}{\sqrt{2}}(1 + a), \quad (11)$$

$$\sin \theta_{13} = \frac{r}{\sqrt{2}}, \quad (12)$$

where s , a , and r are the (s)olar, (a)tmospheric and (r)eactor deviation parameters such that TB mixing [30] is recovered for $s = a = r = 0$. For example, TBC mixing [26] corresponds to $s = a = 0$ and $r = \theta_C$, where θ_C is the Cabibbo angle, which is consistent with data at three sigma as shown in figure 3.

4.2. Tri-maximal mixing and atmospheric sum rules

Tri-maximal mixing is a variation which preserves either the first or the second column of the TB mixing matrix in equation (8), leading to two versions called TM1 or TM2

$$U_{\text{TM1}} = \begin{pmatrix} \frac{2}{\sqrt{3}} & - & - \\ -\frac{1}{\sqrt{6}} & - & - \\ \frac{1}{\sqrt{6}} & - & - \end{pmatrix}, \quad U_{\text{TM2}} = \begin{pmatrix} - & \frac{1}{\sqrt{3}} & - \\ - & \frac{1}{\sqrt{3}} & - \\ - & -\frac{1}{\sqrt{3}} & - \end{pmatrix}. \quad (13)$$

The dashes indicate that the other elements are undetermined. However these are fixed once the reactor angle is specified (it is a free parameter here). These imply the relations

$$\text{TM1:} \quad |U_{e1}| = \sqrt{\frac{2}{3}} \quad \text{and} \quad |U_{\mu 1}| = |U_{\tau 1}| = \frac{1}{\sqrt{6}}; \quad (14)$$

$$\text{TM2:} \quad |U_{e2}| = |U_{\mu 2}| = |U_{\tau 2}| = \frac{1}{\sqrt{3}}. \quad (15)$$

The *atmospheric mixing sum rule*

$$a = \lambda r \cos \delta + \mathcal{O}(a^2, r^2), \quad \text{with } s = \mathcal{O}(a^2, r^2), \quad (16)$$

was first derived in [32] by expanding the PMNS matrix to first order in r, s, a . It also follows from a first order expansion of equations (14) and (15), where $\lambda = 1$ for TM1 and $\lambda = -1/2$ for TM2. The study of correlations of this type, and their application to the discrimination between underlying models, has been shown to be a realistic aim for a next-generation superbeam experiment [34], see for example figure 4.

4.3. Charged lepton corrections and solar sum rules

Now suppose that neutrino mixing U_{TB}^ν obeys TB exactly so that the PMNS matrix according to equation (6) is given by $U_{\text{PMNS}} = U^e U_{\text{TB}}^\nu$ where U_{TB}^ν is equated to equation (8) while U^e encodes some unknown charged lepton corrections which must be small since U_{PMNS} is not far from TB mixing. The *solar mixing sum rule* [35–37] then follows from the assumption that $\theta_{23}^e = \theta_{13}^e = 0$. If the charged lepton mixing matrix involves a Cabibbo-like mixing, then the PMNS matrix is given by

$$U_{\text{PMNS}} = \begin{pmatrix} c_{12}^e & s_{12}^e e^{-i\delta_{12}^e} & 0 \\ -s_{12}^e e^{i\delta_{12}^e} & c_{12}^e & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{\sqrt{2}}{3} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \dots & \dots & \frac{s_{12}^e}{\sqrt{2}} e^{-i\delta_{12}^e} \\ \dots & \dots & \frac{c_{12}^e}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix} \quad (17)$$

Comparing to the PMNS parametrization in equation (1) we identify

$$|U_{e3}| = s_{13} = \frac{s_{12}^e}{\sqrt{2}}, \quad (18)$$

$$|U_{\tau 1}| = |s_{23}s_{12} - s_{13}c_{23}c_{12}e^{i\delta}| = \frac{1}{\sqrt{6}}, \quad (19)$$

$$|U_{\tau 2}| = |-c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta}| = \frac{1}{\sqrt{3}}, \quad (20)$$

$$|U_{\tau 3}| = c_{13}c_{23} = \frac{1}{\sqrt{2}}. \quad (21)$$

The first equation predicts a reactor angle $\theta_{13} \approx 9.2^\circ$ if $\theta_e \approx \theta_C \approx 13^\circ$. The second and fourth equations allow to eliminate θ_{23} to give a new relation between the PMNS parameters, θ_{12} , θ_{13} and δ called a *solar sum rule*, which may be expanded to first order to give the approximate relations

$$\theta_{12} \approx 35.26^\circ + \theta_{13} \cos \delta \quad \text{or} \quad \cos \delta \approx \frac{\theta_{12} - 35.26^\circ}{\theta_{13}} \quad (22)$$

where $35.26^\circ = \sin^{-1} \frac{1}{\sqrt{3}}$, which can be recast as [32],

$$s = r \cos \delta + \mathcal{O}(a^2, r^2). \quad (23)$$

Recently it has been realized that, keeping $\theta_{13}^e = 0$, but allowing $\theta_{23}^e \neq 0$, the following exact result can be obtained by generalizing equation (17) to allow both $\theta_{12}^e, \theta_{23}^e \neq 0$ [38]:




Figure 4. Expectation for the determination of the TM1 atmospheric mixing sum rule $a \approx r \cos \delta$ at one, three and five sigma for a low energy neutrino factory with a magnetized iron detector (for more details see [34]).




Figure 5. Solar sum rules prediction for $\cos \delta$ using the exact result in equation (25) for TB neutrino mixing (for more details see [38]).

$$\frac{|U_{\tau 1}|}{|U_{\tau 2}|} = \frac{|s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta}|}{|-c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta}|} = \frac{1}{\sqrt{2}}, \quad (24)$$

for the previous case $\theta_{23}^e = \theta_{13}^e = 0$ this result follows trivially by taking the ratio of the two equations (19) and (20). Therefore this result applies to that case also. However it turns out that θ_{23}^e cancels in this ratio which is therefore a more general sum rule (though not completely general since it still assumes $\theta_{13}^e = 0$). After some algebra, equation (24) leads to an exact prediction for $\cos \delta$ in terms of the other physical lepton angles

$$\cos \delta = \frac{t_{23}s_{12}^2 + s_{13}^2c_{12}^2/t_{23} - \frac{1}{3}(t_{23} + s_{13}^2/t_{23})}{\sin 2\theta_{12}s_{13}}, \quad (25)$$

as displayed in figure 5.

When expanded to first order, equation (25) reduces to the leading order sum rule in equation (22). This is not too surprising since the previous sum rule case also satisfies equation (24). The leading order sum rule in equation (22) offers a simple way to understand

the results in figure 5. For example from figure 5 it seems that TB neutrino mixing predicts $\cos \delta \approx 0$ if $\theta_{12} \approx 35.26^\circ$, which is obvious from² equation (22). This can also be understood from equation (25) where we see that for $s_{12}^2 = 1/3$ the leading terms $t_{23}s_{12}^2$ and $\frac{1}{3}t_{23}$ in the numerator cancel, leaving $\cos \delta = s_{13}/(2\sqrt{2}t_{23}) \approx 0.05$ which is consistent with the numerical estimates of the error given above for a range of θ_{12} .

Solar sum rules can also be obtained for different types of neutrino mixing such as U_{BM}^ν (which is almost excluded) and U_{GR}^ν (which gives similar results to the case U_{TB}^ν considered here). The general formula given in [38] is

$$\cos \delta = \frac{t_{23}s_{12}^2 + s_{13}^2c_{12}^2/t_{23} - s_{12}^{\nu 2}(t_{23} + s_{13}^2/t_{23})}{\sin 2\theta_{12}s_{13}}, \quad (26)$$

where $s_{12}^{\nu 2} = \frac{1}{3}, \frac{1}{2}$ for $U_{\text{TB,BM}}^\nu$ and so on. The prospects for studying *solar sum rules* at JUNO and LBNF is discussed in [38]. A slightly more lengthy but equivalent formula to equation (25) had been previously derived [39] by an alternative method involving an auxiliary phase ϕ without using the elegant result equation (24),

$$\cos \delta = \frac{t_{23}}{\sin 2\theta_{12}s_{13}} \left[\cos 2\theta_{12}^\nu + (s_{12}^2 - c_{12}^{\nu 2})(1 - \cot^2 \theta_{23}s_{13}^2) \right]. \quad (27)$$

We prefer the simpler form in equation (26) which involves θ_{12}^ν in only one place since it exhibits the cancellation between the terms $t_{23}s_{12}^2$ and $s_{12}^{\nu 2}t_{23}$ when $s_{12}^2 = s_{12}^{\nu 2}$ responsible for the prediction $\cos \delta \approx 0$ in this case. We also advocate the much simpler derivation of equation (26) given in [38].

Finally we give a word of caution that when comparing leading order sum rules to the exact results the ratio $(\cos \delta)_{\text{exact}}/(\cos \delta)_{\text{LO}}$ used in [40] will lead to misleading results when $(\cos \delta)_{\text{LO}} \approx 0$. In general it is safer to compare them using the experimentally relevant quantity $\Delta(\cos \delta) = (\cos \delta)_{\text{exact}} - (\cos \delta)_{\text{LO}}$ defined in [38]. For example we find $\Delta(\cos \delta) \lesssim 0.1$ for TB neutrino mixing corrected by charged lepton mixing. It will take experiment a long time to reach this level of precision, so for present purposes the linear approximation in equation (22) is adequate for TB neutrino mixing.

5. Neutrino mass models

5.1. The open questions from neutrino physics

Despite the great progress coming from neutrino oscillation experiments there are still some outstanding questions. Are the lepton mixing angles consistent with TBC mixing? If not then is the atmospheric angle in the first or second octant? What is the leptonic \mathcal{CP} violating phase δ ? Is the current hint $\delta \sim -\pi/2$ going to hold up? Maybe there is no \mathcal{CP} violation in the lepton sector? Are neutrino mass squared ordered normally or inverted³? What is the lightest neutrino mass? Are neutrino masses Majorana or Dirac in nature? Many neutrino experiments are underway or in the planning stages to address these questions such as T2K, NO ν A, Daya Bay, JUNO, RENO, KATRIN, DUNE and many neutrinoless double beta decay experiments running and planned [25].

² Note that $\cos \delta \sim 0$ is consistent with the current hint $\delta \sim -\pi/2$.

³ It is common but incorrect to refer to this question as the ‘neutrino mass hierarchy’ since the ‘ordering’ question is separate from whether neutrinos are hierarchical in nature or approximately degenerate, which is to do with the lightest neutrino mass.




Figure 6. Roadmap of neutrino mass models.

5.2. Road map of neutrino mass models

Everyone can invent her or his personal roadmap of neutrino mass models, one example being that shown in figure 6. The blue boxes contain experimental questions and the red boxes possible theoretical consequences. In this subsection we shall briefly describe the possible theoretical options for producing neutrino mass.

It is worthwhile to first recall why the observation of non-zero neutrino mass and mixing is evidence for new physics beyond the SM. The most intuitive way to understand why neutrino mass is forbidden in the SM, is to understand that the SM predicts that neutrinos always have a ‘left-handed’ spin—rather like rifle bullets which spin counter clockwise to the direction of travel. In fact this property was first experimentally measured in 1958, two years after the neutrino was discovered, by Maurice Goldhaber, Lee Grodzins and Andrew Sunyar. More accurately, the ‘handedness’ of a particle describes the direction of its spin vector along the direction of motion, and the neutrino being ‘left-handed’ means that its spin vector always points in the opposite direction to its momentum vector. The fact that the neutrino is left-handed, written as ν_L , implies that it must be massless. If the neutrino has mass then, according to special relativity, it can never travel at the speed of light. In principle, a fast moving observer could therefore overtake the spinning massive neutrino and would see it moving in the opposite direction. To the observer, the massive neutrino would therefore appear RH. Since the SM predicts that neutrinos must be strictly left-handed, it follows that neutrinos are massless in the SM. It also follows that the discovery of neutrino mass implies new physics beyond the SM, with profound implications for particle physics and cosmology.

Neutrinos are massless in the SM for three independent reasons:

- There are no RH neutrinos ν_R .
- There are only Higgs doublets (and no Higgs triplets) of $SU(2)_L$.
- There are only renormalizable terms.

In the SM, the three massless neutrinos ν_e, ν_μ, ν_τ are distinguished by separate lepton numbers L_e, L_μ, L_τ . Neutrinos and antineutrinos are distinguished by total conserved lepton number $L = L_e + L_\mu + L_\tau$. To generate neutrino mass we must relax one or more of the above three conditions. For example, by adding RH neutrinos the Higgs mechanism of the SM can give neutrinos the same type of mass as the Dirac electron mass or other charged lepton and quark masses, which would generally break the separate lepton numbers L_e, L_μ, L_τ , but preserve the total lepton number L . However it is also possible for neutrinos to have a new type of mass of a type first proposed by Majorana, which would also break L . There exists a special case where total lepton number L is broken, but the combination $L_e - L_\mu - L_\tau$ is

conserved; such a symmetry would give rise to a neutrino mass matrix with an inverted mass spectrum.

From the theoretical perspective, the main unanswered question is the origin of neutrino mass, and in particular the smallness of neutrino mass. The simplest possibility is that neutrinos have Dirac mass just like the electron mass in the SM, namely due to a term like $y_D \bar{L} H \nu_R$, where L is a lepton doublet containing ν_L , H is a Higgs doublet and ν_R is a RH neutrino. The observed smallness of neutrino masses implies that the Dirac Yukawa coupling y_D must be of order 10^{-12} to achieve a Dirac neutrino mass of about 0.1 eV. Advocates of Dirac masses point out that the electron mass already requires a Yukawa coupling y_e of about 10^{-6} , so we are used to such small Yukawa couplings. In this case, all that is required is to add RH neutrinos ν_R to the SM and we are done. Well, almost. It still needs to be explained why the ν_R have zero Majorana mass, after all they are gauge singlets and so nothing prevents them acquiring (large) Majorana mass terms $M_{RR} \nu_R \nu_R$ where M_{RR} could be as large as the Planck scale. Moreover, Majorana masses offer a unique (and testable) way to generate neutrino masses (since neutrinos do not carry electric charge) even without RH neutrinos. The simplest way to generate Majorana mass is via $y_M \Delta LL$ where Δ is a Higgs triplet and y_M is a Yukawa coupling associated with Majorana mass. Alternatively, at the effective level, Majorana neutrino mass can result from some additional dimension five operators which couple two lepton doublets L to two Higgs doublets H first proposed by Weinberg [41],

$$-\frac{1}{2} H L^T \kappa H L, \quad (28)$$

where κ has dimension $(\text{mass})^{-1}$. This is a non-renormalizable operator, so it violates one of the tenets of the SM. In order to account for a neutrino mass of order 0.1 eV requires $\kappa \sim 10^{-14} \text{ GeV}^{-1}$. This suggests a new high energy mass scale M in physics, a small dimensionless coupling associated with κ , or both.

There are basically five different proposals for the origin of neutrino mass:

- The seesaw mechanisms [27, 42, 43], including low scale seesaw mechanisms [44] (Weinberg operator typically from large Majorana mass $M = M_R$ for RH neutrinos ν_R).
- R-parity violating SUSY [45] (Weinberg operator from TeV scale Majorana mass for neutralinos χ).
- TeV scale loop mechanisms [46–48] (Majorana mass from extra Higgs doublets and singlets at the TeV scale).
- Extra dimensions [49] (Dirac mass with small y_D due to RH neutrinos ν_R in the bulk).
- String theory [50, 51] (new mechanisms for generating large Majorana mass for RH neutrinos ν_R from Planck or string scale physics).

These different mechanisms are reviewed in [52]. Returning to the roadmap in figure 6, we see these mechanisms for neutrino mass represented by the red boxes, being related to the experimental question in the blue boxes.

Since no new physics has yet emerged from the LHC, we are led to consider the seesaw mechanism with RH neutrinos. However even in this case both the number of species and the mass spectrum of RH (or sterile) neutrinos is completely unknown [53]. As shown in figure 7 the mass spectrum can cover the whole range with different physical consequences as indicated. It is one of the goals of neutrino physics to determine this spectrum. In this topical review we shall focus on the classic seesaw mechanism [27] with very heavy RH neutrinos, with masses above the TeV scale, which may be incorporated into a theory of flavour, possibly related to string theory.




Figure 7. Possible mass spectrum of right-handed (or sterile) neutrinos corresponding to the physical implications as shown.

5.3. Seesaw mechanism with two RH neutrinos and SD

In this subsection we consider the high scale (classic) *seesaw* neutrino model involving just *two* RH neutrinos ν_R^{sol} and ν_R^{atm} with Yukawa couplings [6],⁴

$$(H_u/v_u)(a\bar{L}_e + b\bar{L}_\mu + c\bar{L}_\tau)\nu_R^{\text{sol}} + (H_u/v_u)(d\bar{L}_e + e\bar{L}_\mu + f\bar{L}_\tau)\nu_R^{\text{atm}} + \text{h.c.}, \quad (29)$$

where H_u is a Higgs doublet and v_u its VEV. The heavy RH Majorana masses are

$$M_{\text{sol}} \overline{\nu_R^{\text{sol}}} (\nu_R^{\text{sol}})^c + M_{\text{atm}} \overline{\nu_R^{\text{atm}}} (\nu_R^{\text{atm}})^c + \text{h.c.}. \quad (30)$$

In the basis, with rows $(\bar{\nu}_{eL}, \bar{\nu}_{\mu L}, \bar{\nu}_{\tau L})$ and columns $\nu_R^{\text{atm}}, \nu_R^{\text{sol}}$, the resulting Dirac mass matrix is

$$m^D = \begin{pmatrix} d & a \\ e & b \\ f & c \end{pmatrix} \equiv (m_{\text{atm}}^D \quad m_{\text{sol}}^D), \quad m_{\text{atm}}^D \equiv (d \ e \ f)^T, \quad m_{\text{sol}}^D \equiv (a \ b \ c)^T. \quad (31)$$

The (diagonal) RH neutrino heavy Majorana mass matrix M_R with rows $(\overline{\nu_R^{\text{atm}}}, \overline{\nu_R^{\text{sol}}})^T$ and columns $(\nu_R^{\text{atm}}, \nu_R^{\text{sol}})$ is

$$M_R = \begin{pmatrix} M_{\text{atm}} & 0 \\ 0 & M_{\text{sol}} \end{pmatrix}. \quad (32)$$

⁴ We follow the notation of the third paper in [6], which was the first paper to discuss a phenomenologically viable model with two RH neutrinos (see also [7]). Subsequently two RH neutrino models with two texture zeros were proposed in [54], however such two texture zero models are now phenomenologically excluded [55] for the case of a normal neutrino mass hierarchy considered here, while the one texture zero case $d = 0$ (see the second paper in [7]) remains viable.

The seesaw formula is [27],

$$m^\nu = -m^D M_R^{-1} (m^D)^T, \quad (33)$$

where m^ν is the the light effective left-handed Majorana neutrino mass matrix (i.e. the physical neutrino mass matrix), m^D is the Dirac mass matrix in LR convention and M_R is the (heavy) Majorana mass matrix. Using the seesaw formula dropping the overall minus sign which is physically irrelevant, the light effective left-handed Majorana neutrino mass matrix m^ν (i.e. the physical neutrino mass matrix) is, by multiplying the matrices

$$m^\nu = m^D M_R^{-1} (m^D)^T = \begin{pmatrix} \frac{a^2}{M_{\text{sol}}} + \frac{d^2}{M_{\text{atm}}} & \frac{ab}{M_{\text{sol}}} + \frac{de}{M_{\text{atm}}} & \frac{ac}{M_{\text{sol}}} + \frac{df}{M_{\text{atm}}} \\ \frac{ab}{M_{\text{sol}}} + \frac{de}{M_{\text{atm}}} & \frac{b^2}{M_{\text{sol}}} + \frac{e^2}{M_{\text{atm}}} & \frac{bc}{M_{\text{sol}}} + \frac{ef}{M_{\text{atm}}} \\ \frac{ac}{M_{\text{sol}}} + \frac{df}{M_{\text{atm}}} & \frac{bc}{M_{\text{sol}}} + \frac{ef}{M_{\text{atm}}} & \frac{c^2}{M_{\text{sol}}} + \frac{f^2}{M_{\text{atm}}} \end{pmatrix}. \quad (34)$$

The SD [6] assumptions are that $d \ll e, f$ and

$$\frac{(e, f)^2}{M_{\text{atm}}} \gg \frac{(a, b, c)^2}{M_{\text{sol}}}. \quad (35)$$

By explicit calculation, one can check that $\det m^\nu = 0$. Since the determinant of a Hermitian matrix is the product of mass eigenvalues

$$\det(m^\nu m^{\nu\dagger}) = m_1^2 m_2^2 m_3^2,$$

one may deduce that one of the mass eigenvalues of the complex symmetric matrix above is zero, which under the SD assumption is the lightest one $m_1 = 0$ with $m_3 \gg m_2$ since the model approximates to a single RH neutrino model [6]. Hence we see that *SD implies a normal neutrino mass hierarchy*. Including the solar RH neutrino as a perturbation, it can be shown that, for $d = 0$, together with the assumption of a dominant atmospheric RH neutrino in equation (35), leads to the approximate results for the solar and atmospheric angles [6],

$$\tan \theta_{23} \sim \frac{e}{f}, \quad \tan \theta_{12} \sim \frac{\sqrt{2}a}{b-c}. \quad (36)$$

Under the above SD assumption, each of the RH neutrinos contributes uniquely to a particular physical neutrino mass. The SD framework above with $d = 0$ leads to the relations in equation (36) together with the reactor angle bound [7],

$$\theta_{13} \lesssim m_2/m_3. \quad (37)$$

This result shows that SD allows for large values of the reactor angle, consistent with the measured value. Indeed the measured reactor angle, observed a decade after this theoretical bound was derived, approximately saturates the upper limit. In order to understand why this is so, we must go beyond the SD assumptions stated so far, and enter the realms of CSD.

5.4. CSD: the minimal predictive seesaw model

Let us return to equation (31) and set $d = 0$ and $e = f$, with $b = a$ and $c = -a$ [35]. The motivation is that from equation (36) one then approximately expects the good phenomenological relations $t_{23} \sim 1$ and $t_{12} \sim 1/\sqrt{2}$, although the value of the reactor angle bounded by equation (37) remains to be seen. With the above assumption, equation (34) becomes

$$m^\nu = \begin{pmatrix} \frac{a^2}{M_{\text{sol}}} & \frac{a^2}{M_{\text{sol}}} & \frac{-a^2}{M_{\text{sol}}} \\ \frac{a^2}{M_{\text{sol}}} & \frac{a^2}{M_{\text{sol}}} + \frac{e^2}{M_{\text{atm}}} & \frac{-a^2}{M_{\text{sol}}} + \frac{e^2}{M_{\text{atm}}} \\ \frac{-a^2}{M_{\text{sol}}} & \frac{-a^2}{M_{\text{sol}}} + \frac{e^2}{M_{\text{atm}}} & \frac{a^2}{M_{\text{sol}}} + \frac{e^2}{M_{\text{atm}}} \end{pmatrix}. \quad (38)$$

By explicit calculation one then finds that the neutrino mass matrix is exactly diagonalized by the TB mixing matrix in equation (8),

$$U_{\text{TB}}^T m^\nu U_{\text{TB}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{3a^2}{M_{\text{sol}}} & 0 \\ 0 & 0 & \frac{2e^2}{M_{\text{atm}}} \end{pmatrix}. \quad (39)$$

If the charged lepton mass matrix is diagonal, the interpretation is that these constrained couplings $d = 0$, $e = f$ with $b = a$ and $c = -a$ lead to TB mixing, with the lightest neutrino mass $m_1 = 0$, the second lightest neutrino identified as the solar neutrino with mass $m_2 = \frac{3a^2}{M_{\text{sol}}}$ and the heaviest neutrino identified as the atmospheric neutrino with mass $m_3 = \frac{2a^2}{M_{\text{atm}}}$. While TB mixing accurately gives the good relations $t_{23} = 1$ and $t_{12} = 1/\sqrt{2}$, unfortunately it also gives $\theta_{13} = 0$. This is known as CSD [35].

We can generalize the original idea of CSD to other examples of Dirac mass matrix with (in the notation of equation (31)) $d = 0$ and $e = f$ as before, but now with $b = na$ and $c = (n - 2)a$, for any positive integer n , which we refer to as CSD(n). The original CSD in equation (38) with $b = a$ and $c = -a$ is identified as the special case CSD($n = 1$). The motivation for CSD(n) is that for any n equation (36) implies $t_{23} \sim 1$ and $t_{12} \sim 1/\sqrt{2}$, although these results are strongly dependent on the relative phase between the first and second column of the Dirac mass matrix. CSD(n) then corresponds to the following pattern of couplings in the Dirac mass matrix in equation (31):

- CSD(1): $(m_{\text{atm}}^{\text{D}})^T = (0, e, e)$, $(m_{\text{sol}}^{\text{D}})^T = (a, a, -a)$ [35].
- CSD(2): $(m_{\text{atm}}^{\text{D}})^T = (0, e, e)$, $(m_{\text{sol}}^{\text{D}})^T = (a, 2a, 0)$ [56].
- CSD(3): $(m_{\text{atm}}^{\text{D}})^T = (0, e, e)$, $(m_{\text{sol}}^{\text{D}})^T = (a, 3a, a)$ [57].
- CSD(4): $(m_{\text{atm}}^{\text{D}})^T = (0, e, e)$, $(m_{\text{sol}}^{\text{D}})^T = (a, 4a, 2a)$ [57–59].
- CSD(n): $(m_{\text{atm}}^{\text{D}})^T = (0, e, e)$, $(m_{\text{sol}}^{\text{D}})^T = (a, na, (n - 2)a)$ [60].

For the general case of CSD(n) the Dirac mass matrix is then

$$m^{\text{D}} = Y^\nu v_u = \begin{pmatrix} 0 & a \\ e & na \\ e & (n - 2)a \end{pmatrix}. \quad (40)$$

The constrained couplings will be justified later with the help of discrete family symmetry. For now we simply assume these couplings motivated by the desire to obtain an approximately maximal atmospheric angle $\tan \theta_{23} \sim e/f \sim 1$ and trimaximal solar angle $\tan \theta_{12} \sim \sqrt{2}a/(b - c) \sim 1/\sqrt{2}$. Since experiment indicates that the bound $\theta_{13} \lesssim m_2/m_3$ is almost saturated, these schemes require certain phase choices $\arg(a/e)$ in order to achieve the desired reactor angle, leading to predictions for the \mathcal{CP} -violating phase δ_{CP} , discussed below.

In a CSD(n) framework [60], the low energy effective Majorana neutrino mass matrix in equation (34) in the two RH neutrino case may be written as

$$m_{(n)}^\nu = m_a \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} + m_b e^{i\eta} \begin{pmatrix} 1 & n & n-2 \\ n & n^2 & n(n-2) \\ n-2 & n(n-2) & (n-2)^2 \end{pmatrix}, \quad (41)$$

where η is the only physically important phase, which depends on the relative phase between the first and second column of the Dirac mass matrix, $\arg(a/e)$. By comparing equations (38) and (41) for $n = 1$ we identify $m_a = \frac{e^2}{M_{\text{atm}}}$ and $m_b = \frac{a^2}{M_{\text{sol}}}$, which hold for any value of n . This can be thought of as the minimal (two RH neutrino) predictive seesaw model since, for a given n , only three parameters m_a, m_b, η describe the entire neutrino sector (three neutrino masses and the PMNS matrix). CSD(n) with two RH neutrinos always predicts the lightest physical neutrino mass to be zero, $m_1 = 0$. It also immediately predicts TM1 mixing since

$$m_{(n)}^\nu \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}. \quad (42)$$

In other words the column vector $(2, -1, 1)^T$ is an eigenvector of $m_{(n)}^\nu$ with a zero eigenvalue, i.e. it is the first column of the PMNS mixing matrix, corresponding to $m_1 = 0$, which means TM1 mixing in equation (13).

For a given choice of the positive integer n , there are three real input parameters m_a, m_b and η from which two light physical neutrino masses m_2, m_3 , three lepton mixing angles, the \mathcal{CP} -violating phase δ_{CP} and two Majorana phases are derived; a total of nine physical parameters (including the prediction $m_1 = 0$) from three input parameters, i.e. six predictions for each value of n . As the Majorana phases are not known and δ_{CP} is only tentatively constrained by experiment, this leaves five presently measured observables, namely the two neutrino mass squared differences and the three lepton mixing angles, from only three input parameters. Essentially the input parameters m_a and m_b are fixed by the two physical neutrino mass squared differences, which implies that the entire PMNS mixing matrix is determined by only a single parameter, namely the phase η . The resulting best fit predictions for CSD(n) [60] are shown in figure 8 as a function of n . As can be seen, CSD(2) gives a reactor angle which is too small⁵, while CSD($n \geq 5$) gives a reactor angle which is too large. CSD(3) and CSD(4) allow a reactor angle in the desired Goldilocks (experimentally preferred) region $\theta_{13} \sim 8.5^\circ$. This value occurs for special choices of phase $\eta \sim 2\pi/3$ for CSD(3) and $\eta \sim 4\pi/5$ for CSD(4) where positive values of these phases yield negative values of $\delta_{\text{CP}} \sim -90^\circ$ for CSD(3) and $\delta_{\text{CP}} \sim -120^\circ$ for CSD(4), with the mixing angles being independent of the sign of the phase.

5.5. Seesaw mechanism with three RH neutrinos and SD

It is straightforward to extend the previous ideas of SD to the case of three RH neutrinos [6]. The starting assumption is that of the SM supplemented by three RH neutrinos with masses in the classic seesaw range TeV to M_{GUT} in figure 7. Then the light neutrino mass matrix emerges from the seesaw formula in equation (33). The basic idea of SD was given in the framework of the two-right handed neutrino model above, but can now be extended to a third, almost decoupled, RH neutrino, as follows.

Extending the preceding example of two RH neutrinos, it is possible to implement the seesaw mechanism with three RH neutrinos using the SD mechanism [6]. The SD assumption can be made precise as follows. In the basis, $M_{\text{R}} = \text{diag}(M_{\text{atm}}, M_{\text{sol}}, M_{\text{dec}})$ where the Dirac

⁵ This is an example of the general result that two RH neutrino models with two texture zeros and a normal hierarchy are phenomenologically excluded [55].




Figure 8. Best-fit PMNS mixing angles and CP -violating phase with respect to n , for the two right-handed neutrino $CSD(n)$ model. We emphasize that $|\delta_{CP}|$ is a genuine prediction here since we have not used the one sigma hint from experiment as an input constraint. $CSD(3)$ and $CSD(4)$ both yield predictions for mixing angles within the preferred range with differing predictions for the atmospheric angle $\theta_{23} \approx 45^\circ$ and $\theta_{23} \approx 38^\circ$, respectively. Interestingly $CSD(3)$ and $CSD(4)$ lead to predictions for $\delta_{CP} \sim -90^\circ$ and $\delta_{CP} \sim -120^\circ$ with $\eta \sim 2\pi/3$ and $\eta \sim 4\pi/5$ being the best fit values of the input phase.

mass matrix is constructed by extending equation (31) to the three columns $m^D = (m_{atm}^D, m_{sol}^D, m_{dec}^D)$, where $m_{atm}^D = (d, e, f)^T$, $m_{sol}^D = (a, b, c)^T$, etc, the SD assumption in equation (35) generalizes to:

$$\frac{(m_{atm}^D)^\dagger m_{atm}^D}{M_{atm}} \gg \frac{(m_{sol}^D)^\dagger m_{sol}^D}{M_{sol}} \gg \frac{(m_{dec}^D)^\dagger m_{dec}^D}{M_{dec}}. \tag{43}$$

Equation (43) immediately predicts a normal neutrino mass hierarchy:

$$m_3 \gg m_2 \gg m_1 \tag{44}$$

which is the main consequence of SD. The lightest physical neutrino mass m_1 is much smaller than the others since the corresponding RH neutrino ν_R^{dec} being approximately decoupled from the seesaw mechanism. The heaviest physical neutrino has mass m_3 much larger than m_2 since the atmospheric RH neutrino makes the dominant contribution to the seesaw mechanism. The model approximates to the two RH neutrino case (as considered previously) where $m_1 = 0$. The SM with three such RH neutrinos is depicted in figure 9.




Figure 9. The standard model with three right-handed neutrinos defined as $(\nu_R^{\text{atm}}, \nu_R^{\text{sol}}, \nu_R^{\text{dec}})$ which in sequential dominance are mainly responsible for the m_3, m_2, m_1 physical neutrino masses, respectively.

6. Towards a theory of flavour

The flavour problem may be defined as the following collection of puzzles left unanswered by the SM:

- Why are there three families of quarks and leptons?
- Why are all charged fermion masses so hierarchical with down-type quark masses being of the same order as charged lepton masses, and up-type quark masses are much more hierarchical?
- Why are at least two neutrino masses not very hierarchical?
- What is the origin of the neutrino mass?
- Why are neutrino masses so tiny compared to charged fermion masses?
- What is the origin of fermion mixing (both CKM and PMNS matrices)?
- Why are CKM mixing angles smaller than PMNS mixing angles apart from the Cabibbo angle which is of the same order as the reactor angle?
- What is the origin of \mathcal{CP} violation in the quark (and lepton) sectors?

These questions motivate the search for a theory of flavour beyond the SM. In this section we explore possible directions towards a theory of flavour based on the ideas of symmetry, in particular unification together with family symmetry.

6.1. Grand unified theories

One of the exciting things about the discovery of neutrino masses and mixing angles is that this provides additional information about the flavour problem—the problem of understanding the origin of three families of quarks and leptons and their masses and mixing angles. In the framework of the seesaw mechanism, new physics beyond the SM is required to violate lepton number and generate RH neutrino masses which may be as large as the GUT scale. This is also exciting since it implies that the origin of neutrino masses is also related to some GUT symmetry group G_{GUT} , which unifies the fermions within each family. Some possible candidate unified gauge groups are shown in figure 10.

Let us take $G_{\text{GUT}} = SU(5)$ as an example [61]. Each family of quarks (with colour r, b, g) and leptons fits nicely into $SU(5)$ representations of left-handed (L) fermions, $F = \bar{5}$ and




Figure 10. Some possible candidate unified gauge groups.

$T = 10$

$$F = \begin{pmatrix} d_r^c \\ d_b^c \\ d_g^c \\ e^- \\ -\nu_e \end{pmatrix}_L, \quad T = \begin{pmatrix} 0 & u_g^c & -u_b^c & u_r & d_r \\ \cdot & 0 & u_r^c & u_b & d_b \\ \cdot & \cdot & 0 & u_g & d_g \\ \cdot & \cdot & \cdot & 0 & e^c \\ \cdot & \cdot & \cdot & \cdot & 0 \end{pmatrix}_L, \quad (45)$$

where c denotes \mathcal{CP} conjugated fermions. The $SU(5)$ representations $F = \bar{5}$ and $T = 10$ decompose into multiplets of the SM gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ as $F = (d^c, L)$, corresponding to

$$\bar{5} = (\bar{3}, 1, 1/3) \oplus (1, \bar{2}, -1/2), \quad (46)$$

and $T = (u^c, Q, e^c)$, corresponding to

$$10 = (\bar{3}, 1, -2/3) \oplus (3, 2, 1/6) \oplus (1, 1, 1). \quad (47)$$

Thus a complete quark and lepton SM family (Q, u^c, d^c, L, e^c) is accommodated in the $F = \bar{5}$ and $T = 10$ representations, with RH neutrinos, whose \mathcal{CP} conjugates are denoted as ν^c , being singlets of $SU(5)$, $\nu^c = 1$. The Higgs doublets H_u and H_d which break electroweak symmetry in a two Higgs doublet model are contained in the $SU(5)$ multiplets H_5 and $H_{\bar{5}}$.

The Yukawa couplings for one family of quarks and leptons are given by

$$y_u H_{5i} T_{jk} T_{lm} \epsilon^{ijklm} + y_\nu H_{5i} F^i \nu^c + y_d H_{\bar{5}}^i T_{ij} F^j, \quad (48)$$

where ϵ^{ijklm} is the totally antisymmetric tensor of $SU(5)$ with $i, j, k, l = 1, \dots, 5$, which decompose into the SM Yukawa couplings

$$y_u H_u Q u^c + y_\nu H_\nu L \nu^c + y_d (H_d Q d^c + H_d e^c L). \quad (49)$$

Notice that the Yukawa couplings for down quarks and charged leptons are equal at the GUT scale. Generalizing this relation to all three families we find the $SU(5)$ prediction for Yukawa matrices

$$Y_d = Y_e^T, \quad (50)$$

which is successful for the third family, but fails badly for the first and second families. Georgi and Jarlskog [62] suggested to include a higher Higgs representation $H_{\bar{25}}$ which is responsible for the 2-2 entry of the down and charged lepton Yukawa matrices. Dropping $SU(5)$ indices for clarity

$$(Y_d)_{22} H_{\overline{45}} T_2 F_2, \quad (51)$$

decomposes into the second family SM Yukawa couplings

$$(Y_d)_{22} (H_d Q_2 d_2^c - 3H_d e_2^c L_2), \quad (52)$$

where the factor of -3 is an $SU(5)$ Clebsch–Gordan coefficient⁶. Assuming a hierarchical Yukawa matrix with a zero Yukawa element (texture) in the 1-1 position, results in the GUT scale Yukawa relations

$$y_b = y_\tau, \quad y_s = \frac{y_\mu}{3}, \quad y_d = 3y_e, \quad (53)$$

which, after renormalization group running effects are taken into account, are consistent with the low energy masses. The precise viability of these relations has been widely discussed in the light of recent progress in lattice theory which enable more precise values of quark masses to be determined, especially the strange quark mass (see, e.g., [63]). In SUSY theories with low values of the ratio of Higgs VEVs, the relation for the third generation $y_b = y_\tau$ at the GUT scale remains viable, but a viable GUT scale ratio of y_μ/y_s is more accurately achieved within SUSY $SU(5)$ GUTs using a Clebsch factor of $9/2$, as proposed in [64], which is 50% higher than the Georgi–Jarlskog prediction of 3. For other Clebsch relations see [65].

6.2. Discrete family symmetry

As already remarked, it is a remarkable fact that the smallest leptonic mixing angle, the reactor angle, is of a similar magnitude to the largest quark mixing angle, the Cabibbo angle, indeed they may even be equal to each other up to a factor of $\sqrt{2}$. Such relationships may be a hint of a connection between leptonic mixing and quark mixing, where such a connection might be achieved using GUTs [66, 67]. For example, the Georgi–Jarlskog relations discussed above already lead to the left-handed charged lepton mixing angle having a simple relation with the RH down-type quark mixing angle $\theta_{12}^{e_L} \approx \theta_{12}^{d_R}/3$ where the approximation assumes hierarchical Yukawa matrices, with the 1-1 elements being approximately zero. If the upper 2×2 Yukawa matrices are symmetric (as motivated by the successful Gatto–Sartori–Tonin (GST) relation [68] which relates the 12 mixing $\theta_{12}^{d_{L,R}}$ to the down and strange mass by $\theta_{12}^{d_{L,R}} \approx \sqrt{m_d/m_s}$) then we may drop the L, R subscripts and this relation simply becomes $\theta_{12}^e = \theta_{12}^d/3$. In large classes of models, the quark mixing originates predominantly from the down-type quark sector, in which case this relation becomes $\theta_{12}^e = \theta_C/3$. If one starts from TB mixing in the neutrino sector, resulting from some discrete family symmetry, then, using the results in section 4.3 such a charged lepton correction results in a reactor angle in the lepton sector of $\theta_{13} \approx \theta_C/(3\sqrt{2})$. This is a factor of 3 too small to account for the observed reactor angle, but it illustrates how the reactor angle could possibly be related to the Cabibbo angle using GUTs. Indeed it has been suggested that perhaps the charged lepton mixing angle is exactly equal to the Cabibbo angle in some GUT model, leading to $\theta_{13} \approx \theta_C/\sqrt{2}$ [26, 69]. However it is non-trivial to reconcile such large charged lepton mixing with the successful relationships between charged lepton and down-type quark masses, and it seems more likely that charged lepton mixing is not entirely responsible for the reactor angle.

The above discussion provides an additional motivation for combining GUTs with discrete family symmetry in order to account for the reactor angle. Putting these two ideas together we are suggestively led to a framework of new physics beyond the SM based on commuting GUT and family symmetry groups

⁶ In this setup, H_d is the light linear combination of the electroweak doublets contained in $H_{\overline{5}}$ and $H_{\overline{45}}$.




Figure 11. Quark and lepton masses lego plot (true heights need to be scaled by the factors shown) indicating the directions that GUT and family symmetries are acting.




Figure 12. Some popular family symmetries and their relationships.

$$G_{\text{GUT}} \times G_{\text{FAM}}. \tag{54}$$

The spectrum of quark and lepton masses may also provide some motivation for considering a family symmetry as well as a grand unified symmetry, acting in different directions, as illustrated in figure 11. The (scaled) heights of the towers representing the fermion masses, show vast hierarchies which are completely mysterious in the SM. Some popular family symmetries which admit triplet representations are shown in figure 12. The mathematics of these and other groups has recently been reviewed in [11, 14, 15] to which we refer the interested reader for more details.

Here we just mention the family symmetry A_4 as it is the smallest non-Abelian finite group with an irreducible triplet representation. A_4 is the symmetry group of the tetrahedron. There are 12 independent transformations of the tetrahedron and hence 12 group elements as follows:

- Four rotations by 120° clockwise (seen from a vertex) which are T -type.
- Four rotations by 120° anti-clockwise (seen from a vertex) which are T -type.
- Three rotations by 180° which are S -type.
- One unit operator \mathcal{I} .

The generators of the A_4 group, can be written as S and T with $S^2 = T^3 = (ST)^3 = \mathcal{I}$. All 12 group elements can be formed by multiplying together these two generators in all possible ways.

A_4 has four irreducible representations, three singlets $1, 1'$ and $1''$ and one triplet. The products of singlets are:

$$1 \otimes 1 = 1, 1' \otimes 1'' = 1, 1' \otimes 1' = 1'' \otimes 1'' = 1'. \quad (55)$$

Later we shall sometimes work in the real basis of the triplet representation [70],

$$S = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \quad (56)$$

which generate 12 real 3×3 matrix group elements after multiplying these two matrices together in all possible ways [70]. In this basis one has the following Clebsch rules for the multiplication of two triplets, $3 \times 3 = 1 + 1' + 1'' + 3_1 + 3_2$, with

$$\begin{aligned} (ab)_1 &= a_1b_1 + a_2b_2 + a_3b_3; \\ (ab)_{1'} &= a_1b_1 + \omega^2a_2b_2 + \omega a_3b_3; \\ (ab)_{1''} &= a_1b_1 + \omega a_2b_2 + \omega^2a_3b_3; \\ (ab)_{3_1} &= (a_2b_3, a_3b_1, a_1b_2); \\ (ab)_{3_2} &= (a_3b_2, a_1b_3, a_2b_1), \end{aligned} \quad (57)$$

where $a = (a_1, a_2, a_3)$ and $b = (b_1, b_2, b_3)$ are the two triplets and $\omega^3 = 1$.

6.3. Klein symmetry

The starting point for family symmetry models is to consider the Klein symmetry of the neutrino mass matrix. First consider the phase symmetry of the diagonal charged lepton mass matrix M_e ,

$$T^\dagger (M_e^\dagger M_e) T = M_e^\dagger M_e, \quad (58)$$

where $T = \text{diag}(1, \omega, \omega^2)$ and $\omega = e^{2\pi i/n}$. For example for $n = 3$ clearly T generates the group Z_3^T . In any case, the Klein symmetry of the neutrino mass matrix, in this basis, is given by

$$m^\nu = S^T m^\nu S, \quad m^\nu = U^T m^\nu U, \quad (59)$$

where [71]

$$S = U_{\text{PMNS}}^* \text{diag}(+1, -1, -1) U_{\text{PMNS}}^T, \quad (60)$$

$$U = U_{\text{PMNS}}^* \text{diag}(-1, +1, -1) U_{\text{PMNS}}^T, \quad (61)$$

$$SU = U_{\text{PMNS}}^* \text{diag}(-1, -1, +1) U_{\text{PMNS}}^T \quad (62)$$

and

$$\mathcal{K} = \{1, S, U, SU\} \quad (63)$$

is called the Klein symmetry $Z_2^S \times Z_2^U$.

6.4. Direct models

The idea of direct models is that the three generators S, T, U introduced above are embedded into a discrete family symmetry G which is broken by new Higgs fields called ‘flavons’ of two types: ϕ^l whose VEVs preserve T and ϕ^ν whose VEVs preserve S, U . These flavons are segregated such that ϕ^l only appears in the charged lepton sector and ϕ^ν only appears in the

⁷ Note that this is not the same basis as equation (56) since T is diagonal in this basis (but still traceless since $1 + \omega + \omega^2 = 0$).




Figure 13. The direct approach to models of lepton mixing.

neutrino sector as depicted in figure 13, thereby enforcing the symmetries of the mass matrices. Note that the full Klein symmetry $Z_2^S \times Z_2^U$ of the neutrino mass matrix is enforced by symmetry in the direct approach.

Following the measurement of the reactor angle, it has emerged that the only viable direct models are those based on $\Delta(6N^2)$ [72–74]. Unfortunately large N is required in order to achieve the desired reactor angle. Moreover such models generally predict the \mathcal{CP} phase $\delta = 0, \pi$ resulting in the atmospheric sum rule [73],

$$\theta_{23} = 45^\circ \mp \theta_{13}/\sqrt{2}. \tag{64}$$

which follows since the PMNS matrix has the TM2 form shown in equation (13).

6.5. Spontaneous \mathcal{CP} violation

The inclusion of discrete family symmetry and GUTs into a theory of flavour offers the possibility of having spontaneously broken \mathcal{CP} symmetry. The idea is that the high energy theory respects \mathcal{CP} but it becomes spontaneously broken along with the discrete family symmetry and GUT symmetry. As with all types of spontaneous symmetry breaking, this offers the possibility of understanding the origin of \mathcal{CP} violation, and relating \mathcal{CP} violation in the quark and lepton sectors. For example, it is possible that the \mathcal{CP} violating phases in the quark and lepton sectors may be predicted within this kind of approach.

The direct approach can be generalized to the case of a conserved \mathcal{CP} (see [75] and references therein) which is spontaneously broken as shown in figure 14. This approach has been studied for $\Delta(6N^2)$ in [76]. However since we already know that $\delta = 0, \pi$ in this case it only fixes the Majorana phases.

The generalized \mathcal{CP} approach has also been used in the semi-direct approach (defined below) where the phase δ is undetermined without \mathcal{CP} . Here the results are more interesting since for the smaller groups like A_4 and S_4 one generally predicts a discrete choice including $\delta = \pm\pi/2$ [77]. However for larger groups in the series $\Delta(6N^2)$ and $\Delta(3N^2)$, broken in a semi-direct way, the discrete predictions for δ proliferate [78]. Motivated by the the good experimental prospects for measuring leptonic \mathcal{CP} violation, there has been considerable theoretical interest in \mathcal{CP} symmetry in different approaches to family symmetry models [79].




Figure 14. The direct approach including \mathcal{CP} . The idea is that the original high energy theory conserves \mathcal{CP} but \mathcal{CP} is spontaneously broken in the low energy theory. Nevertheless one may define separate \mathcal{CP} symmetries which are preserved in the charged lepton and neutrino sectors, which survive along with preserved subgroups of the original family symmetry in each of these sectors.

Recently an invariant approach \mathcal{CP} symmetry in family symmetry models has been discussed [80]. It is worthwhile to first recap how the invariant approach works for any theory where the Lagrangian is specified. Following [81], to study \mathcal{CP} symmetry in any model one divides a given Lagrangian as follows $\mathcal{L} = \mathcal{L}_{\mathcal{CP}} + \mathcal{L}_{\text{rem}}$ where $\mathcal{L}_{\mathcal{CP}}$ is the part that automatically conserves \mathcal{CP} (like the kinetic terms and gauge interactions) while \mathcal{L}_{rem} includes the \mathcal{CP} violating non-gauge interactions such as the Yukawa couplings. Then one considers the most general \mathcal{CP} transformation that leaves $\mathcal{L}_{\mathcal{CP}}$ invariant and check if invariance under \mathcal{CP} restricts \mathcal{L}_{rem} —only if this is the case can \mathcal{L} violate \mathcal{CP} .

In the presence of a family symmetry G , one may check if a given vacuum leads to spontaneous \mathcal{CP} violation, as follows. Consider a Lagrangian invariant under G and \mathcal{CP} , containing a series of scalars which under \mathcal{CP} transform as $(\mathcal{CP})\phi_i(\mathcal{CP})^{-1} = U_{ij}\phi_j^*$. In order for the vacuum to be \mathcal{CP} invariant, the following relation has to be satisfied: $\langle 0 | \phi_i | 0 \rangle = U_{ij} \langle 0 | \phi_j^* | 0 \rangle$ [82]. The presence of G usually allows for many choices for U . If (and only if) no choice of U exists which satisfies the previous condition, will the vacuum violate \mathcal{CP} , leading to spontaneous \mathcal{CP} violation. In order to prove that no choice of U exists one can construct \mathcal{CP} -odd invariants.

As a brief review of how to derive \mathcal{CP} -odd invariants, consider the Lagrangian of the leptonic part of the SM extended by Majorana neutrino masses. After electroweak breaking at low energies, the most general mass terms are as in equation (3) which we rewrite in matrix form as

$$\mathcal{L}^{\text{lepton}} = -\bar{e}_L m_l e_R - \frac{1}{2} \bar{\nu}_L m_\nu \nu_L^c + \text{h.c.}, \quad (65)$$

due to the $SU(2)_L$ structure, the most general \mathcal{CP} transformation which leaves the leptonic gauge interactions invariant are (ignoring spin)

$$L(x) \rightarrow UL^*(x_P), \quad e_R(x) \rightarrow Ve_R^*(x_P), \quad (66)$$

where $L = (\nu_L, e_L)$ are the left-handed neutrino and charged lepton fields in a weak basis, and x_P are the parity (three-space) inverted coordinates.

In order for $\mathcal{L}^{\text{lepton}}$ to be \mathcal{CP} invariant under equation (66), the terms shown in the equation (65) go into their respective h.c. terms and vice versa:

$$U^\dagger m_\nu U^* = m_\nu^*, \quad U^\dagger m_l V = m_l^*. \quad (67)$$

From equation (67) one can infer how to build combinations of the mass matrices that will result in equations where U and V cancel entirely. The condition for \mathcal{CP} to be conserved is [81]:

$$I_1 \equiv \text{Tr} \left([H_\nu, H_l]^3 \right) = \text{Tr} \left([H_\nu H_l - H_l H_\nu]^3 \right) = 0, \quad (68)$$

where $H_\nu \equiv m_\nu m_\nu^\dagger$ and $H_l \equiv m_l m_l^\dagger$. This equation is a necessary and sufficient condition for Dirac \mathcal{CP} invariance, since it follows from the existence of \mathcal{CP} transformations in equation (67). If the mass matrices are chosen such that $I_1 = 0$ then Dirac type \mathcal{CP} is conserved while if $I_1 \neq 0$ then Dirac type \mathcal{CP} is violated⁸.

As pointed out in [80], once a Lagrangian is specified, which is invariant under a family symmetry G and some \mathcal{CP} transformation, then the consistency relations [75, 77] are automatically satisfied. In order to prove this it is sufficient to consider some generic Lagrangian invariant under a family symmetry transformation, involving some mass term m (Dirac or Majorana), then define $H = mm^\dagger$. Under some G transformation, $\rho(g)$, the mass term remains unchanged implying:

$$\rho(g)^\dagger H \rho(g) = H. \quad (69)$$

Invariance of the Lagrangian under \mathcal{CP} transformation U requires the mass term to swap with its h.c., hence:

$$U^\dagger H U = H^*. \quad (70)$$

Taking the complex conjugate of equation (69) we find

$$\left(\rho(g)^\dagger \right)^* H^* \rho(g)^* = H^* = U^\dagger H U, \quad (71)$$

using equation (70) for the last equality. Using equation (70) again:

$$\left(\rho(g)^\dagger \right)^* U^\dagger H U \rho(g)^* = U^\dagger H U. \quad (72)$$

Hence by using once more equation (69) for a g' , we find

$$U \left(\rho(g)^\dagger \right)^* U^\dagger H U \rho(g)^* U^\dagger = H = \rho(g')^\dagger H \rho(g'). \quad (73)$$

Comparing both sides of equation (73) we see

$$U \rho(g)^* U^\dagger = \rho(g') \quad (74)$$

which is just the consistency relation [75, 77]. In other words, if we consider equations (69) and (70) we do not need to consider the consistency condition separately since it always follows.

The above considerations about whether \mathcal{CP} is conserved or violated apply separately both to the original theory (defined by some high energy Lagrangian, above the scale of symmetry breaking) and to the spontaneously broken theory (defined by some low energy effective Lagrangian, below the scale of symmetry breaking). We are mainly interested in theories which respect \mathcal{CP} at high energy, but where \mathcal{CP} is spontaneously broken, since these

⁸ This is called Dirac type \mathcal{CP} violation since it occurs both when neutrinos are Dirac and Majorana, where the latter case is assumed above. There are two further necessary and sufficient conditions for low energy leptonic \mathcal{CP} invariance which are peculiar to the Majorana sector [83].




Figure 15. The semi-direct approach to models of lepton mixing.

allow for the possibility of being able to predict the amount of \mathcal{CP} violation (e.g. the physical \mathcal{CP} violating phases in some basis).

6.6. Semi-direct models

Taking a less constrained approach to model building one may suppose that we start from only smaller discrete family groups such as S_4 , which leads to either TB or BM mixing at leading order, or A_5 which leads to GR mixing at leading order as shown in figure 15. Then we suppose that at higher order, one or more of the generators S , T , U is broken, which is necessary in this approach since the resulting BM, TB and GR mixing patterns discussed in equations (7)–(9) are excluded. There are two interesting possibilities depicted in figure 15 as follows:

- (1) If the T generator protecting the charged lepton mass matrix is broken, then we can expect charged lepton corrections leading to the solar sum rules discussed in section 4.3.
- (2) If the U generator is broken then this leads to either TM1 or TM2 mixing depending on whether SU or S is preserved, leading to atmospheric sum rules as discussed in section 4.2.

The semi-direct approach was first used in [84, 85] for A_4 where there is no U generator to start with and also S_4 which is broken to A_4 at higher order [85]. It was subsequently generalized to von Dyck groups in [86]. In all cases the reactor angle is not predicted but described by a free parameter. This is a retreat from the original goal of predicting lepton mixing angles using symmetry.

6.7. Indirect models

The final logical possibility is that the family symmetry is completely broken in both the neutrino and charged lepton sectors as shown in figure 16 for the example of the smallest




Figure 16. The indirect approach to models of lepton mixing. The notation is such that ϕ^l refers to $\phi_{e,\mu,\tau}$ and ϕ^ν refers to $\phi_{\text{atm}}, \phi_{\text{sol}},$ and ϕ_{dec} .

family symmetry group that admits triplet representations, namely A_4 . In this approach, we allow the flavons ϕ^l and ϕ^ν to have not only symmetry preserving vacuum alignments, but also new alignments which are orthogonal to them and break the symmetry. In the following, ϕ^l refers to $\phi_{e,\mu,\tau}$ which only enter the charged lepton sector and are responsible for a diagonal charged lepton mass matrix, while ϕ^ν refers to $\phi_{\text{atm}}, \phi_{\text{sol}},$ and ϕ_{dec} which only enter the neutrino sector and are responsible for a Dirac mass matrix of the CSD(n) form.

The advantages of the indirect approach over the previous approaches are:

- (1) It can involve a small family symmetry group such as A_4 (unlike the direct approach which involves large family symmetry groups).
- (2) It is highly predictive since it can yield CSD(n) where the entire PMNS matrix is predicted in terms of one input parameter (unlike the semi-direct approach which only predicts sum rules).

The basic starting point is to consider some small family symmetry such as A_4 which admits triplet representations. The family symmetry is broken by triplet flavons ϕ_i whose vacuum alignment will control the structure of the Yukawa couplings. To illustrate how this works, we sketch a model, where the relevant operators responsible for the Yukawa structure in the neutrino sector are

$$\frac{1}{\Lambda}H_u(L \cdot \phi_{\text{atm}})\nu_{\text{atm}}^c + \frac{1}{\Lambda}H_u(L \cdot \phi_{\text{sol}})\nu_{\text{sol}}^c + \frac{1}{\Lambda}H_u(L \cdot \phi_{\text{dec}})\nu_{\text{dec}}^c, \quad (75)$$

where L is the $SU(2)$ lepton doublet, assumed to transform as a triplet under the family symmetry, while $\nu_{\text{atm}}^c, \nu_{\text{sol}}^c, \nu_{\text{dec}}^c$ are \mathcal{CP} conjugates of the RH neutrinos and H_u is the electroweak scale up-type Higgs field, the latter being family symmetry singlets but distinguished by some additional quantum numbers. In the charged-lepton sector, we consider the operators

$$\frac{1}{\Lambda}H_d(L \cdot \phi_e)e^c + \frac{1}{\Lambda}H_d(L \cdot \phi_\mu)\mu^c + \frac{1}{\Lambda}H_d(L \cdot \phi_\tau)\tau^c, \quad (76)$$

where e^c , μ^c , τ^c are the \mathcal{CP} conjugated RH electron, muon and tau respectively. The RH neutrino Majorana superpotential is typically chosen to give a diagonal mass matrix

$$M_R = \text{diag}(M_{\text{atm}}, M_{\text{sol}}, M_{\text{dec}}). \quad (77)$$

Details of the origin of these operators (e.g. in terms of Majoron fields), the relative values of M_{atm} , M_{sol} , M_{dec} as well as the inclusion of any off-diagonal terms in M_R will all depend on the additional specifications of the model beyond our sketchy model here (we shall consider real models shortly in the next section 7).

The idea is that $\text{CSD}(n)$ discussed in section 5.4 emerges from flavon vacuum alignments in the effective operators involving three flavon fields ϕ_{atm} , ϕ_{sol} , and ϕ_{dec} which are triplets under the flavour symmetry and acquire VEVs that break the family symmetry completely in both the neutrino and charged lepton sectors. The subscripts are chosen by noting that ϕ_{atm} correlates with the atmospheric neutrino mass m_3 , ϕ_{sol} with the solar neutrino mass m_2 , and ϕ_{dec} with the lightest neutrino mass m_1 , which in CSD is light enough that the associated third RH neutrino can, to good approximation, be thought of as decoupled from the theory [6]. $\text{CSD}(n)$ corresponds to the choice of vacuum alignments

$$\langle \phi_{\text{atm}} \rangle = v_{\text{atm}} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}, \quad \langle \phi_{\text{sol}} \rangle = v_{\text{sol}} \begin{pmatrix} 1 \\ \text{in} \\ n-2 \end{pmatrix}, \quad \langle \phi_{\text{dec}} \rangle = v_{\text{dec}} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad (78)$$

where n is a positive integer, and the only phases allowed are in the overall proportionality constants. Such vacuum alignments arise from symmetry preserving alignments together with orthogonality conditions [57, 58], as discussed below.

The starting point for understanding the alignments in equation (78) are the symmetry preserving vacuum alignments of A_4 , namely:

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} \pm 1 \\ \pm 1 \\ \pm 1 \end{pmatrix}, \quad (79)$$

which each preserve some subgroup of A_4 in a basis where the 12 group elements in the triplet representation are real as in equation (56) [70] (i.e. each alignment in equation (79) is an eigenvector of at least one non-trivial group element with eigenvalue +1.) The first alignment in equation (78), which completely breaks the A_4 symmetry, arises from the orthogonality conditions

$$\begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \perp \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad (80)$$

involving two symmetry preserving alignments selected from equation (79). The following symmetry breaking alignment may be obtained which is orthogonal to the alignment in equation (80) and one of the symmetry preserving alignments

$$\begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \perp \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}. \quad (81)$$

The $\text{CSD}(n)$ alignment in equation (78) is orthogonal to the above alignment in equation (81),

$$\begin{pmatrix} 1 \\ n \\ n-2 \end{pmatrix} \perp \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}, \quad (82)$$

where the orthogonality in equation (82) is maintained for any value of n (not necessarily integer). To pin down the value of n and show that it is a particular integer requires a further orthogonality condition.

For example, for $n = 3$, the desired alignment is obtained from the two orthogonality conditions

$$\begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix} \perp \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \quad (83)$$

where the first condition above is a particular case of equation (82) and the second condition involves a new alignment, obtained from two of the symmetry preserving alignments in equation (79),

$$\begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \perp \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}. \quad (84)$$

Using equation (75), the vacuum alignments in equation (78) make up the columns of the Dirac neutrino Yukawa matrix $Y^\nu \propto (\langle \phi_{\text{atm}} \rangle, \langle \phi_{\text{sol}} \rangle, \langle \phi_{\text{dec}} \rangle)$, giving a Dirac mass matrix

$$m^D = Y^\nu v_u = \begin{pmatrix} 0 & a & 0 \\ e & na & 0 \\ e & (n-2)a & c \end{pmatrix}, \quad (85)$$

which is an extension of equation (40) to include a third (decoupled) RH neutrino, where $m^D = (m_{\text{atm}}^D, m_{\text{sol}}^D, m_{\text{dec}}^D)$ and the coefficients e , a , and c are generally complex. The charged-lepton Yukawa matrix is chosen to be diagonal (up to model-dependent corrections, assumed small), corresponding to the existence of three flavons ϕ_e , ϕ_μ and ϕ_τ in the charged-lepton sector which acquire VEVs with alignments [57, 58]

$$\langle \phi_e \rangle = v_e \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \langle \phi_\mu \rangle = v_\mu \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \langle \phi_\tau \rangle = v_\tau \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (86)$$

Given this choice, it is clear that Y^e is diagonal, hence U_{eL} is the identity matrix up to diagonal phase rotations, and that $U_{\text{PMNS}} = U_{\nu L}^\dagger$, i.e. simply the matrix that diagonalizes the neutrino mass matrix, up to charged lepton phase rotations.

7. Realistic theories of flavour

In this section we briefly review two realistic indirect models involving the family symmetry A_4 . The first model involves the PS gauge group with CSD(4), while the second model involves $SU(5)$ GUT with CSD(3). We also discuss leptogenesis in these two models.

7.1. A to Z of flavour with PS

As an example of an ‘indirect’ model, an ‘A to Z of flavour with PS’ based on the PS gauge group has been proposed [87] as sketched in figure 17. The PS symmetry leads to $Y^u = Y^\nu$, where the columns of the Yukawa matrices are determined as in equation (75) with the flavon alignments as in equation (78) for the case $n = 4$. The first column is proportional to the alignment $(0, e, e)$ the second column proportional to the CSD(4) orthogonal alignment, $(a, 4a, 2a)$ and the third column is proportional to the alignment $(0, 0, c)$, where $e \ll a \ll c$ gives the hierarchy $m_u \ll m_c \ll m_t$. This structure predicts a Cabibbo angle $\theta_C \approx 1/4$ in the




Figure 17. A to Z of flavour with Pati–Salam, where $A \equiv A_4$ and $Z \equiv Z_5$. The left-handed families form a triplet of A_4 and are doublets of $SU(2)_L$. The right-handed families are distinguished by Z_5 and are doublets of $SU(2)_R$. The $SU(4)_C$ unifies the quarks and leptons with leptons as the fourth colour, depicted here as white.

diagonal $Y^d \sim Y^e$ basis enforced by the first three alignments in equation (79). It also predicts a normal neutrino mass hierarchy with $\theta_{13} \approx 9^\circ$, $\theta_{23} \approx 45^\circ$ and $\delta \approx 260^\circ$ [87].

The model is based on the PS gauge group, with $A_4 \times Z_5$ (A to Z) family symmetry

$$SU(4)_C \times SU(2)_L \times SU(2)_R \times A_4 \times Z_5. \quad (87)$$

The quarks and leptons are unified in the PS representations as follows

$$F_i = (4, 2, 1)_i = \begin{pmatrix} u & u & u & \nu \\ d & d & d & e \end{pmatrix}_i \rightarrow (Q_i, L_i),$$

$$F_i^c = (\bar{4}, 1, 2)_i = \begin{pmatrix} u^c & u^c & u^c & \nu^c \\ d^c & d^c & d^c & e^c \end{pmatrix}_i \rightarrow (u_i^c, d_i^c, \nu_i^c, e_i^c), \quad (88)$$

where the SM multiplets $Q_i, L_i, u_i^c, d_i^c, \nu_i^c, e_i^c$ resulting from PS breaking are also shown and the subscript i ($= 1, 2, 3$) denotes the family index. The left-handed quarks and leptons form an A_4 triplet F , while the three (\mathcal{CP} conjugated) RH fields F_i^c are A_4 singlets, distinguished by Z_5 charges $\alpha, \alpha^3, 1$, for $i = 1, 2, 3$, respectively. Clearly the PS model cannot be embedded into an $SO(10)$ GUT since different components of the 16-dimensional representation of $SO(10)$ would have to transform differently under $A_4 \times Z_5$, which is impossible. On the other hand, the PS gauge group and A_4 could emerge directly from string theory.

The PS gauge group is broken at the GUT scale to the SM

$$SU(4)_C \times SU(2)_L \times SU(2)_R \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y, \quad (89)$$

by PS Higgs, H^c and \bar{H}^c ,

$$H^c = (\bar{4}, 1, 2) = (u_H^c, d_H^c, \nu_H^c, e_H^c),$$

$$\bar{H}^c = (4, 1, 2) = (\bar{u}_H^c, \bar{d}_H^c, \bar{\nu}_H^c, \bar{e}_H^c). \quad (90)$$

These acquire VEVs in the ‘RH neutrino’ directions, with equal VEVs close to the GUT scale 2×10^{16} GeV,

$$\langle H^c \rangle = \langle \nu_H^c \rangle = \langle \bar{H}^c \rangle = \langle \bar{\nu}_H^c \rangle \sim 2 \times 10^{16} \text{ GeV}, \quad (91)$$

so as to maintain SUSY gauge coupling unification.

Our starting point is to assume that the high energy theory, above the PS breaking scale, conserves \mathcal{CP} symmetry. Under a \mathcal{CP} transformation, the A_4 singlet fields ξ, Σ_u, Σ_d transform into their complex conjugates

$$\xi \rightarrow \xi^*, \quad \Sigma_u \rightarrow \Sigma_u^*, \quad \Sigma_d \rightarrow \Sigma_d^*, \quad (92)$$

where the complex conjugate fields transform in the complex conjugate representations under $A_4 \times Z_5$. For example if $\xi \sim \alpha^4$, under Z_5 , then $\xi^* \sim \alpha$. Similarly if $\Sigma_u \sim 1'$, $\Sigma_d \sim 1''$, under A_4 , then $\Sigma_u^* \sim 1''$, $\Sigma_d^* \sim 1'$. On the other hand, in a particular basis, for A_4 triplets $\phi \sim (\phi_1, \phi_2, \phi_3)$, a consistent definition of \mathcal{CP} symmetry requires the second and third triplet components to swap under \mathcal{CP} ,

$$\phi \rightarrow (\phi_1^*, \phi_3^*, \phi_2^*). \quad (93)$$

With the above definition of \mathcal{CP} , all coupling constants g and explicit masses m are real due to CP conservation and the only source of phases can be the VEVs of fields which break $A_4 \times Z_5$. In the model of interest, all the physically interesting \mathcal{CP} phases will arise from Z_5 breaking.

Let us now consider the A_4 triplet fields ϕ which also carry Z_5 charges. In the full model there are four such triplet fields, or ‘flavons’, denoted as $\phi_1^u, \phi_2^u, \phi_1^d, \phi_2^d$. The idea is that ϕ_i^u are responsible for up-type quark flavour, while ϕ_i^d are responsible for down-type quark flavour.

The structure of the Yukawa matrices depends on the so-called CSD(4) vacuum alignments of these flavons, with the overall phases quantized due to Z_5 ,

$$\langle \phi_1^u \rangle = \frac{V_1^u}{\sqrt{2}} e^{im\pi/5} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}, \quad \langle \phi_2^u \rangle = \frac{V_2^u}{\sqrt{21}} e^{im\pi/5} \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix}, \quad (94)$$

and

$$\langle \phi_1^d \rangle = V_1^d e^{in\pi/5} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \langle \phi_2^d \rangle = V_2^d e^{in\pi/5} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}. \quad (95)$$

We note here that the vacuum alignments in equation (95) and the first alignment in equation (94) are fairly ‘standard’ alignments that are encountered in TB mixing models, while the second alignment in equation (94) is obtained using orthogonality arguments, as discussed in section 6.7. In particular we are using the vacuum alignments in equation (78) for the case $n = 4$, where we identify ϕ_{atm} and ϕ_{sol} with ϕ_1^u and ϕ_2^u . We also use the alignments in equation (86) where we identify ϕ_e and ϕ_μ with ϕ_1^d and ϕ_2^d .

The model will involve Higgs bi-doublets of two kinds, h_u which lead to up-type quark and neutrino Yukawa couplings and h_d which lead to down-type quark and charged lepton Yukawa couplings. In addition a Higgs bidoublet h_3 , which is also an A_4 triplet, is used to give the third family Yukawa couplings.

After the PS and A_4 breaking, most of these Higgs bi-doublets will get high scale masses and will not appear in the low energy spectrum. In fact only two light Higgs doublets will survive down to the TeV scale, namely H_u and H_d . The basic idea is that the light Higgs doublet H_u with hypercharge $Y = +1/2$, which couples to up-type quarks and neutrinos, is a linear combination of components of the Higgs bi-doublets of the kind h_u and h_3 , while the light Higgs doublet H_d with hypercharge $Y = -1/2$, which couples to down-type quarks and charged leptons, is a linear combination of components of Higgs bi-doublets of the kind h_d and h_3 ,

$$h_u, h_3 \rightarrow H_u, \quad h_d, h_3 \rightarrow H_d. \quad (96)$$

The renormalizable Yukawa operators, which respect PS and A_4 symmetries, have the following form, leading to the third family Yukawa couplings shown, using equations (88), (96),

$$F. h_3 F_3^c \rightarrow Q_3 H_u u_3^c + Q_3 H_d d_3^c + L_3 H_u \nu_3^c + L_3 H_d e_3^c, \quad (97)$$

where we have used equations (88), (96). The non-renormalizable operators, which respect PS and A_4 symmetries, have the following form

$$F. \phi_i^u h_u F_i^c \rightarrow Q. \langle \phi_i^u \rangle H_u u_i^c + L. \langle \phi_i^u \rangle H_u \nu_i^c, \quad (98)$$

$$F. \phi_i^d h_d F_i^c \rightarrow Q. \langle \phi_i^d \rangle H_d d_i^c + L. \langle \phi_i^d \rangle H_d e_i^c, \quad (99)$$

where $i = 1$ gives the first column of each Yukawa matrix, while $i = 2$ gives the second column and we have used equations (88), (96). Thus the third family masses are naturally larger since they correspond to renormalizable operators, while the hierarchy between first and second families arises from a hierarchy of flavon VEVs. The lepton operators in equations (98), (99) may be compared to the operators in equations (75), (76).

Inserting the vacuum alignments in equations (94) and (95) into equations (98) and (99), together with the renormalizable third family couplings in equation (97), gives the Yukawa matrices of the form

$$Y^u v_u = Y^\nu v_u = \begin{pmatrix} 0 & a & 0 \\ e & 4a & 0 \\ e & 2a & c \end{pmatrix}, \quad Y^d \sim Y^e \sim \begin{pmatrix} y_d^0 & 0 & 0 \\ 0 & y_s^0 & 0 \\ 0 & 0 & y_b^0 \end{pmatrix}. \quad (100)$$

The PS unification predicts the equality of Yukawa matrices $Y^u = Y^\nu$ and $Y^d \sim Y^e$, while the A_4 vacuum alignment predicts the structure of each Yukawa matrix, essentially identifying the first two columns with the vacuum alignments in equations (94) and (95). With a diagonal RH Majorana mass matrix, Y^ν leads to a successful prediction of the PMNS mixing parameters. Also the Cabibbo angle is given by $\theta_C \approx 1/4$ [59]. Thus equation (100) is a good starting point for a theory of quark and lepton masses and mixing, although the other quark mixing angles and the quark \mathcal{CP} phase are approximately zero. However the above discussion ignores the effect of Clebsch factors which will alter the relationship between elements of Y^d and Y^e , which also include off-diagonal elements responsible for small quark mixing angles in the full model discussed in [87].

In realistic unified models involving an $SO(10)$ -inspired pattern of Dirac and heavy RH neutrino masses, assuming the type I seesaw, the lightest RH neutrino N_1 is too light to yield successful thermal leptogenesis, barring highly fine-tuned solutions, while the second heaviest RH neutrino N_2 is typically in the correct mass range. In [88] we discussed N_2 dominated leptogenesis in the A to Z model, where N_1 is identified with $\nu_1^c = \nu_{\text{atm}}^c$, while N_2 is identified with $\nu_2^c = \nu_{\text{sol}}^c$ and N_3 is identified with $\nu_3^c = \nu_{\text{dec}}^c$, as depicted in figure 17. In the A to Z model the neutrino Dirac mass matrix is equal to the up-type quark mass matrix and has the particular constrained structure in equation (100). We showed that flavour coupling effects in the Boltzmann equations are crucial to the success of such N_2 dominated leptogenesis in this model, by helping to ensure that the flavour asymmetries produced at the N_2 scale survive N_1 washout. The numerical results, supported by analytical insight, showed that in order to achieve successful N_2 leptogenesis, consistent with neutrino phenomenology, requires a ‘flavour swap scenario’ whereby the asymmetry generated in the tauon flavour emerges as a surviving asymmetry dominantly in the muon flavour. However successful leptogenesis

requires a less hierarchical pattern of RH neutrino masses than naively expected, at the expense of some mild fine-tuning involving significant off-diagonal elements of the heavy RH Majorana mass matrix. This leads to large deviations from the CSD(4) predictions, including a NO neutrino spectrum with an atmospheric neutrino mixing angle well into the second octant and a Dirac phase $\delta_{\text{CP}} \approx 20^\circ$, a set of predictions that will be tested soon in neutrino oscillation experiments, as discussed in [88].

7.2. Towards a complete $A_4 \times SU(5)$ SUSY GUT

In this section we describe a fairly complete $A_4 \times SU(5)$ SUSY GUT model which implements CSD(3) with two RH neutrinos [89]. This model has the following virtues:

- It is fully renormalizable at the GUT scale, with an explicit $SU(5)$ breaking sector and a spontaneously broken \mathcal{CP} symmetry.
- The MSSM is reproduced with R-parity emerging from a discrete \mathbb{Z}_4^R .
- Doublet-triplet splitting is achieved through the missing partner mechanism [90].
- A μ term is generated at the correct scale.
- Proton decay is sufficiently suppressed.
- It solves the strong \mathcal{CP} problem through the Nelson–Barr mechanism [91, 92].
- It explains the hierarchies in the quark sector, and successfully fits all of the quark masses, mixing angles and the \mathcal{CP} phase, using only $\mathcal{O}(1)$ parameters.
- It justifies the CSD(3) alignment which accurately predicts the leptonic mixing angles, as well as a normal neutrino mass hierarchy.
- It involves two RH neutrinos with the lighter one dominantly responsible for the atmospheric neutrino mass.
- There is only one physical phase in the model, called η , which is responsible for \mathcal{CP} violation in both leptogenesis and neutrino oscillations.
- A \mathbb{Z}_9 flavour symmetry fixes the phase η to be one of ninth roots of unity [93].

Apart from $A_4 \times SU(5)$ the model also involves the discrete symmetries $\mathbb{Z}_9 \times \mathbb{Z}_6 \times \mathbb{Z}_4^R$. It is renormalizable at the GUT scale, but many effects, including most fermion masses, come from non-renormalizable terms that arise when heavy messenger fields are integrated out. Unwanted or potentially dangerous terms are forbidden by the symmetries and the prescribed messenger sector, including any terms that would generate proton decay or strong \mathcal{CP} violation. Such terms may arise from Planck scale suppressed terms, but prove to be sufficiently small. Due to the completeness of the model, the field content is too big to be listed here, but the superfields relevant for quarks, leptons and Higgs, including flavons, are shown in table 1.

The SM fermions are contained within superfields F and T_i . The MSSM Higgs doublet H_u originates from a combination of H_5 and H_{45} , and H_d from a combination of H_5 and H_{45} . Having the Higgs doublets inside these different representations generates the correct relations between down-type quarks and charged leptons. Doublet-triplet splitting is achieved by the missing partner mechanism [90].

The field ξ which gains a VEV $v_\xi \sim 0.06 M_{\text{GUT}}$ generates a hierarchical fermion mass structure in the up-type quark sector through terms like $v_u T_i T_j (v_\xi/M)^{6-i-j}$, where v_u is the VEV of H_u . It also partially contributes to the mass hierarchy for down-type quarks and charged leptons and provides the mass scales for the RH neutrinos as discussed later. It further produces a highly suppressed μ term $\sim (v_\xi/M)^8 M_{\text{GUT}}$. The resulting symmetric Yukawa matrix for up-type quarks is

Table 1. Superfields containing SM fermions, the Higgses and relevant flavons.

Field	Representation				
	A_4	$SU(5)$	\mathbb{Z}_9	\mathbb{Z}_6	\mathbb{Z}_4^R
F	3	$\bar{5}$	0	0	1
T_1	1	10	5	0	1
T_2	1	10	7	0	1
T_3	1	10	0	0	1
N_{atm}^c	1	1	7	3	1
N_{sol}^c	1	1	8	3	1
Γ	1	1	0	3	1
H_5	1	5	0	0	0
\bar{H}_5	1	$\bar{5}$	2	0	0
H_{45}	1	45	4	0	2
\bar{H}_{45}	1	$\bar{45}$	5	0	0
ξ	1	1	2	0	0
θ_2	1	1	1	4	0
ϕ_{atm}	3	1	3	1	0
ϕ_{sol}	3	1	2	1	0

$$Y_{ij}^u = u_{ij} \left(\frac{\langle \xi \rangle}{M} \right)^{n_{ij}} \sim \begin{pmatrix} \tilde{\xi}^4 & \tilde{\xi}^3 & \tilde{\xi}^2 \\ & \tilde{\xi}^2 & \tilde{\xi} \\ & & 1 \end{pmatrix}, \quad (101)$$

where $\tilde{\xi} = \langle \xi \rangle / M \sim 0.1$. The up-type Yukawa matrix Y^u is highly nondiagonal while the down-type and charged lepton Yukawa matrices $Y^d \sim Y^e$, derived from terms like $F\phi TH$, are nearly diagonal

$$Y_{\text{LR}}^d \sim Y_{\text{RL}}^e \sim \begin{pmatrix} \frac{\langle \xi \rangle v_e}{v_{\Lambda_{24}}^2} & \frac{\langle \xi \rangle v_\mu}{v_{\Lambda_{24}} v_{H_{24}}} & 0 \\ 0 & \frac{v_{H_{24}} v_\mu}{M^2} & 0 \\ 0 & 0 & \frac{v_\tau}{M} \end{pmatrix}, \quad (102)$$

where $v_{e,\mu,\tau}$ are charged lepton flavon VEVs as in equation (86), while $v_{\Lambda_{24}}$ and $v_{H_{24}}$ are the respective VEVs of heavy Higgs Λ_{24} and H_{24} , and we include the subscripts LR to emphasize the role of the off-diagonal term to left-handed mixing from Y^d . The off-diagonal term in Y^e also provides a tiny contribution to left-handed charged lepton mixing $\theta_{12}^e \sim m_e/m_\mu$ which may safely be neglected. It also introduces \mathcal{CP} violation to the CKM matrix via the phase of $\langle \xi \rangle$.

The relevant terms in the superpotential giving neutrino masses are

$$W_\nu = y_1 H_5 F \frac{\phi_{\text{atm}}}{\langle \theta_2 \rangle} N_{\text{atm}}^c + y_2 H_5 F \frac{\phi_{\text{sol}}}{\langle \theta_2 \rangle} N_{\text{sol}}^c + y_3 \frac{\xi^2}{M_\Gamma} N_{\text{atm}}^c N_{\text{atm}}^c + y_4 \xi N_{\text{sol}}^c N_{\text{sol}}^c, \quad (103)$$

where the y_i are dimensionless couplings, expected to be $\mathcal{O}(1)$. The alignment of the flavon vacuum is fixed by the form of the superpotential, with ϕ_{atm} and ϕ_{sol} gaining VEVs according to CSD(3) in equation (78):

Table 2. Best fit parameters and predictions for an $A_4 \times SU(5)$ SUSY GUT with CSD (3) and a fixed phase $\eta = 2\pi/3$, as described in [89]. The spectrum is NO with lightest neutrino mass $m_1 = 0$ and hence the remaining Majorana phase (predicted but not indicated) will be practically impossible to measure.

n	m_a (meV)	m_b (meV)	η (rad)	θ_{12} ($^\circ$)	θ_{13} ($^\circ$)	θ_{23} ($^\circ$)	δ_{CP} ($^\circ$)	m_2 (meV)	m_3 (meV)
3	26.57	2.684	$\frac{2\pi}{3}$	34.3	8.67	45.8	-86.7	8.59	49.8

$$\langle \phi_{\text{atm}} \rangle = v_{\text{atm}} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}, \quad \langle \phi_{\text{sol}} \rangle = v_{\text{sol}} \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix}. \quad (104)$$

This results in a low energy effective Majorana mass matrix of the form in equation (41) for $n = 3$, namely

$$m^\nu = m_a \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} + m_b e^{i\eta} \begin{pmatrix} 1 & 3 & 1 \\ 3 & 9 & 3 \\ 1 & 3 & 1 \end{pmatrix}, \quad (105)$$

where η is the only physically important phase, which depends on the relative phase between the first and second column of the Dirac mass matrix in the flavour basis. The phase η is responsible for \mathcal{CP} violation in both leptogenesis and neutrino oscillations. We identify

$$m_a = \left| \frac{y_1^2 v_u^2 v_{\text{atm}}^2 M_\Gamma}{y_3 \langle \theta_2 \rangle^2 v_\xi^2} \right|, \quad m_b = \left| \frac{y_2^2 v_u^2 v_{\text{sol}}^2}{y_4 \langle \theta_2 \rangle^2 v_\xi} \right|. \quad (106)$$

The Abelian flavour symmetry \mathbb{Z}_9 fixes the phase η to be one of the ninth roots of unity, through a variant of the mechanism used in [93]. The particular choice $\eta = 2\pi/3$ can give the neutrino mixing angles with great accuracy. Furthermore, this phase corresponds to $\delta_{CP} \approx -\pi/2$, consistent with hints from experimental data.

The relevant best fit parameters from our model are given in table 2, along with the model predictions for the leptonic mixing angles and neutrino masses, for $\tan \beta = 5$.

Using the above estimates, in [94] we estimated the baryon asymmetry of the Universe (BAU) for this model resulting from N_1 leptogenesis:

$$Y_B \approx 2.5 \times 10^{-11} \sin \eta \left[\frac{M_1}{10^{10} \text{ GeV}} \right]. \quad (107)$$

Using $\eta = 2\pi/3$ and the observed value of Y_B fixes the lightest RH neutrino mass:

$$M_1 \approx 3.9 \times 10^{10} \text{ GeV}. \quad (108)$$

Note that the phase η controls the BAU via leptogenesis in equation (107). The phase η also controls the entire PMNS matrix, including all the lepton mixing angles as well as all low energy \mathcal{CP} violation. The single phase η is the therefore the source of all \mathcal{CP} violation in this model, including both \mathcal{CP} violation in neutrino oscillations and in leptogenesis, providing a direct link between these two phenomena in this model. We not only have a correlation between the sign of the BAU and the sign of low energy leptonic \mathcal{CP} violation, but we actually know the value of the leptogenesis phase: it is $\eta = 2\pi/3$ which leads to the observed excess of matter over antimatter for $M_1 \approx 4.10^{10}$ GeV together with an observable neutrino oscillation phase $\delta_{CP} \approx -\pi/2$.

8. F-theory origin of SUSY GUTs with discrete family symmetry

F-theory models have attracted considerable interest over the recent years [95]. For example, SUSY GUTs based on $SU(5)$ have been shown to emerge naturally from F-theory. However, in the F-theory context, the $SU(5)$ GUT group is only one part of a larger symmetry. The other parts manifest themselves at low energies as Abelian and/or non-Abelian discrete symmetries, which can be identified as family symmetries, leading to significant constraints in the effective superpotential (for a review see e.g. [96]).

In [97] non-Abelian fluxes were conjectured to give rise to non-Abelian discrete family symmetries in the low energy effective theory. The origin of such a symmetry is the non-Abelian $SU(5)_\perp$ which accompanies $SU(5)_{\text{GUT}}$ at the E_8 point of enhancement. Whether a non-Abelian symmetry survives in the low energy theory will depend on the geometry of the compactified space and the fluxes present. The usual assumption is that the $SU(5)_\perp$ is first broken to a product of $U(1)_\perp$ groups which are then further broken by the action of discrete symmetries associated with the monodromy group. Instead it was conjectured in [97] that *non-Abelian fluxes* can break $SU(5)_\perp$ first to a non-Abelian discrete group S_4 then to a smaller group such as A_4 , D_4 and so on which act as a family symmetry group in the low energy effective theory [98]. This could provide the origin of the $A_4 \times SU(5)$ SUSY GUT model discussed in section 7.2.

9. Conclusion

In conclusion, although the reactor angle has been accurately measured, which rules out simple patterns of lepton mixing such as BM, TB and GR, it is still possible to have simple patterns of lepton mixing with the first or second column of the TB matrix preserved, namely TM1 or TM2, with atmospheric sum rules. It is also possible to maintain BM, TB and GR mixing for *neutrinos* with the reactor angle is due to charged lepton corrections, leading to solar sum rules.

Although adding RH neutrinos is a very simple and minimal thing to do, the number of RH (sterile) neutrinos is undetermined by anomaly cancellation, and their mass spectrum is completely unknown. The classic seesaw mechanism would correspond to having three RH neutrinos with masses in the range $\text{TeV}-M_{\text{GUT}}$.

SD continues to provide an elegant and natural way to understand neutrino mixing angles, with the dominant RH neutrino couplings providing the atmospheric mixing angle, the sub-dominant solar RH neutrino couplings providing the solar mixing angle and the decoupled RH neutrino couplings being irrelevant. The main predictions of SD are a normal neutrino mass hierarchy and the bound on the reactor angle $\theta_{13} \lesssim m_2/m_3$, which indicated the potential largeness of the reactor angle a decade before it was measured. Spurred on by the success of SD, recent versions of CSD have been proposed which explain why the reactor angle bound is saturated. A particular class of such models called CSD(n) give a successful description of the PMNS matrix in terms of a small number of input parameters with good fits of lepton mixing angles for $n = 3, 4$.

Turning to theories of flavour, a very promising approach is the combination of GUT and family symmetry. The large lepton mixing angles suggest some sort of discrete family symmetry at work, although not in the most simple direct way imagined before the reactor angle was measured. In particular the direct symmetry approach in which the symmetries of the mass matrices are directly embedded into the family symmetry, drives us to family

symmetry groups in the $\Delta(6N^2)$ series with large N values necessary in order to explain the reactor angle.

One possibility is that only part of the symmetries of the mass matrices can be found in the family symmetry group, which is called the semi-direct approach. This allows smaller family groups such as S_4 , A_4 , A_5 whose generators S , T , U may only partly survive. For example if T is broken but the Klein symmetry S , U survives in the neutrino sector this would correspond to BM, TB or GR neutrino mixing but with charged lepton corrections, leading to solar sum rules. If T is preserved but U is broken then this corresponds to TM1 or TM2 mixing with atmospheric sum rules.

An attractive alternative is the indirect approach where a small family symmetry such as A_4 is completely broken. In this case new vacuum alignments are possible which can be used to give interesting Yukawa couplings corresponding to different types of CSD(n), leading to highly predictive models. We have given two examples of such models, namely an A to Z of flavour with PS gauge group and a rather complete $A_4 \times SU(5)$ SUSY GUT of flavour, which could originate from F-theory. The A to Z model allows N_2 leptogenesis at the expense of large deviations from the CSD(4) predictions, but still remains very predictive, with NO neutrino masses, an atmospheric angle in the second octant and a Dirac phase $\delta_{CP} \sim 20^\circ$, predictions that will be tested soon by neutrino oscillation experiments.

The $A_4 \times SU(5)$ SUSY GUT is consistent with N_1 leptogenesis arising from the minimal (two-right handed neutrino) predictive seesaw model, and accurately reproduces the CSD(3) predictions with $\eta = 2\pi/3$ being the only source of \mathcal{CP} violation for low energy neutrino physics as well as in the early Universe. This model therefore provides a direct link between matter–antimatter asymmetry and \mathcal{CP} violation in neutrino oscillation experiments. The neutrino masses and PMNS matrix (nine observables) are fixed in this model by three input parameters leading to the predictions in table 2. For example the model predicts a NO spectrum with maximal atmospheric angle $\theta_{23} \sim \pi/4$ and leptonic \mathcal{CP} violation $\delta_{CP} \sim -\pi/2$ in agreement with current experimental hints. Experiment will soon decide if this model is on the right track.

In conclusion, the discovery of neutrino mass and mixing continues to offer tantalizing clues that may help to unravel the mystery of fermion flavour, mass, mixing and \mathcal{CP} violation. The SM is clearly unable to answer such questions, or provide an explanation of the origin of neutrino mass, dark matter or matter–antimatter asymmetry. The history of physics suggests that the answer to these questions will involve symmetry. The largeness of atmospheric and solar mixing, which resemble TB mixing, motivates the use of non-Abelian discrete family symmetries, where such approaches must and can allow Cabibbo sized reactor mixing as we have discussed. The combination of GUTs and discrete family symmetry, together with spontaneous \mathcal{CP} violation, continues to provide promising and testable candidate theories of flavour capable of answering the intriguing puzzles left in the wake of the SM. We have seen that such theories may provide a link between matter–antimatter asymmetry and \mathcal{CP} violation in neutrino oscillation experiments, accompanied by a set of precise predictions which are readily testable by neutrino oscillation experiments.

Acknowledgments

SFK acknowledges partial support from the STFC Consolidated ST/J000396/1 grant and the European Union FP7 ITN-INVISIBLES (Marie Curie Actions, PITN- GA-2011-289442). I would also like to congratulate my colleagues Takaaki Kajita who is the scientist in charge of the University of Tokyo node of ITN-INVISIBLES and Arthur B McDonald (Sudbury

Neutrino Observatory) on the joint award of the 2015 Nobel Prize for Physics ‘for the discovery of neutrino oscillations, which shows that neutrinos have mass’.

References

- [1] Hirsch M, Pas H and Porod W 2013 (<http://scientificamerican.com/article/ghostly-beacons-of-new-physics-extreme-physics-special/>)
- [2] Altarelli G and Feruglio F 1999 *Phys. Rep.* **320** 295
- [3] King S F 2004 *Rep. Prog. Phys.* **67** 107
- [4] Mohapatra R N and Smirnov A Y 2006 *Ann. Rev. Nucl. Part. Sci.* **56** 569
- [5] Strumia A and Vissani F 2006 (arXiv:[hep-ph/0606054](https://arxiv.org/abs/hep-ph/0606054))
- [6] King S F 1998 *Phys. Lett. B* **439** 350
King S F 1999 *Nucl. Phys. B* **562** 57
King S F 2000 *Nucl. Phys. B* **576** 85
- [7] King S F 2002 *J. High Energy Phys.* [JHEP09\(2002\)011](https://arxiv.org/abs/hep-th/0208081)
King S F 2003 *Phys. Rev. D* **67** 113010
- [8] King S F and Ross G G 2001 *Phys. Lett. B* **520** 243
King S F and Ross G G 2003 *Phys. Lett. B* **574** 239
- [9] Altarelli G and Feruglio F 2005 *Nucl. Phys. B* **720** 64
Altarelli G and Feruglio F 2006 *Nucl. Phys. B* **741** 215
de Medeiros Varzielas I, King S F and Ross G G 2007 *Phys. Lett. B* **644** 153
de Medeiros Varzielas I, King S F and Ross G G 2007 *Phys. Lett. B* **648** 201
- [10] Altarelli G and Feruglio F 2010 *Rev. Mod. Phys.* **82** 2701
- [11] Ishimori H, Kobayashi T, Ohki H, Shimizu Y, Okada H and Tanimoto M 2010 *Prog. Theor. Phys. Suppl.* **183** 1
Ishimori H, Kobayashi T, Ohki H, Shimizu Y, Okada H and Tanimoto M 2012 *Lect. Notes Phys.* **858** 1
Ishimori H, Kobayashi T, Ohki H, Shimizu Y, Okada H and Tanimoto M 2013 *Fortsch. Phys.* **61** 441
- [12] Hall L J, Murayama H and Weiner N 2000 *Phys. Rev. Lett.* **84** 2572
de Gouvea A and Murayama H 2015 *Phys. Lett. B* **747** 479
- [13] Hirsch M and King S F 2001 *Phys. Lett. B* **516** 103
Altarelli G, Feruglio F and Masina I 2003 *J. High Energy Phys.* [JHEP01\(2003\)035](https://arxiv.org/abs/hep-th/0303157)
de Gouvea A and Murayama H 2003 *Phys. Lett. B* **573** 94
- [14] King S F and Luhn C 2013 *Rep. Prog. Phys.* **76** 056201
- [15] King S F, Merle A, Morisi S, Shimizu Y and Tanimoto M 2014 *New J. Phys.* **16** 045018
- [16] Fukugita M and Yanagida T 1986 *Phys. Lett. B* **174** 45
Bari P Di 2012 *Contemp. Phys.* **53** 4 315
Blanchet S and Bari P Di 2012 *New J. Phys.* **14** 125012
- [17] Merle A 2013 *Int. J. Mod. Phys. D* **22** 1330020
- [18] Murayama H, Suzuki H, Yanagida T and Yokoyama J 1993 *Phys. Rev. Lett.* **70** 1912
Ellis J R 2004 *Nucl. Phys. B* **137** 190
Antusch S, Bastero-Gil M, King S F and Shafi Q 2005 *Phys. Rev. D* **71** 083519
- [19] Barbieri R, Hall L J, Oliver S J and Strumia A 2005 *Phys. Lett. B* **625** 189
- [20] Verma S 2015 *Adv. High Energy Phys.* **2015** 385968
- [21] Hu B Z *et al* (Daya Bay Collaboration) 2015 *Phys. Rev. Lett.* **115** 111802
- [22] Gonzalez-Garcia M C, Maltoni M and Schwetz T 2014 *J. High Energy Phys.* [JHEP1411\(2014\)052](https://arxiv.org/abs/1404.7573)
- [23] Capozzi F, Fogli G L, Lisi E, Marrone A, Montanino D and Palazzo A 2014 *Phys. Rev. D* **89** 093018
- [24] Forero D V, Tortola M and Valle J W F 2014 *Phys. Rev. D* **90** 093006
- [25] For a non-exhaustive list of neutrino experiments see https://en.wikipedia.org/wiki/List_of_neutrino_experiments
- [26] King S F 2012 *Phys. Lett. B* **718** 136
- [27] Minkowski P 1977 *Phys. Lett. B* **67** 421
Gell-Mann M, Ramond P and Slansky R 1979 *Supergravity* (Amsterdam: North-Holland)
Yanagida T 1979 *Proc. Workshop on Unified Theory and Baryon Number of the Universe (KEK, Japan)*

- Glashow S L 1979 *Cargese Lecture*
Mohapatra R N and Senjanovic G 1980 *Phys. Rev. Lett.* **44** 912
- [28] Olive K A *et al* (Particle data group) 2014 *Chin. Phys. C* **38** 090001
- [29] Fukugita M, Tanimoto M and Yanagida T 1998 *Phys. Rev. D* **57** 4429
Barger V D, Pakvasa S, Weiler T J and Whisnant K 1998 *Phys. Lett. B* **437** 107
Davidson S and King S F 1998 *Phys. Lett. B* **445** 191
- [30] Harrison P F, Perkins D H and Scott W G 2002 *Phys. Lett. B* **530** 167
- [31] Datta A, Ling F S and Ramond P 2003 *Nucl. Phys. B* **671** 383
Everett L L and Stuart A J 2009 *Phys. Rev. D* **79** 085005
Feruglio F and Paris A 2011 *J. High Energy Phys.* **JHEP10(2011)03**
- [32] King S F 2008 *Phys. Lett. B* **659** 244
- [33] Pakvasa S, Rodejohann W and Weiler T J 2008 *Phys. Rev. Lett.* **100** 111801
- [34] Ballett P, King S F, Luhn C, Pascoli S and Schmidt M A 2014 *Phys. Rev. D* **89** 016016
- [35] King S F 2005 *J. High Energy Phys.* **JHEP08(2005)105**
- [36] Masina I 2006 *Phys. Lett. B* **633** 134
- [37] Antusch S and King S F 2005 *Phys. Lett. B* **631** 42
Antusch S, Huber P, King S F and Schwetz T 2007 *J. High Energy Phys.* **JHEP04(2007)060**
- [38] Ballett P, King S F, Luhn C, Pascoli S and Schmidt M A 2014 *J. High Energy Phys.* **JHEP1412(2014)122**
- [39] Marzocca D, Petcov S T, Romanino A and Sevilla M C 2013 *J. High Energy Phys.* **JHEP05(2013)073**
Petcov S T 2015 *Nucl. Phys. B* **892** 400
- [40] Girardi I, Petcov S T and Titov A V 2015 *Nucl. Phys. B* **894** 733
- [41] Weinberg S 1980 *Phys. Rev. D* **22** 1694
- [42] Magg M and Wetterich C 1980 *Phys. Lett. B* **94** 61
Schechter J and Valle J W F 1980 *Phys. Rev. D* **22** 2227
Lazarides G, Shafi Q and Wetterich C 1981 *Nucl. Phys. B* **181** 287
Mohapatra R N and Senjanovic G 1981 *Phys. Rev. D* **23** 165
Wetterich C 1981 *Nucl. Phys. B* **187** 343
- [43] Foot R, Lew H, He X G and Joshi G C 1989 *Z. Phys. C* **44** 441
- [44] Morisi S and Valle J W F 2013 *Fortsch. Phys.* **61** 466
- [45] Hall L J and Suzuki M 1984 *Nucl. Phys. B* **231** 419
Grossman Y and Haber H E 1997 *Phys. Rev. Lett.* **78** 3438
Diaz M A, Romao J C and Valle J W F 1998 *Nucl. Phys. B* **524** 23
Dreiner H K 1997 (arXiv:hep-ph/9707435)
Diaz M A 1997 (arXiv:hep-ph/9712213)
Drees M, Pakvasa S, Tata X and ter Veldhuis T 1998 *Phys. Rev. D* **57** 5335
Faessler A, Kovalenko S and Simkovic F 1998 *Phys. Rev. D* **58** 055004
Chun E J 1999 *Phys. Lett. B* **454** 304
Abada A and Losada M 2000 *Nucl. Phys. B* **585** 45
Hirsch M, Diaz M A, Porod W, Romao J C and Valle J W F 2000 *Phys. Rev. D* **62** 113008
Hirsch M, Diaz M A, Porod W, Romao J C and Valle J W F 2002 *Phys. Rev. D* **65** 119901 (erratum)
- De Campos F, Diaz M A, Eboli O J P, Magro M B and Mercadante P G 2002 *Nucl. Phys. B* **623** 47
- Diaz M A, Hirsch M, Porod W, Romao J C and Valle J W F 2003 *Phys. Rev. D* **68** 013009
Diaz M A, Hirsch M, Porod W, Romao J C and Valle J W F 2005 *Phys. Rev. D* **71** 059904 (erratum)
- Hirsch M and Porod W 2003 *Phys. Rev. D* **68** 115007
- Barbier R *et al* 2005 *Phys. Rep.* **420** 1
- de Campos F, Diaz M A, Eboli O J P, Lineros R A, Magro M B and Mercadante P G 2005 *Phys. Rev. D* **71** 055008
- Aristizabal Sierra D, Hirsch M and Porod W 2005 *J. High Energy Phys.* **JHEP09(2005)033**
Chun E J and Park S C 2005 *J. High Energy Phys.* **JHEP01(2005)009**
- Diaz M A, Mora C and Zerwekh A R 2005 *Eur. Phys. J. C* **44** 277
- Dedes A, Rimmer S and Rosiek J 2006 *J. High Energy Phys.* **JHEP08(2006)005**
- Dreiner H K, Luhn C, Murayama H and Thormeier M 2007 *Nucl. Phys. B* **774** 127
- Allanach B C and Kom C H 2008 *J. High Energy Phys.* **JHEP04(2008)081**

- Dreiner H K, Hanussek M and Grab S 2010 *Phys. Rev. D* **82** 055027
Dreiner H K, Hanussek M, Kim J-S and Kom C H 2011 *Phys. Rev. D* **84** 113005
Allanach B C, Kom C H and Hanussek M 2012 *Comput. Phys. Commun.* **183** 785
Bazzocchi F, Morisi S, Peinado E, Valle J W F and Vicente A 2013 *J. High Energy Phys.* **JHEP01(2013)033**
Dreiner H K, Hanussek M and Luhn C 2012 *Phys. Rev. D* **86** 055012
Peinado E and Vicente A 2012 *Phys. Rev. D* **86** 093024
- [46] Zee A 1980 *Phys. Lett. B* **93** 389
Zee A 1980 *Phys. Lett. B* **95** 461 (erratum)
Cheng T P and Li L-F 1980 *Phys. Rev. D* **22** 2860
Petcov S T 1982 *Phys. Lett. B* **115** 401
Babu K S 1988 *Phys. Lett. B* **203** 132
Deshpande N G, Gunion J F, Kayser B and Olness F I 1991 *Phys. Rev. D* **44** 837
Jarlskog C, Matsuda M, Skadhauge S and Tanimoto M 1999 *Phys. Lett. B* **449** 240
Koide Y and Ghosal A 2001 *Phys. Rev. D* **63** 037301
Balaji K R S, Grimus W and Schwetz T 2001 *Phys. Lett. B* **508** 301
Kitabayashi T and Yasue M 2002 *Phys. Lett. B* **524** 308
Kitabayashi T and Yasue M 2002 *Int. J. Mod. Phys. A* **17** 2519
Chun E J, Lee K Y and Park S C 2003 *Phys. Lett. B* **566** 142
He X-G 2004 *Eur. Phys. J. C* **34** 371
Cirigliano V, Kurylov A, Ramsey-Musolf M J and Vogel P 2004 *Phys. Rev. D* **70** 075007
Akeroyd A G and Aoki M 2005 *Phys. Rev. D* **72** 035011
Bouchand R and Merle A 2012 *J. High Energy Phys.* **JHEP07(2012)084**
- [47] Ma E, Natale A and Rashed A 2012 *Int. J. Mod. Phys. A* **27** 1250134
[48] King S F, Merle A and Panizzi L 2014 *J. High Energy Phys.* **JHEP11(2014)124**
[49] Arkani-Hamed N, Dimopoulos S and Dvali G R 1998 *Phys. Lett. B* **429** 263
Dienes K R, Dudas E and Gherghetta T 1999 *Nucl. Phys. B* **557** 25
Arkani-Hamed N, Dimopoulos S, Dvali G R and March-Russell J 2002 *Phys. Rev. D* **65** 024032
Faraggi A E and Pospelov M 1999 *Phys. Lett. B* **458** 237
Dvali G R and Smirnov A Y 1999 *Nucl. Phys. B* **563** 63
Mohapatra R N, Nandi S and Perez-Lorenzana A 1999 *Phys. Lett. B* **466** 115
Barbieri R, Creminelli P and Strumia A 2000 *Nucl. Phys. B* **585** 28
- [50] Mohapatra R N and Valle J W 1986 *Phys. Rev. D* **34** 1642
[51] Faraggi A E and Halyo E 1993 *Phys. Lett. B* **307** 311
Haba N, Hattori C, Matsuda M, Matsuoka T and Mochinaga D 1994 *Phys. Lett. B* **337** 63
Haba N, Hattori C, Matsuda M, Matsuoka T and Mochinaga D 1994 *Prog. Theor. Phys.* **92** 153
Cleaver G, Cvetic M, Espinosa J R, Everett L L and Langacker P 1998 *Phys. Rev. D* **57** 2701
Ellis J R, Leontaris G K, Lola S and Nanopoulos D V 1998 *Phys. Lett. B* **425** 86
Langacker P 1998 *Phys. Rev. D* **58** 093017
Ellis J R, Leontaris G K, Lola S and Nanopoulos D V 1999 *Eur. Phys. J. C* **9** 389
Ibanez L E, Munoz C and Rigolin S 1999 *Nucl. Phys. B* **553** 43
Coriano C and Faraggi A E 2004 *Phys. Lett. B* **581** 99
Kim J E 2004 *Phys. Lett. B* **591** 119
Giedt J, Kane G L, Langacker P and Nelson B D 2005 *Phys. Rev. D* **71** 115013
Antusch S, Eyton-Williams O J and King S F 2005 *J. High Energy Phys.* **JHEP08(2005)103**
Langacker P and Nelson B D 2005 *Phys. Rev. D* **72** 053013
- [52] Bandyopadhyay A *et al* (ISS Physics Working Group Collaboration) 2009 *Rep. Prog. Phys.* **72** 106201
- [53] Drewes M 2015 (arXiv:1502.06891)
[54] Frampton P H, Glashow S L and Yanagida T 2002 *Phys. Lett. B* **548** 119
[55] Harigaya K, Ibe M and Yanagida T T 2012 *Phys. Rev. D* **86** 013002
[56] Antusch S, King S F, Luhn C and Spinrath M 2012 *Nucl. Phys. B* **856** 328
[57] King S F 2013 *J. High Energy Phys.* **JHEP07(2013)137**
[58] King S F 2013 *Phys. Lett. B* **724** 92
[59] King S F 2014 *J. High Energy Phys.* **JHEP01(2014)119**
[60] Bjorkerth F and King S F 2014 (arXiv:1412.6996)
[61] Georgi H and Glashow S L 1974 *Phys. Rev. Lett.* **32** 438
[62] Georgi H and Jarlskog C 1979 *Phys. Lett. B* **86** 297

- [63] Ross G and Serna M 2008 *Phys. Lett. B* **664** 97
- [64] Antusch S and Spinrath M 2009 *Phys. Rev. D* **79** 095004
- [65] Antusch S, King S F and Spinrath M 2014 *Phys. Rev. D* **89** 055027
- [66] Antusch S and Maurer V 2011 *Phys. Rev. D* **84** 117301
- [67] Marzocca D, Petcov S T, Romanino A and Spinrath M 2011 *J. High Energy Phys.* **JHEP11(2011)009**
- [68] Gatto R, Sartori G and Tonin M 1968 *Phys. Lett. B* **28** 128
- [69] Zhang X and Ma B-Q 2012 *Phys. Rev. D* **86** 093002
- [70] Ma E and Rajasekaran G 2001 *Phys. Rev. D* **64** 113012
- [71] King S F and Luhn C 2009 *J. High Energy Phys.* **JHEP10(2009)093**
- [72] Holthausen M, Lim K S and Lindner M 2013 *Phys. Lett. B* **721** 61
- [73] King S F, Neder T and Stuart A J 2013 *Phys. Lett. B* **726** 312
- [74] Fonseca R M and Grimus W 2014 *J. High Energy Phys.* **JHEP09(2014)033**
- [75] Holthausen M, Lindner M and Schmidt M A 2013 *J. High Energy Phys.* **JHEP04(2013)122**
- [76] King S F and Neder T 2014 *Phys. Lett. B* **736** 308
- [77] Feruglio F, Hagedorn C and Ziegler R 2013 *J. High Energy Phys.* **JHEP07(2013)027**
Feruglio F, Hagedorn C and Ziegler R 2014 *Eur. Phys. J. C* **74** 2753
Ding G-J, King S F, Luhn C and Stuart A J 2013 *J. High Energy Phys.* **JHEP05(2013)084**
Ding G-J, King S F and Stuart A J 2013 *J. High Energy Phys.* **JHEP12(2013)006**
- [78] Hagedorn C, Meroni A and Molinaro E 2015 *Nucl. Phys. B* **891** 499
Ding G J, King S F and Neder T 2014 *J. High Energy Phys.* **JHEP12(2014)007**
- [79] Hall L J and Ross G G 2013 *J. High Energy Phys.* **JHEP11(2013)091**
Antusch S, King S F and Spinrath M 2013 *Phys. Rev. D* **87** 096018 9
Antusch S, Holthausen M, Schmidt M A and Spinrath M 2013 *Nucl. Phys. B* **877** 752
de Medeiros Varzielas I and Emmanuel-Costa D 2011 *Phys. Rev. D* **84** 117901
de Medeiros Varzielas I, Emmanuel-Costa D and Leser P 2012 *Phys. Lett. B* **716** 193
de Medeiros Varzielas I 2012 *J. High Energy Phys.* **JHEP08(2012)055**
Bhattacharyya G, de Medeiros Varzielas I and Leser P 2012 *Phys. Rev. Lett.* **109** 241603
de Medeiros Varzielas I and Pidt D 2014 *J. Phys. G: Nucl. Part. Phys.* **41** 025004
de Medeiros Varzielas I and Pidt D 2013 *J. High Energy Phys.* **JHEP11(2013)206**
Chen M C, Fallbacher M, Mahanthappa K T, Ratz M and Trautner A 2014 *Nucl. Phys. B* **883** 267
Chen M C and Mahanthappa K T 2009 *Phys. Lett. B* **681** 444
Ding G J and King S F 2014 *Phys. Rev. D* **89** 093020
Li C C and Ding G J 2015 *J. High Energy Phys.* **JHEP05(2015)100**
Li C C and Ding G J 2015 *J. High Energy Phys.* **JHEP08(2015)017**
Ding G J and Zhou Y L 2014 *J. High Energy Phys.* **JHEP06(2014)023**
Li C C and Ding G J 2014 *Nucl. Phys. B* **881** 206
Luhn C 2013 *Nucl. Phys. B* **875** 80
Girardi I, Meroni A, Petcov S T and Spinrath M 2014 *J. High Energy Phys.* **JHEP02(2014)050**
Di Iura A, Hagedorn C and Meloni D 2015 (arXiv:1503.04140)
Everett L L, Garon T and Stuart A J 2015 *J. High Energy Phys.* **JHEP04(2015)069**
Ferreira P M, Grimus W, Lavoura L and Ludl P O 2012 *J. High Energy Phys.* **JHEP09(2012)128**
- [80] Branco G C, de Medeiros Varzielas I and King S F 2015 *Phys. Rev. D* **92** 036007
Branco G C, de Medeiros Varzielas I and King S F 2015 *Nucl. Phys. B* **899** 14
- [81] Bernabeu J, Branco G C and Gronau M 1986 *Phys. Lett. B* **169** 243
Jarlskog C 1985 *Phys. Rev. Lett.* **55** 1039
- [82] Branco G C, Gerard J M and Grimus W 1984 *Phys. Lett. B* **136** 383
- [83] Branco G C, Lavoura L and Rebelo M N 1986 *Phys. Lett. B* **180** 264
Branco G C, Rebelo M N and Silva-Marcos J I 1999 *Phys. Rev. Lett.* **82** 683
Dreiner H K, Kim J S, Lebedev O and Thormeier M 2007 *Phys. Rev. D* **76** 015006
- [84] Shimizu Y, Tanimoto M and Watanabe A 2011 *Prog. Theor. Phys.* **126** 81
- [85] King S F and Luhn C 2011 *J. High Energy Phys.* **JHEP09(2011)042**
- [86] Hernandez D and Smirnov A Y 2012 *Phys. Rev. D* **86** 053014
Hernandez D and Smirnov A Y 2013 *Phys. Rev. D* **87** 053005
- [87] King S F 2014 *J. High Energy Phys.* **JHEP08(2014)130**
- [88] Di Bari P and King S F 2015 *J. Cosmol. Astropart. Phys.* **JCAP10(2015)008**
- [89] Bjorkerth F, de Anda F J, de Medeiros Varzielas I and King S F 2015 *J. High Energy Phys.* **JHEP06(2015)141**

- [90] Masiero A, Nanopoulos D V, Tamvakis K and Yanagida T 1982 *Phys. Lett. B* **115** 380
Grinstein B 1982 *Nucl. Phys. B* **206** 387
- [91] Nelson A E 1984 *Phys. Lett. B* **136** 387
Nelson A E 1984 *Phys. Lett. B* **143** 165
- [92] Barr S M 1984 *Phys. Rev. Lett.* **53** 329
Barr S M 1984 *Phys. Rev. D* **30** 1805
- [93] Ross G G, Velasco-Sevilla L and Vives O 2004 *Nucl. Phys. B* **692** 50
Antusch S, King S F and Malinsky M 2008 *J. High Energy Phys.* [JHEP06\(2008\)068](#)
Antusch S, King S F, Luhn C and Spinrath M 2011 *Nucl. Phys. B* **850** 477
- [94] Bjorkerth F, de Anda F J, Varzielas I d M and King S F 2015 *J. High Energy Phys.* [JHEP1510\(2015\)104](#)
- [95] Beasley C, Heckman J J and Vafa C 2009 GUTs and exceptional branes in f-theory: II. Experimental predictions *J. High Energy Phys.* [JHEP01\(2009\)059](#)
- [96] Leontaris G K 2011 *PoS CORFU* vol 2011 p 095
- [97] Antoniadis I and Leontaris G K 2013 *Eur. Phys. J. C* **73** 2670
- [98] Karozas A, King S F, Leontaris G K and Meadowcroft A 2014 *J. High Energy Phys.* [JHEP09\(2014\)107](#)
Karozas A, King S F, Leontaris G K and Meadowcroft A K 2015 *J. High Energy Phys.* [JHEP1510\(2015\)041](#)